

THESE TERMS GOVERN YOUR USE OF THIS DOCUMENT

Your use of this Ontario Geological Survey document (the “Content”) is governed by the terms set out on this page (“Terms of Use”). By downloading this Content, you (the “User”) have accepted, and have agreed to be bound by, the Terms of Use.

Content: This Content is offered by the Province of Ontario’s *Ministry of Northern Development and Mines* (MNDM) as a public service, on an “as-is” basis. Recommendations and statements of opinion expressed in the Content are those of the author or authors and are not to be construed as statement of government policy. You are solely responsible for your use of the Content. You should not rely on the Content for legal advice nor as authoritative in your particular circumstances. Users should verify the accuracy and applicability of any Content before acting on it. MNDM does not guarantee, or make any warranty express or implied, that the Content is current, accurate, complete or reliable. MNDM is not responsible for any damage however caused, which results, directly or indirectly, from your use of the Content. MNDM assumes no legal liability or responsibility for the Content whatsoever.

Links to Other Web Sites: This Content may contain links, to Web sites that are not operated by MNDM. Linked Web sites may not be available in French. MNDM neither endorses nor assumes any responsibility for the safety, accuracy or availability of linked Web sites or the information contained on them. The linked Web sites, their operation and content are the responsibility of the person or entity for which they were created or maintained (the “Owner”). Both your use of a linked Web site, and your right to use or reproduce information or materials from a linked Web site, are subject to the terms of use governing that particular Web site. Any comments or inquiries regarding a linked Web site must be directed to its Owner.

Copyright: Canadian and international intellectual property laws protect the Content. Unless otherwise indicated, copyright is held by the Queen’s Printer for Ontario.

It is recommended that reference to the Content be made in the following form: <Author’s last name>, <Initials> <year of publication>. <Content title>; Ontario Geological Survey, <Content publication series and number>, <total number of pages>p.

Use and Reproduction of Content: The Content may be used and reproduced only in accordance with applicable intellectual property laws. *Non-commercial* use of unsubstantial excerpts of the Content is permitted provided that appropriate credit is given and Crown copyright is acknowledged. Any substantial reproduction of the Content or any *commercial* use of all or part of the Content is prohibited without the prior written permission of MNDM. Substantial reproduction includes the reproduction of any illustration or figure, such as, but not limited to graphs, charts and maps. Commercial use includes commercial distribution of the Content, the reproduction of multiple copies of the Content for any purpose whether or not commercial, use of the Content in commercial publications, and the creation of value-added products using the Content.

Contact:

FOR FURTHER INFORMATION ON	PLEASE CONTACT:	BY TELEPHONE:	BY E-MAIL:
The Reproduction of Content	MNDM Publication Services	Local: (705) 670-5691 Toll Free: 1-888-415-9845, ext. 5691 (inside Canada, United States)	Pubsales@ndm.gov.on.ca
The Purchase of MNDM Publications	MNDM Publication Sales	Local: (705) 670-5691 Toll Free: 1-888-415-9845, ext. 5691 (inside Canada, United States)	Pubsales@ndm.gov.on.ca
Crown Copyright	Queen’s Printer	Local: (416) 326-2678 Toll Free: 1-800-668-9938 (inside Canada, United States)	Copyright@gov.on.ca

LES CONDITIONS CI-DESSOUS RÉGISSENT L'UTILISATION DU PRÉSENT DOCUMENT.

Votre utilisation de ce document de la Commission géologique de l'Ontario (le « contenu ») est régie par les conditions décrites sur cette page (« conditions d'utilisation »). En téléchargeant ce contenu, vous (l'« utilisateur ») signifiez que vous avez accepté d'être lié par les présentes conditions d'utilisation.

Contenu : Ce contenu est offert en l'état comme service public par le *ministère du Développement du Nord et des Mines* (MDNM) de la province de l'Ontario. Les recommandations et les opinions exprimées dans le contenu sont celles de l'auteur ou des auteurs et ne doivent pas être interprétées comme des énoncés officiels de politique gouvernementale. Vous êtes entièrement responsable de l'utilisation que vous en faites. Le contenu ne constitue pas une source fiable de conseils juridiques et ne peut en aucun cas faire autorité dans votre situation particulière. Les utilisateurs sont tenus de vérifier l'exactitude et l'applicabilité de tout contenu avant de l'utiliser. Le MDNM n'offre aucune garantie expresse ou implicite relativement à la mise à jour, à l'exactitude, à l'intégralité ou à la fiabilité du contenu. Le MDNM ne peut être tenu responsable de tout dommage, quelle qu'en soit la cause, résultant directement ou indirectement de l'utilisation du contenu. Le MDNM n'assume aucune responsabilité légale de quelque nature que ce soit en ce qui a trait au contenu.

Liens vers d'autres sites Web : Ce contenu peut comporter des liens vers des sites Web qui ne sont pas exploités par le MDNM. Certains de ces sites pourraient ne pas être offerts en français. Le MDNM se dégage de toute responsabilité quant à la sûreté, à l'exactitude ou à la disponibilité des sites Web ainsi reliés ou à l'information qu'ils contiennent. La responsabilité des sites Web ainsi reliés, de leur exploitation et de leur contenu incombe à la personne ou à l'entité pour lesquelles ils ont été créés ou sont entretenus (le « propriétaire »). Votre utilisation de ces sites Web ainsi que votre droit d'utiliser ou de reproduire leur contenu sont assujettis aux conditions d'utilisation propres à chacun de ces sites. Tout commentaire ou toute question concernant l'un de ces sites doivent être adressés au propriétaire du site.

Droits d'auteur : Le contenu est protégé par les lois canadiennes et internationales sur la propriété intellectuelle. Sauf indication contraire, les droits d'auteurs appartiennent à l'Imprimeur de la Reine pour l'Ontario.

Nous recommandons de faire paraître ainsi toute référence au contenu : nom de famille de l'auteur, initiales, année de publication, titre du document, Commission géologique de l'Ontario, série et numéro de publication, nombre de pages.

Utilisation et reproduction du contenu : Le contenu ne peut être utilisé et reproduit qu'en conformité avec les lois sur la propriété intellectuelle applicables. L'utilisation de courts extraits du contenu à des fins *non commerciales* est autorisée, à condition de faire une mention de source appropriée reconnaissant les droits d'auteurs de la Couronne. Toute reproduction importante du contenu ou toute utilisation, en tout ou en partie, du contenu à des fins *commerciales* est interdite sans l'autorisation écrite préalable du MDNM. Une reproduction jugée importante comprend la reproduction de toute illustration ou figure comme les graphiques, les diagrammes, les cartes, etc. L'utilisation commerciale comprend la distribution du contenu à des fins commerciales, la reproduction de copies multiples du contenu à des fins commerciales ou non, l'utilisation du contenu dans des publications commerciales et la création de produits à valeur ajoutée à l'aide du contenu.

Renseignements :

POUR PLUS DE RENSEIGNEMENTS SUR	VEUILLEZ VOUS ADRESSER À :	PAR TÉLÉPHONE :	PAR COURRIEL :
la reproduction du contenu	Services de publication du MDNM	Local : (705) 670-5691 Numéro sans frais : 1 888 415-9845, poste 5691 (au Canada et aux États-Unis)	Pubsales@ndm.gov.on.ca
l'achat des publications du MDNM	Vente de publications du MDNM	Local : (705) 670-5691 Numéro sans frais : 1 888 415-9845, poste 5691 (au Canada et aux États-Unis)	Pubsales@ndm.gov.on.ca
les droits d'auteurs de la Couronne	Imprimeur de la Reine	Local : 416 326-2678 Numéro sans frais : 1 800 668-9938 (au Canada et aux États-Unis)	Copyright@gov.on.ca



Ontario

Ministry of
Northern Development
and Mines

**Ontario Geological Survey
Open File Report 5829**

**Geology of the
Kamiskotia Area**

1992

3268
ISSN 0826-9580
ISBN 0-7729-9870-1

Geology of the Kamiskotia Area

by

C.T. Barrie¹

1 Geologist, Precambrian Geoscience Section, Ontario Geological Survey

Manuscript approved for publication by B.O. Dressler, Acting Section Chief, Precambrian Geoscience Section, Ontario Geological Survey, January 20, 1992.

Critical Reader: A. Fyon

This report is published with the approval of V.G. Milne, Director, Ontario Geological Survey.



Ministry of
Northern Development
and Mines

Ontario

ONTARIO GEOLOGICAL SURVEY

Open File Report 5829

Geology of the Kamiskotia Area

By

C.T. Barrie

1992

Parts of this publication may be quoted if credit is given. It is recommended that reference to this publication be made in the following form:

Barrie, C.T. 1992. Geology of the Kamiskotia area; Ontario Geological Survey, Open File Report 5829, 180p.

Ontario Geological Survey

OPEN FILE REPORT

Open File Reports are made available to the public subject to the following conditions:

This report is unedited. Discrepancies may occur for which the Ontario Geological Survey does not assume liability. Recommendations and statements of opinions expressed are those of the author or authors and are not to be construed as statements of government policy.

This Open File Report is available for viewing at the following locations:

- (1) Mines Library
Ministry of Northern Development and Mines
8th floor, 77 Grenville Street
Toronto, Ontario M7A 1W4
- (2) The office of the Regional or Resident Geologist in whose district the area covered by this report is located.


Copies of this report may be obtained at the user's expense from a commercial printing house. For the address and instructions to order, contact the appropriate Regional or Resident Geologist's office(s) or the Mines Library. Microfiche copies (42x reduction) of this report are available for \$2.00 each plus provincial sales tax at the Mines Library or the Public Information Centre, Ministry of Natural Resources, W-1640, 99 Wellesley Street West, Toronto.

Handwritten notes and sketches may be made from this report. Check with the Mines Library or Regional/Resident Geologist's office whether there is a copy of this report that may be borrowed. A copy of this report is available for Inter-Library loan.

This report is available for viewing at the following Regional or Resident Geologist's offices:

Cobalt - Box 230, Presley St., Cobalt, Ontario P0J 1C0
Timmins - 60 Wilson Ave., Timmins, Ontario P4N 2S7
Sudbury - 2nd Floor, 159 Cedar St., Sudbury Ontario P3E 6A5

The right to reproduce this report is reserved by the Ontario Ministry of Northern Development and Mines. Permission for other reproductions must be obtained in writing from the Director, Ontario Geological Survey.



V.G. Milne, Director
Ontario Geological Survey

Foreword

This report describes the geology of the Kamikotia area located 15 km west of Timmins and is based on a doctoral thesis by the author at the University of Toronto and additional field investigations for the Ontario Geological Survey.

The report puts particular emphasis on geochemistry and economic geology of the Kamikotia Gabbroic and Volcanic Complexes. Four massive sulphide deposits in the volcanic complex have been mined and approximately 7 million tons of ore have been recovered. One significant, mesothermal gold deposit and several prospects are located along the Destor Porcupine Fault Zone in the southeastern part of the area and along a splay of this fault zone.

Exploration efforts should focus on potential massive sulphide deposits in the metavolcanic rocks and on gold in areas where rocks have been subjected to brittle-ductile deformation.

V.G.Milne
Director, Geoscience Branch
Ontario Geological Survey

CONTENTS

	PAGE
Abstract.....	xvii
Introduction.....	1
Purpose.....	1
Location, Accessibility and Physiography.....	2
Previous Work.....	3
Acknowledgements.....	4
Geology.....	6
Regional Geological Setting.....	6
General Geology of the Kamiskotia Area.....	6
Kamiskotia Gabbroic Complex.....	8
Definition.....	8
Name, Historical Background.....	8
Choice of Lithodemic Rank.....	9
Physical Boundaries.....	10
Geology.....	10
Contacts.....	11
Alteration.....	13
Petrography.....	13
Lower Zone.....	13
Middle Zone.....	14
Mixed Magma Outcrops.....	15
Upper Zone.....	15
Granophyre.....	16
Kamiskotia Volcanic Complex.....	17
Definition.....	17
Physical boundaries.....	17
Geology.....	18
Petrography.....	19
Lower Volcanic Suite.....	20
Metasedimentary Rocks.....	20
Granitoid Rocks.....	21
Bristol Township Lamprophyre Suite.....	22
Structure.....	23
Bedding and Layering, and Regional Folding.....	23
Structural Fabric Analysis.....	26
Results.....	26
Interpretation.....	28
Summary.....	31
Geochronology.....	32
Results of U-Pb Geochronology.....	32
Sm-Nd and Rb-Sr Isochron and Regression Ages.....	33
Magmatic and Structural History.....	35
Geochemistry.....	35
Alteration.....	36
KGC Cumulates.....	36
KGC Chilled Rocks.....	39
Granophyric Rocks.....	41

Kamiskotia Volcanic Complex.....	42
Basalts, Evolved Basalts and Andesites.....	42
Rhyolites.....	43
Lower Volcanic Suite.....	44
Granitoid Rocks.....	44
Bristol Township Lamprophyre Suite.....	45
Geochemical Modeling of KGC Magmatic Processes.....	47
Phase Diagrams.....	48
Mass Balance Calculations.....	48
Assimilation - Fractional Crystallization Modeling.....	50
Nd Isotope Signatures.....	54
Petrogenesis of the KGC, and KVC basalts.....	55
Petrogenesis of KVC rhyolites.....	57
Petrogenesis of Bristol Township Lamprophyre Suite.....	61
Economic Geology.....	63
Volcanogenic Cu-Zn Deposits.....	63
Kam-Kotia and Jameland Mines.....	64
Canadian Jamieson Mine.....	66
Genex Mine.....	67
Mesothermal Au Deposits.....	69
Holmer Property.....	69
Au-REE Occurrence: The Croxall Property.....	70
De Santis Property.....	71
Magmatic Ni-Cu Occurrences.....	72
Western Whitesides Township Occurrences.....	72
Bean Lake - Pirsson Lake Occurrences.....	73
Northwest Carscallen Township Occurrences.....	73
Potential for Mineralization.....	73
Volcanogenic Cu-Zn Deposits.....	74
Mesothermal Au Deposits.....	75
Magmatic Ni-Cu Occurrences.....	76
Discussion.....	77
Gabbroic Complex - Granitoid Relationships	
in the Southern Superior Province.....	77
Chemical Comparison with other Synvolcanic	
Mafic Intrusions in the Southern Superior Province.....	78
Comparison with the Skaergaard Intrusion	
and Eastern Iceland Volcanic Fields.....	79
Cogenesis of the KGC and KVC,	
and Significance with Respect to VMS Deposits.....	81
Kamiskotia - Kidd Creek Relationships.....	82
Time - Stratigraphic Correlation	
in the Southern Abitibi Subprovince.....	83
Late Transpression across the	
Southern Superior Province.....	84
Timing of Magmatism in the Southern Superior Province.....	84
Late Thermal Events in the Kamiskotia Area.....	86
Comparison to Modern Tectonic Comparison to Modern Tectonic Settings:	
Synthesis.....	88
References.....	93

LIST OF TABLES

Table 1. Lithologic units for the Kamiskotia area.....	116
Table 2. Summary of U-Pb geochronology.....	119
Table 3. Summary of Rb-Sr and Sm-Nd isochron/regression ages.....	120
Table 4. Geochemistry of Kamiskotia cumulates.....	121
Table 5. Geochemistry of gabbro chill samples and Kamiskotia basalts	126
Table 6. Geochemistry of the Lower volcanic suite.....	127
Table 7. Geochemistry of granitoid rocks.....	129
Table 8. Geochemistry of the Bristol Township lamprophyre suite....	131
Table 9. Ni-Cu-PGE abundances in magmatic sulphide occurrences.....	134
Conversion Table	180

LIST OF FIGURES

Figure 1. Location map.	135
Figure 2. Distribution of outcrops in Kamiskotia area.	136
Figure 3. Example of density of Matachewan dikes in Kamiskotia area.	137
Figure 4. Geology of the Kamiskotia Gabbroic Complex and nearby rocks.	138
Figure 5. Microprobe analyses for selected samples in the Kamiskotia Gabbroic Complex.	139
Figure 6. Foliation map.	140
Figure 7. Foliation trajectory map.	141
Figure 8. Lineation map.	142
Figure 9. Apparent total mineral strain fabric map.	143
Figure 10. Location of U-Pb samples in western Abitibi Subprovince.	144
Figure 11. Summary of the U-Pb ages in the Kamiskotia - Kidd Creek area.	145
Figure 12. Locations of samples and traverses in the Kamiskotia Gabbroic Complex.	146
Figure 13. Petrographic traverse for cumulate rocks.	147
Figure 14. Mg', normative An, TiO ₂ and P ₂ O ₅ versus stratigraphic height.	148
Figure 15. Ni and Sc versus stratigraphic height.	149

Figure 16. Incompatible trace elements and trace element ratios: La, Yb, La/Yb, Zr, 150 Y, Zr/Y; versus stratigraphic height.	
Figure 17. Rare earth element profiles for Kamiskotia Gabbroic Complex rocks and 151 related basalts.	
Figure 18. Geochemistry of Kamiskotia and Kidd Creek rhyolites. 152	
Figure 19. Rare earth element profiles for selected granitoid rocks. 153	
Figure 20. X-Y plots for Bristol Township lamprophyre suite 154	
Figure 21. Rare earth element profiles for Bristol Township lamprophyre suite. 156	
Figure 22. CMAS-type tetrahedron projections for Kamiskotia basalts. 157	
Figure 23. Mass balance calculations using the REE. 158	
Figure 24. X-Y plots for liquid compositions with AFC modeling curves. 159	
Figure 25. Primitive mantle-normalized profiles for Kamiskotia Gabbroic Complex 160 average chill composition.	
Figure 26. Geology of the Kam-Kotia mine. 161	
Figure 27. Geology of the Canadian Jamieson mine property. 162	
Figure 28. Geology of the Genex Mine area. 163	
Figure 29. Geology of the Genex Mine. 164	
Figure 30. Cross-section of A and H zones, Genex mine. 165	
Figure 31. Cross-section of the C zone, Genex mine. 166	
Figure 32. Geology of the Holmer gold property. 167	
Figure 33. Geology of the De Santis gold property. 168	
Figure 34. Geology of the Croxall gold-REE property. 169	
Figure 35. Compilation of U-Pb ages for southern Superior Province by rock type. 170	
Figure 36. Tectonic model for the Kamiskotia area and parts of the southwestern 171 Abitibi Subprovince.	

LIST OF PLATES

Plate 1. Kamiskotia gabbro cumulates.....	173
Plate 2. Kamiskotia gabbro contact zone textures.....	174
Plate 3. Kamiskotia gabbro mixed magma outcrop textures.....	175
Plate 4. Photomicrographs of Kamiskotia intrusive rocks.....	176
Plate 5. Kamiskotia volcanic rock textures.....	177
Plate 6. Deformation textures.....	178
Plate 7. Croxall property ultramafic lamprophyre suite.....	179

GEOLOGICAL MAP

Geology of the Kamiskotia Area, in back pocket
District of Cochrane
Scale 1:50,000

ABSTRACT

This report describes the geology, geochemistry, structural history and economic geology of the Kamiskotia area, with particular emphasis on the Kamiskotia Gabbroic and Volcanic Complexes. The area is located from 15 km to 40 km west of Timmins, Ontario in the western Abitibi Subprovince.

The Kamiskotia Gabbroic Complex (KGC) is centrally located in the Kamiskotia area. The KGC is a large (170 km²) tholeiitic intrusion and is divided into four stratigraphic units. The Lower Zone is composed of adcumulus and mesocumulus peridotite, troctolite, and gabbro-norite, with Mg numbers from 73 to 86. The Middle Zone is composed of mesocumulus gabbro-norites, with Mg numbers from 60 to 74. The Upper Zone is composed of mesocumulus and orthocumulus gabbro-norites and ferroan gabbro-norites, with Mg numbers from 37 to 64. Modal layering is observed locally, in the Lower and Upper Zones. The fourth zone is a quartz-bearing granophyre of intermediate to felsic composition, and is characterized by granophyric textures and mariolitic cavities. It both overlies and is along strike with Upper Zone cumulates. The mafic cumulates have flat REE patterns ($La_N/Yb_N = 0.4 - 2.6$), with chondrite-normalized REE abundances (REE_N) ranging from 0.6 - 2 for olivine-bearing Lower Zone adcumulates, to 10 - 25 for Upper Zone orthocumulates. Europium anomalies are strongly positive for the Lower Zone and diminish upsection to slightly negative for the highest stratigraphic level in the Upper Zone. The granophyres have very high incompatible element contents, with most samples containing 500 to 800 ppm Zr, $REE_N = 100$ to 200; and have nearly flat REE patterns ($La_N/Yb_N = 1.6$ to 2.6) with strong negative Eu anomalies. Chilled rocks have Mg numbers from 54 to 58, $REE_N = 20$ to 30, and $La_N/Yb_N = 0.8$ to 1.2.

stratigraphy; 5) regional sub-horizontal, bulk north-south shortening that formed a non-penetrative east-west flattening fabric; with an element of 6) late, predominantly dextral transpression, that formed high strain zones related to a splay of the western Destor Porcupine Fault Zone (DPFZ) within the Lower volcanic suite; and 7) emplacement of the Bristol Township lamprophyre suite along the DPFZ.

U-Pb ages from related studies help substantiate this chronology and aid in the correlation of Kamiskotia rocks across the southern Abitibi Subprovince. Zircon ages for the KGC and a KVC rhyolite are 2707 ± 2 Ma and 2705 ± 2 Ma, respectively. The KVC rhyolite is 10 Ma younger than the chemically similar, massive sulphide-bearing Kidd Creek rhyolite located 25 km to the northwest. Two of the granitoid intrusions have zircon ages of 2696 ± 2 Ma and 2694 ± 4 Ma. Two zircon fractions from the younger granitoid intrusion exhibit inheritance, with $^{207}\text{Pb}/^{206}\text{Pb}$ minimum ages up to 2926 Ma, indicating the presence of significantly older crust in this region. A minimum age for the formation of the DPFZ in this area is given by a U-Pb garnet - titanite age for a late-tectonic, garnetite dike, part of the Bristol Township lamprophyre suite within the DPFZ, at 2687 ± 3 Ma.

Four volcanogenic Cu + Zn $\pm$ Ag $\pm$ Au deposits in the KVC have been mined, with >7 million tons of ore recovered. One significant, mesothermal gold deposit and several prospects are located along the DPFZ in the southeastern part of the Kamiskotia area, and along a splay of the DPFZ in the Lower volcanic suite. Exposed on one gold prospect are lamprophyre dikes that contain high rare earth element (REE) concentrations (up to 0.43 wt.% RE₂O₃). Several areas in the Lower Zone of the KGC have minor, low grade Ni-Cu concentrations, with very low platinum group element (PGE) contents.

Exploration efforts should focus on the potential of Cu-Zn deposits in the KVC, with particular emphasis on locating and exploring near synvolcanic faults, in areas with strong chloritic alteration, and in the vicinity of incompatible element-enriched

The Kamiskotia Volcanic Complex (KVC) overlies and is intruded by the KGC. The KVC is comprised of a bimodal assemblage, including tholeiitic basalts and subordinate basaltic andesites and andesites; and high silica rhyolites. Basalts occur as massive and pillowed flows, and as pillow breccias, hyaloclastite and hyaloclastitic tuffs, with plagioclase phenocrysts up to 15% locally. They have Mg numbers from 47 to 67, $REE_N = 17$ to 37, and $La_N/Yb_N = 1.4$ to 1.8. Rhyolites are pyroclastic tuffs, agglomerates, welded tuffs and flow-lobes, and are generally quartz- and feldspar-phyric. They generally have 300 to 400 ppm Zr, $REE_N = 70$ to 280, $La_N/Yb_N = 2$ to 3. Basaltic andesites and andesites are similar in texture to the basalts, and are notable for their discordant, dike- and sill-like masses that cut stratigraphy proximal to massive sulphide deposits. They have Mg numbers from 31 to 47, and are unusually enriched in their incompatible element contents, with 180 to 480 ppm Zr, $REE_N = 25$ to 84, $La_N/Yb_N = 1.2$ to 2.8.

Geochemical modelling of the cumulate rocks and basalts using major and trace elements and radiogenic isotope signatures, indicate that the KGC was derived from a primitive parental magma that fractionated at low pressures. Rhyolitic compositions may be derived from basaltic compositions by greater than 90% fractionation of mafic minerals, plagioclase and Fe-Ti oxide. There is no chemical or isotopic evidence for assimilation of any chemically distinct or isotopically enriched crustal material.

Field observations support the following sequence of magmatic and structural events in the Kamiskotia area: 1) formation of the Lower volcanic suite, stratigraphically below the KGC; 2) emplacement of the KGC and KVC; and coeval intrusion of granitoid A in Turnbull Township into the crystallizing KGC; 3) regional, bulk rotation about a sub-horizontal axis of this stratigraphy to form a northeast-facing monocline; possibly synchronous with 4) emplacement of three large granitoid intrusions, superimposing their respective contact strain aureoles or zones on the

basaltic andesite dikes, sills and flows. Gold exploration should concentrate in areas where rocks have been subjected to late tectonic, brittle-ductile deformation, such as the splay of the DPFZ in the Lower volcanic suite, and along the Kamiskotia Highway fault north of the Canadian Jamieson Mine. The potential for significant Ni-Cu-PGE mineralization in the KGC is considered poor for the following reasons: there is little evidence for large-scale contamination by a more siliceous or oxide-rich crust, generally the cause for sulphide immiscibility that leads to magmatic Ni-Cu deposits; sulphide-bearing sulphide-poor, cumulus and pegmatitic rocks from across the intrusion have very low platinum and palladium abundances. The potential for a bulk-tonnage, low grade REE-gold deposit in the Bristol Township lamprophyre suite should be considered, given that the lamprophyre suite is much more extensive than previously recognized.

INTRODUCTION

PURPOSE

This report describes the geology, geochemistry, geochronology, structural history and economic geology of Late Archean rocks in the Kamiskotia area, with emphasis on the Kamiskotia Gabbroic Complex (KGC) and the Kamiskotia Volcanic Complex (KVC). Included within this report are formal definitions for the KGC and the KVC, using the guidelines of the North American Commission on Stratigraphic Nomenclature (1983) (Table 1).

Large mafic-ultramafic intrusions such as the KGC are a main constituent of Precambrian terranes, and they represent an important source for magmatic Ni-Cu-PGE and Cr deposits (Windley 1976; Naldrett 1981). The majority of these intrusions are stratiform and generally coeval with greenstone belt rocks, in contrast to the Bushveld Complex, South Africa and the Great Dyke, Zimbabwe, which are significantly younger than their country rocks. A feature common to many stratiform intrusions is their geologic setting at the boundaries between regional granitoid and greenstone belt terranes. Examples are found in the Murchison and Pietersburg belts of the Kaapvaal craton, South Africa and Swaziland (e.g., Rooiwater igneous complex and other mafic-ultramafic sills of the Pietersburg Sequence, and the Usushwana complex, Pongola Group), in the Rhodesian craton of Zimbabwe (Mashaba and Shabani mafic-ultramafic complexes), and in the Pilbara block of Western Australia (e.g., Munni Munni and Millindina complexes). Numerous examples are found across the Superior Province, with particularly large ones in the Wabigoon Subprovince of Northwestern Ontario (e.g., Mulcahy Lake and related intrusions; Bad Vermilion and Grassy Portage intrusions), and in the Abitibi Subprovince of Ontario and Quebec (e.g., KGC, Montcalm, Doré Lake, and Bell River complexes).

This report contributes to the understanding of the petrogenesis of Archean strata-bound mafic-ultramafic intrusions, particularly those found at granitoid-greenstone boundaries, in the context of greenstone belt tectonic evolution, with particular emphasis on the KGC. For this study, lithologic and structural mapping of the KGC and surrounding granitoid-greenstone rocks provides a foundation for geochemical studies. Precise chronologic constraints on magmatic and tectonic events in the Kamiskotia area are provided by U-Pb zircon, titanite and garnet geochronology. These events are placed within the well-constrained sequence of tectonic events documented across the southern Abitibi Subprovince and southern Superior Province. Supracrustal magmatic processes for the KGC are investigated using detailed geochemical traverses of the cumulate stratigraphy and geochemistry of the KVC. Primitive source characteristics are described for the KGC from recent trace element and Nd isotope geochemistry studies (Barrie and Shirey, in prep.). Additionally, post-crystallization thermal events are documented using internal Nd and Sr isotope systematics.

This report also describes the KVC in terms of its physical volcanology and geochemistry, and considers its close temporal and genetic relationship with volcanic massive sulphide deposits, and the KGC.

Finally, the largest of the copper-zinc, lode gold, rare earth element and nickel-copper-platinum group element mines, deposits and occurrences in the area are described, in light of current metallogenetic concepts. Following these descriptions, suggestions are made for future mineral exploration in the Kamiskotia area.

LOCATION, ACCESSIBILITY AND PHYSIOGRAPHY

The study area is located 15 to 40 km west of Timmins, Ontario. It encompasses the area contained by Longs.81°30' W to 82°00' W and Lats.81°22'30" N to 48°37'30" N, including all of Whitesides, Carscallen, Massey, Turnbull, Cote and Robb townships and parts of Enid,

Fortune, Bristol, Jamieson, Frey and Godfrey townships, in the district of Cochrane (Figure 1). Principal access routes are from Highway 101 in the southeastern part, from the Kamiskotia Highway (Highway 576) in the northeastern part, and from a major, east-west logging road through the central part that intersects Highway 101. Many subsidiary, all-weather logging roads are present throughout the study area, and in Robb Township, numerous trails and winter roads cover the area south and southeast of Kamiskotia Lake.

Generalized outcrop locations and the locations of prominent esker ridges are given in Figure 2. Also given in Figure 2, is the location of ground coverage for this study. Much of the information for the eastern quarter of the map is compiled from previous studies listed below. In Figure 3, the density of early Proterozoic Matachewan dikes is shown, which is representative of their density for approximately half of the field area. Matachewan dikes are not shown on the accompanying 1:50,000 geological map.

PREVIOUS WORK

Townships in the Kamiskotia area that have been mapped at a scale of 1:15,840 by the Ontario Department of Mines: Bristol (Ferguson 1957a), Carscallen (Ferguson 1957b), Whitesides (Leahy 1968), Cote (Bright and Hunt 1973); Robb and Jamieson (presented together at 1:31,680: Middleton 1973a), Godfrey (Hogg 1954) and Turnbull and Godfrey (presented together at 1:31,680: Middleton 1973b). The geology and mineral occurrences of Turnbull and Godfrey townships have been described in an open file report (Middleton 1975), and ground magnetic survey maps have been published for Robb, Jamieson, Godfrey and Turnbull townships (Middleton 1969, 1970, 1971a, 1971b). Middleton also produced a geophysical report for Robb and Jamieson townships (Middleton 1973c) which describes the prospecting and mining activities in these townships in detail. Wolfe produced a report and an accompanying map on the geology and distribution of Ni, Cu and Co in the Kamiskotia Gabbroic Complex (Wolfe 1970, 1971). Pyke (1982) included the geology of the Kamiskotia

Volcanic Complex in his regional synthesis of the Timmins area. A series of high resolution aeromagnetic maps were produced by the Ontario Geological Survey that cover the eastern half of the area (Barlow 1988).

One master's thesis investigated the geochemistry of the Kamiskotia Volcanic Complex (Hart 1984), and another has studied the Genex Cu deposit in Godfrey Township (Legault 1985). This geological report is accompanied by a map at a scale of 1:50,000, a structural and U-Pb geochronologic study (Barrie and Davis 1990) and an Nd-Sr isotopic study (Barrie and Shirey 1989) on the KGC and surrounding Archean rocks, which stem from doctoral thesis work at the University of Toronto (Barrie 1990).

ACKNOWLEDGEMENTS

There are many people who have helped during the course of this project. Those who were particularly helpful in the scientific research are: Tony Naldrett and Mike Gorton of the University of Toronto, Don Davis of the Royal Ontario Museum (ROM); Steve Shirey of the Department of Terrestrial Magnetism, Carnegie Institution of Washington, D.C. (DTM); and Tony Green of Falconbridge, Limited. I also thank Tom Hart and Marc Legault for permission to use geochemistry and figures from their M.Sc. theses.

In the field, I was accompanied by a number of Falconbridge Limited employees: Mike Kerwin, Doug Hurst, Murray Jerome, Jamie Cecchetto, George MacTaggart, Paul Roos, and Kim Wyotiuk. Their assistance and good humor are greatly appreciated. Additionally, Dave Comba, Bob Stewart, Ted Barnett, Scott McLean, Paul Binney, and Phil Day all of Falconbridge Limited were helpful in providing samples or logistical support. Others were helpful by providing access to drill core or exploration properties, including Dr. Mathew Blecha of Teck Explorations Limited, Dr. Stew Fumerton of Chevron Canada Resources, Limited, and Jim Croxall, of Timmins, Ontario. I am grateful to Wil Doherty who provided high quality ICP-MS data for several challenging, REE-enriched samples.

I would like to thank the following corporations and institutions for their support: the Carnegie Institution of Washington, D.C., Falconbridge Limited, the Jack Satterley Laboratory of the ROM, the University of Toronto, and Teck Explorations Limited.

I thank the following geologists of the Ontario Geological Survey for their reviews which have improved this report: M. E. Cherry, J. A. Fyon, S. L. Jackson, and M. Sanborn-Barrie.

GEOLOGY

REGIONAL GEOLOGICAL SETTING

The Kamiskotia area is located in the westernmost part of the Abitibi Subprovince of the Superior Province (Figure 1). Late Archean metavolcanic, metasedimentary and intrusive rocks of the Ontario portion of the Abitibi Subprovince (herein termed the western Abitibi Subprovince) have been subjected to variable deformation and metamorphism.

Metamorphism ranges from sub-greenschist to middle amphibolite facies, with higher metamorphic grade generally proximal to the margins of large felsic granitoid intrusions. The geology and evolution of parts of the western Abitibi Subprovince have been described and interpreted by numerous authors, most notably Dimroth et al. (1982, 1983a, 1983b), Pyke (1982), Corfu et al. (1989), Jackson and Sutcliffe (1990) and Barrie and Davis (1990).

From U-Pb geochronology, Late Archean rocks of the western Abitibi Subprovince range in age from 2747 ± 2 Ma for the Pacaud Group metavolcanic rocks (Mortensen, personal communication, 1989) south of Kirkland Lake, to $2673 \pm 6/-2$ for an albitite dike in the McIntyre Mine of the Porcupine gold camp (Corfu et al. 1989). There is evidence for older crustal rocks in the vicinity, from xenocrystic zircons in granitic rocks and lamprophyres (Barrie and Davis 1990; Corfu et al., in prep.), from detrital zircons in the Pontiac metasedimentary rocks (Gariépy et al. 1984), and, indirectly, from Nd isotopic signatures (Barrie 1990). Late Archean rocks are cut by the Early Proterozoic Matachewan dikes, which have U-Pb zircon and baddeleyite ages of approximately 2450 Ma (Heaman 1989) (Table 2).

GENERAL GEOLOGY OF THE KAMISKOTIA AREA

The KGC is a large, deformed tholeiitic intrusive complex situated centrally in the Kamiskotia area (Figure 4). It is overlain by, and in part gradational with, metavolcanic

rocks of the KVC; including basalt and rhyolite, and with volumetrically minor evolved basalt and andesite, some of which occur as hypabyssal sills. The KVC hosts four volcanogenic massive sulphide deposits (see Economic Geology section). KGC footwall rocks are informally termed the Lower volcanic suite, which is capped by a 2 m thick cherty oxide-sulphide iron formation. In a general sense, this stratigraphic succession is near-vertical and faces to the north and east. Four granitoid masses composed of hornblende $\pm$ biotite tonalite to granite, and locally rimmed with contact intrusive breccia, have intruded the stratigraphy in the Kamiskotia area. These include granitoid A (Turnbull Township tonalite), predominantly of tonalitic composition, which exhibits textures indicative of magma mixing with fine-grained KGC rocks. Granitoid A is interpreted to have intruded the base or the margin of the KGC during KGC crystallization (described below). Granitoid B (Cote Township tonalite) is predominantly tonalitic, and has a well-developed foliation parallel to its margin. Granitoids C and D, to the west and south of the KGC, are composed of several discrete plutons that range in composition from trondhjemite-tonalite to granodiorite-granite. Regional metamorphism up to the lower greenschist facies has affected the stratigraphy, except within 1 to 3 km of the granitoid - greenstone contacts that have well-developed foliations, where the metamorphic grade is up to middle amphibolite facies.

A major ductile deformation zone is present to the east and south of granitoid D, with a splay extending to the north of granitoid D through the Lower volcanic rocks. The deformation zone may represent parts of the westernmost Destor-Porcupine Fault Zone (DPFZ), a major fault boundary that extends hundreds of kilometres across the Abitibi Subprovince. Alkalic magmatic activity, represented by a lamprophyre suite with allikitic affinities, and mesothermal lode gold mineralization, are associated with the fault zone in this region. The north-northwest-trending Matachewan mafic dike suite cuts all stratigraphy, including deformed rocks in the DPFZ.

KAMISKOTIA GABBROIC COMPLEX

DEFINITION

In this report, the Kamiskotia Gabbroic Complex (KGC) is defined in accordance with the North American Stratigraphic Code (1983). The purpose for a formal definition is to provide stability in nomenclature for easy referencing, and for future studies on the KGC, and for studies concerning lithostratigraphic and time-stratigraphic correlation in the western Abitibi Subprovince. In order to define a rock unit, the Code requires a discussion of the name, historical background and unit rank; descriptions of the rock types, physical boundaries, shape, and other regional aspects including possible correlation with other rocks, and the age and petrogenesis. In this section, the name, historical background and unit rank are discussed, and descriptions of the physical boundaries and rock types are given; the other aspects are discussed in detail elsewhere in this report.

Name, Historical Background

The name Kamiskotia is taken from Kamiskotia Lake and the Kamiskotia River, the most prominent bodies of water in the map area. The first reference to this area and the gabbroic rocks was in 1900, in William A. Parks' survey of Niven's Base Line, from Night Hawk Lake to Missanabie Lake (Parks 1900). The following excerpts are from his report:

The Kamiskotaia- Sagaigan River enters the Mattagami on the west side, about six miles below the three sandy portages. It is about 100 feet wide at its mouth but its navigability is interrupted by rapids...On this portage occurs a peculiar hard schistose rock, striking a little south of east. It presents various shades of pink and green, and weathers out with white dots, owing to a decomposed feldspar. This rock runs up a

hill of considerable height (referring to rhyolites of Kamiskotia hill in Jamieson Township)...

Kamiskotaia Lake is a fine body of water of from two to three miles in diameter, and containing several rocky islands. The rock at the head of the river is hard massive green to black rock, resembling diorite, but contains a large amount of quartz. Under the microscope it shows decomposed plagioclase crystals and blebs of quartz, all imbedded in a fine grained matrix, consisting largely of quartz with minute grains of a dark alteration product. It is probably an altered quartz diorite... (altered KGC quartz gabbro).

Kerr reported on his exploration in the Mattagami valley a few years later, and attempted, without success, to change the name of the Kamiskotia River (Kerr 1906):

About four miles below Niven's second base the Mattagami receives a rather considerable tributary from the west, which I have named the Coffey (Kamiskotia River), the present Indian name, Kamiskotaia-Sagaigan, being altogether too cumbrous. We ascended this river and some of its tributaries to Lake Kamiskotaia...

Choice of Lithodemic Unit Rank

The choice of "complex" as the lithodemic unit rank is based on several criteria. The North American Stratigraphic Code (1983) defines "complex" as "an assemblage or mixture of rocks of two or more genetic classes, igneous, metamorphic or sedimentary, with or without highly complicated structure". The mafic cumulates and the granophyres of intermediate to felsic composition are considered to represent two separate genetic classes of igneous rocks, particularly considering the different magmatic processes that were operative during their

formation. (There is geochemical evidence that supports their comagmatic nature; see Geochemistry section). "Complex" has been used for this intrusion consistently in the literature (e.g., Wolfe 1970, 1971; Middleton 1969, 1970, 1973c, 1975; Pyke 1982; Campbell et al. 1981; 1982; 1984; Hart 1984, Legault 1985; Barrie 1990), although there has been variation on the rock name given and the usage of capitalization. Additionally, the Kamiskotia intrusive rocks have many similarities to well-known intrusions that are termed complexes (e.g., Bushveld Complex, South Africa, Stillwater Complex, Montana, U.S.A.; Bell River and Doré Lake Complexes, Quebec).

Physical Boundaries

The KGC occupies 170 km², in northeastern Carscallen, northern Whitesides, eastern and central Massey, northeastern Enid, central Cote, southern Robb, western and northern Turnbull, and western Godfrey Townships (see 1:50,000 map). It is bounded to the south by the Lower volcanic suite, to the west by granitoid C, to the north and east by the KVC; and internally by granitoids A and B, in Turnbull and Cote townships, respectively. Contact relationships with adjacent rock types are described below.

GEOLOGY

Terminology for the rocks of the KGC is from Irvine (1982). The majority of KGC rocks are termed cumulates, defined as "an igneous rock characterized by a framework of touching mineral crystals and grains that evidently were concentrated through fractional crystallization of their parental magmatic liquids." (Irvine 1982). The packing of cumulus crystals is described by the terms adcumulus (<7% intercumulus minerals), mesocumulus (7 to 25% intercumulus minerals), or orthocumulus (25 to 50+ % intercumulus minerals). Cumulus processes are processes related to the deposition or growth of cumulus crystals or

grains, whereas postcumulus processes refer to processes that post-date cumulus processes, and may be either pre- or post-solidus (Irvine 1982).

The KGC is subdivided into four zones on the basis of field and petrographic observations and geochemistry: partly layered, olivine-bearing cumulates of the Lower Zone (LZ) along the southern and western margin (Figure 4); gabbro-norite and anorthositic gabbro-norite cumulates of the Middle Zone (MZ); partly layered, ferroan gabbro-norite, anorthositic gabbro-norite and hornblende gabbro cumulates of the Upper Zone (UZ); and granophyric rocks of intermediate and felsic composition above and along strike with the UZ cumulates. The UZ - granophyre contact is irregular, with stoped blocks of partially hybridized granophyric rock within chloritized, quartz-rich UZ gabbro locally (Hart 1984).

Facing directions within the LZ and the UZ are from cumulates with cross-bedding structures, or from pyroxene-rich to plagioclase-rich gradations within individual cumulus layers (Plates 1a, b). It is possible that a synform with a north-trending axis is present in the southwestern KGC. Granitoids B, C and D represent antiformal structures, and in general facing directions are away from their margins.

Contacts

Contacts with adjacent rocks exhibit a variety of textures. The adjacent rocks display partial melt zones, and brecciated and agmatitic textures. In the KGC, textures related to the presence or migration of volatiles are found within LZ rocks, and within roof pendants and hybridized blocks of roof-rock material in the granophyre.

Along the southern boundary of the KGC, thermally metamorphosed pillow-shaped structures are present within 100 m of the contact with Lower volcanic rocks (Plate 2a). These textures are gradational with a zone 50 to 200 m wide of contorted, migmatites, with local patches up to several square metres of strong epidote - chlorite - actinolite alteration in basaltic and plagioclase-porphyrific basaltic material. The lower contact with KVC rhyolites

has similar migmatitic textures; also agmatitic textures, where partially melted rhyolite was injected into quenched gabbroic material (similar to agmatitic textures in mixed magma outcrops: Plate 3b).

Within the basal KGC, post-solidus or post-cumulus textures reflect the influence of volatiles, apparently derived from partially melted wall rocks. At some localities, volatile-rich fluids have penetrated along dilatant fractures (post-solidus?: Plate 2b); elsewhere fluids apparently migrated along grain boundaries, and terminated in mushroom-like structures of pegmatitic gabbro and haloes of epidote and clay minerals (post-cumulus?: Plate 2c). Drill core that penetrated layered LZ in western Whitesides Township exhibits another unusual texture, similar to the "tennis ball marker horizon" in the eastern Bushveld Complex. The drill core has spherical, medium-grained plagioclase aggregates up to 10 cm in diameter, within a matrix of sub-pegmatitic gabbro-norite (Plate 2d). This phenomenon is interpreted to have resulted from the aggregation of plagioclase in the presence of a "bridging liquid", which coats loose cumulate phases and bonds them into spheres with minimum surface tension at the base of the magma chamber (Lee and Sharpe 1979). The bridging liquid is apparently a siliceous partial melt of footwall rocks (in this case, cherty iron formation, and Lower volcanic suite rocks) that remains immiscible from the resident magma. Such "bridging liquids" are used in the metallurgical industry to separate particulate matter from melts (Lee and Sharpe 1979).

Blocks of felsic roof rock material have been described within the granophyre and parts of the UZ, in Godfrey and Robb townships (Hogg 1954; Middleton 1975; Hart 1984). These blocks have hybridized margins where they are in contact with more mafic intrusive rocks. Blocks of diorite within the UZ in southern Robb Township south of Kamiskotia Lake have textures similar to those found in the mixed magma outcrops in Turnbull Township, where the felsic intrusive rocks are clearly distinguished from the granophyre on the basis of trace element geochemistry. This area is reinterpreted as late injection of tonalitic liquid into the crystallizing UZ rocks (see below).

Alteration

All KGC rocks have been subjected to varying degrees of post-solidus alteration. For example, clinopyroxene is commonly partly or completely replaced by tremolite-actinolite and hornblende, and plagioclase is commonly altered to sericite, chlorite and clay. Generally the UZ cumulates are slightly altered, whereas the MZ and LZ cumulates are moderately altered. Granophyre rocks are moderately to highly altered, with abundant chlorite in the more intermediate compositions, and chlorite + epidote + sericite assemblages in felsic granophyres. Except for rare sheared outcrops, primary textures are readily discerned in outcrop and hand specimen. The following petrological descriptions emphasize the primary cumulus and post-cumulus mineralogy as determined from outcrop, hand specimen, and petrographic observations, supplemented by microprobe wavelength dispersive analyses of minerals (reported in Barrie 1990). Microprobe mineral analyses are presented in Figure 5.

PETROGRAPHY

Lower Zone

The LZ is distinguished from other rocks of the KGC by the sporadic presence of olivine as a cumulus phase, and by whole-rock Mg numbers that are generally > 75 (whole rock Mg number = mol percent MgO/(MgO + FeO), with $\text{FeO} = 0.85 \text{ FeO}_{\text{total}}$). Modal layering and cross-bedding features are observed locally (Plates 1a,b). LZ rocks include peridotite, troctolite, olivine gabbro, magnesian gabbro-norite, and gabbroic anorthosite (Plate 1c) as medium-grained meso- to adcumulate, with clinopyroxene, orthopyroxene, olivine and plagioclase as cumulus phases. Chromite and sulphides occur sporadically in trace amounts in the cumulates (Plate 4a), and the LZ hosts several minor occurrences of low grade, Ni-

Cu-sulphide mineralization. Olivine ranges in composition from Fo₇₇ to Fo₈₁; clinopyroxene Mg numbers (mineral Mg number = mol percent MgO/(MgO + FeO_{total})) range from 77 to 82, and plagioclase ranges from An₇₉ to An₈₉ (Figure 5: samples A-D). Less primitive orthocumulus to mesocumulus gabbroic rocks predominate along the southern contact (sample F: Figure 5). A large outcrop (300 m by 300 m) of altered peridotite occurs stands in high relief in northwest Carscallen and northeast Whitesides townships. The majority of the outcrop is comprised of talc, serpentine, chlorite and magnetite; relict cumulus olivine is observed locally. The margins of the outcrop are schistose, and contact relationships with layered troctolites and olivine gabbros to the south, and gabbro-norites to the north, are not discernable. The peridotite outcrop is believed to be concordant with surrounding KGC rocks.

Middle Zone

The MZ is comprised of gabbro, gabbro-norite, anorthositic gabbro, and gabbroic anorthosite as massive, medium- and coarse-grained mesocumulate to adcumulate, with clinopyroxene, orthopyroxene and plagioclase as the cumulus phases. Ti-magnetite, ilmenite and sulphides occur as accessory, intercumulus phases. Sub-pegmatitic and pegmatitic textures are common. The majority of MZ rocks are cumulates that do not exhibit modal layering; however, mafic enclaves of basaltic or fine-grained melagabbroic material are found within anorthositic gabbro in northern Whitesides Township (Plate 1d), and mixed magma textures with granitoid A tonalite are present in western and northern Turnbull Township (described below). For cumulus MZ rocks, plagioclase ranges in composition from An₄₆ to An₇₀ and clinopyroxene Mg numbers range from 51 to 85 (samples G-I, Figure 5).

Mixed Magma Outcrops

A wide variety of mixed magma textures between MZ and UZ gabbroic and tonalitic - granodioritic rocks occur over a 1 km by 5 km area, in western Turnbull Township (Figure 4). Agmatitic rocks with >60% felsic intrusive material are found near and within the bounds of the granitoid A (Plates 3a, b). Well-preserved mafic pillow-shaped structures up to 1 m by 1 m, with quenched rims that contain fine-grained radiating, acicular clinopyroxene and plagioclase aggregates, are found within coarse-grained, felsic intrusive rocks (Plates 3c-f, 4a). The pillow-shaped structures have aphyric margins that grade into uniform, medium-grained cores. In Robb Township south of Kamiskotia Lake, tonalitic and gabbroic material are intercalated as alternating sill-like structures from 1 to 10 m thick. Here the grain size in the gabbro becomes increasingly fine-grained toward the tonalitic sills, and the gabbro is highly chloritic. These textures are similar to those found in well-documented mixed magma zones in mafic intrusions, such as the Tigalak and Newark Island layered intrusions, Labrador (Weibe and Wild 1983; Weibe 1987).

Upper Zone

The UZ is comprised of ferroan gabbro, quartz gabbro, hornblende gabbro and hornblendite, as massive and layered, medium- and coarse-grained meso- and orthocumulate. Plagioclase, clinopyroxene, orthopyroxene, pigeonite, and locally Ti-magnetite are cumulus phases; apatite, and biotite occur as intercumulus phases (Plate 4b). Wolfe (1970) reported hornblende gabbro and hornblendite outcrops in the southeastern part of the UZ. Plagioclase commonly comprises greater than 70% of the mode, and several metre-thick anorthosite layers (>90% plagioclase) are observed locally. Inverted pigeonite with clinopyroxene exsolution is common. Generally UZ rocks have a seriate texture and show no

petrographic evidence for post-cumulus overgrowths. Plagioclase compositions from petrographic observations and microprobe analyses range from An₄₆ to An₆₀; Mg numbers for clinopyroxene and orthopyroxene range from 59 to 70, and 57 to 75, respectively (samples J-L: Figure 5). Cumulus magnetite contains up to 2.5 wt.% V₂O₃ (Barrie 1990).

Granophyre

A wide variety of rocks are included in the granophyre zone, including diorite, quartz diorite, quartz monzodiorite, granodiorite, tonalite, quartz monzonite quartz-feldspar porphyry, and granite. These rocks are considered as one unit because the majority of them exhibit granophyric textures (Plate 4c), and because of their similar high trace element abundances (see geochemistry section). Hogg (1954) has provided the most detailed mapping and petrologic descriptions of the granophyre zone. His units: "Intermediate Intrusives, Granite, Granophyre, and Porphyritic Intrusives" are included in the granophyre zone of this report (Figure 5). Contacts of granophyre zone rocks have irregular patterns. This is in part due to felsic granophyres that grade into KVC rocks: Hogg (1954) noted that some felsic granophyre rocks with layered spherulitic structures are gradational into rhyolites in northeastern Godfrey Township. Granophyres of intermediate composition (diorite, quartz diorite), located in northwestern Godfrey Township, are composed of acicular hornblende and actinolite, chlorite, trellis-textured leucoxene after Ti-magnetite, and quartz and feldspar. Granophyres of felsic composition (quartz monzodiorite, granodiorite, tonalite, quartz monzonite quartz-feldspar porphyry, and granite) in northwest Godfrey and southeast Jamieson townships have fine-grained margins and granophyric and locally mariolitic textures in the interior.

KAMISKOTIA VOLCANIC COMPLEX

DEFINITION

The Kamiskotia Volcanic Complex (KVC) is defined here, in accordance with the North American Stratigraphic Code (1983). The historical background, and reasons for the choice of name and lithodemic unit rank are essentially the same as those mentioned above for the KGC. Regional aspects, possible correlation with other rocks, and the age and petrogenesis of these metavolcanic rocks are discussed in detail elsewhere in this report. In this section, the descriptions of the physical boundaries and rock types are given.

Physical Boundaries

The KVC occupies 210 km², in Carscallen, Bristol, a eastern Turnbull, western Godfrey, southwestern Jamieson, southern Robb, and central Cote townships (see accompanying 1:50,000 map, also Figure 5). It is bounded to the south by the Lower volcanic suite; to the west by the KGC and granitoid A in Turnbull Township; to the north and east by a line that is parallel to stratigraphy and extends from a point 2 km north of the Kam-Kotia Mine to a point 2 km east of the Genex Mine, extending into western Loveland Township to the north and toward central Bristol Township. This line represents a demarcation between metavolcanic rocks with few ground or airborne geophysical conductors to the west, and metavolcanic - metasedimentary rocks with numerous conductors to the east (Barlow 1988). The boundaries are well-defined in outcrop where they are represented by contacts with the KGC, but are less well-constrained in areas of poor exposure to the south, east and north. These boundaries are slightly more restrictive than those proposed by Middleton (1975). It is noted that the Reid Township rhyolite, with similar U-Pb age (see

Geochronology section) and geochemistry to KVC rhyolites, may be a distal airfall deposit related to KVC magmatism.

GEOLOGY

The KVC is divided into two units: felsic rocks, composed of massive or poorly bedded pyroclastic deposits with lesser block and ash flow material and flow-lobe complexes; and mafic rocks, comprised of massive and pillowed flows, pillow breccias and hyaloclastite tuffs and breccias. These units are observed to be intercalated on a scale of tens of metres in several locations with excellent exposure: at Mt. Jamieson in Jamieson Township, and the Shell outcrops 3 km to the northwest, and the Canadian Jamieson and Genex Mine properties in Godfrey Township. This implies that some mafic and felsic units are coeval.

Felsic rocks predominate in Carscallen and southern Godfrey townships. A thick sequence of pyroclastic tuffs and flows are present in Carscallen Township, with several individual units having apparent thicknesses of hundreds of metres, divided by agglomeratic units tens of metres in thickness. In northern Bristol Township, where exposures are poor, descriptions of rocks in drill core indicate a significant component of mafic metavolcanic rocks and lesser metasedimentary rocks. Some of the tuffs have fiamme-like textures that may represent primary welding. In Godfrey and eastern Turnbull townships, several layers of felsic agglomerates are present (Plate 5a), within massive, crystal-ash tuffs. A welded quartz-crystal lapilli tuff in central southern Godfrey Township has unusual andesitic fragments with quenched rims (Comba et al. 1986), suggesting that the andesitic material quenched during a predominantly felsic pyroclastic eruption (Plate 5b). This provides additional evidence for the coeval eruption and deposition of magmas of intermediate and felsic composition. In Jamieson Township, rhyolite occurs in lobe-shaped flows (with restricted widths and thicknesses on a scale of tens to hundreds of metres). These constitute most of the outcrop ridges that extend from Mt. Jamieson, 3 km to the northeast (Comba et al. 1986). The lobe-

shaped rhyolite flows are interlayered with mafic flows. They are flow-banded, locally spherulitic, and occur locally as hyaloclastite, with large amygdules or lithophysa up to 3 cm in diameter. Spherulites may comprise up to 50% of a rhyolite flow on a given outcrop (Plate 5c).

KVC mafic rocks predominate in Robb Township, where a nearly uninterrupted, 2 km stratigraphic thickness is present west of the Kam-Kotia Mine. In this area, massive and pillowed basalts have relatively few amygdules (< 10%). Amygdules are larger and more abundant along strike and upsection to the southeast in Godfrey Township. Thick, massive flows and sills are more common along strike to the west, whereas thinner units, including pillowed basalts and hyaloclastites are more common toward the southeast (Middleton 1973c). Mafic metavolcanic rocks on the well-exposed Canadian Jamieson Mine property are: massive flows, pillowed flows (Plate 4d), some with amygdules up to 0.5 mm (comprising up to 2% of the rock), and pillow breccia and hyaloclastite. Mafic rocks at the Genex Mine property are massive and pillowed flows, and pillow breccias.

PETROGRAPHY

Primary mineralogy in KVC rocks is in general difficult to discern due to alteration. Less altered felsic tuffs and flows generally contain 5 to 15% quartz phenocrysts; feldspar phenocrysts are less common and range up to 15%. The matrix is aphanitic, and comprised of sericite, chlorite, quartz, potassium-feldspar and clay minerals; trace phases may include zircon, apatite, titanite, rutile, pyrite, and leucoxene after Ti-magnetite. In more altered felsic rocks, these minerals and epidote and carbonate may be present. Spherules are composed of fine-grained, radiating intergrowths of quartz and feldspar, with the feldspar commonly altered to sericite and clay minerals.

Mafic rocks contain plagioclase phenocrysts up to 20% locally, although > 10% plagioclase phenocryst contents are rare. Primary mineralogy is particularly difficult to

determine in mafic rocks of the KVC. Hart (1984) reported that sericite and epidote alteration of plagioclase does not allow for optical determination of their compositions. The matrix is generally chlorite, altered feldspar, amphibole, carbonate, epidote, possibly pyroxene and quartz; trace phases may include titanite, leucoxene after Ti-magnetite, and sulphide.

LOWER VOLCANIC SUITE

The Lower volcanic suite occurs in central Whitesides, and southern Carscallen and Bristol townships. Rock types include massive and pillowed and mafic rocks, and tuffaceous(?) rocks of intermediate and felsic composition. Mafic rocks predominate and comprise approximately 80% of the suite. At least one cherty sulphide-oxide iron formation 2 to 10 m thick is intercalated with the Lower volcanic suite (present in outcrop at the southeast end of Carscallen Lake), and a separate iron formation conformably overlies the suite. Ductile deformation is evident in many outcrops and obscures or obliterates primary textures (See Structural Geology section). Where deformed, the mafic rocks are hornblende-actinolite-plagioclase schists. Felsic rocks are principally tuffaceous and are comprised of sericite, quartz, and feldspar and clay minerals. Strong chlorite alteration is prevalent in felsic rocks below the iron formation in western Carscallen Township, and is similar in appearance to syn-volcanic alteration generally attributed to sea-floor hydrothermal systems. Carbonate $\pm$ fuchsite alteration is present adjacent to quartz - carbonate - sulphide veins that cut deformed Lower volcanic suite rocks in several locations, including at the Union Mine property (Figure 4).

METASEDIMENTARY ROCKS

Metasedimentary rocks are located in southern Bristol Township, and central Carscallen and Whitesides townships. They are comprised of metagreywacke, argillite,

metachert, and oxide- sulphide-bearing iron formation. At the Holmer Mine property in Bristol Township, the northeast part of the main outcrops is metagreywacke that consists of metre-thick units with grains that grade from coarse sand-sized to silt-sized, indicating facing directions to the south. These may represent thin turbidite deposits. Most iron formation and metachert occur in several outcrops along strike, from southern Carscallen Township to western Whitesides Township. This iron formation represents a single unit and contains up to 25% pyrite and 25% magnetite, either heavily disseminated through metachert or as discrete layers up to 20 cm thick. In central Carscallen Township, this iron formation rests conformably on pillowed basalts of the Lower volcanic suite, with magnetite and sulphide interstitial to relatively unaltered basalt pillows locally. Iron formation is also present in outcrop in southwestern Carscallen Township south of Carscallen Lake: this is believed to represent a separate unit.

GRANITOID ROCKS

Granitoid rocks are quartz-bearing, leucocratic intrusive rocks that are not related to the KGC or the KVC. They have been divided into four units. Three have been given informal names for the 1:50,000 map: the Turnbull Township tonalite, the Cote Township tonalite, and the Groundhog river tonalite (in Enid, Frey, and western Massey and Whitesides townships). In this report, these granitoid masses are termed granitoids A, B, C, respectively; a fourth granitoid mass, in southern Whitesides and Carscallen townships, is termed granitoid D (Figure 4). Other granitoid rocks are undifferentiated.

Granitoid A is characterized by mixed magma textures with chilled rocks of the KGC, and exhibits a wide variety of textural and modal characteristics. It is predominantly composed of medium-to coarse-grained tonalite and granodiorite, with lesser, quartz diorite, quartz monzodiorite, quartz monzonite and granite, and local quartz-feldspar aplite and pegmatite. Much of the granitoid A region is underlain by hybrid or agmatitic rocks where

KGC gabbroic material comprises blocks, fragments and pillow-like structures within a matrix of granitoid rock (Plates 3a-f).

Granitoids B and C are comprised of medium- to coarse-grained quartz diorite, tonalite, quartz monzodiorite, granodiorite, quartz monzonite, granite; and felsic aplite and pegmatite dikes or masses. Both intrusions have well-developed flattening fabrics at their margins, with prominent foliations and locally developed gneissic fabrics that parallel the intrusion contacts. The margins are characterized by intrusive breccia zones up to 3 km thick which contain flattened, angular blocks of wall rock. In granitoid B, angular blocks of anorthositic gabbro are found up to five km from the southern margin, and it appears that much of the intrusion contains large inclusions of gabbro.

BRISTOL TOWNSHIP LAMPROPHYRE SUITE

An unusual lamprophyre suite is found within the DPFZ in southwestern Bristol Township, on the Croxall and Holmer properties (Figure 4; see also Figures 32 and 34) (Plate 7). For this report, the suite is informally named the Bristol Township lamprophyre suite. The suite occurs within deformed and carbonate-altered metabasalts and metasedimentary rocks. Recently exhumed exposures on the Croxall property have revealed that the suite is greater than 60 m wide and has sub-vertical contacts. Ground magnetic surveys suggest that the suite extends under glacial cover to the southwest for several kilometres with widths up to 1 km. The suite is also present in outcrop 1.5 km to the northeast of the Croxall property on the Holmer property, where trace-element enriched, altered clinopyroxenites are found adjacent to and Au mineralization.

The Bristol Township lamprophyre suite is unique in its mineralogy and geochemistry in the southern Superior Province. It is comprised of three principal lithologic types, with mutually gradational and/or crosscutting relationships. The most abundant (50% of the exposures) is biotite lamprophyre found at the margins of the suite, with the groundmass

comprised of fine-grained ferroan carbonate, serpentine, biotite and apatite as principal constituents, and minor to trace amounts of potassium-feldspar, chlorite, magnetite, baddeleyite, sulphate and sulphide comprising the remainder. The next most abundant lithology (47% of the exposures) is a diopside-rich rock, comprised of medium- to very coarse-grained diopside in a distinctive bladed crystal habit, with interstitial chessboard albite (after potassium-feldspar), garnet, phlogopite, magnetite, and epidote. Garnetite and garnet-rich feldspathic dikes are the third lithology of the lamprophyre suite (3% of the exposures). These dikes contain from 7% to 85% red-brown garnet as fine-grained euhedra to very coarse-grained masses. Initial microprobe analyses indicate high calcium and iron contents, and > 5 wt. % TiO_2 and up to 3500 ppm Zr. This indicates that they are melanite-andradite (with a kimzeyitic component), common to ultramafic lamprophyres (Rock 1986). Light brown, anisotropic hydrogarnet or clinopyroxene is found rimming garnet, and is present up to 30%. Other minerals present are diopside, calcite, biotite, apatite; and minor potassium-feldspar, tremolite, titanite, allanite, magnetite, and chlorite and sulphides. There is one exposure of highly altered garnetite, with an octagonal-shaped, coarse-grained mineral (garnet?) as the principal constituent in fine-grained matrix of chlorite, biotite, epidote, magnetite, rutile and sulphides (pyrite, chalcopyrite). The garnet has been completely replaced by fine-grained calcite and dolomite, potassium-feldspar, albite, sericite, and very fine-grained rutile, baddeleyite and/or zircon.

STRUCTURE

BEDDING AND LAYERING, AND REGIONAL FOLDING

Facing directions are present in the metavolcanic strata and in KGC cumulates (Figure 4). In the Lower volcanic suite, facing directions from pillowed mafic metavolcanic rocks are consistently to the north through Whitesides, Carscallen and Bristol townships. In several of

these outcrops, bedding is overturned and dips steeply to the south. Grain size gradations in KVC pyroclastic crystal ash deposits indicate facing directions to the northeast in southern Godfrey Township, and face east in recently stripped outcrops in central Godfrey Township. These facing directions are consistent with well-exposed outcrops of KVC pillowed mafic metavolcanic rocks, which are east-facing in central Godfrey Township, and face to the northeast in northern part of this township, and in southern Robb and Jamieson townships. Transposition of bedding has occurred in the vicinity of the DPFZ, with a predominantly dextral sense of displacement on the horizontal plane; and near the fault that parallels Highway 576 near the Canadian Jamieson Mine, also with a dextral sense of displacement. The transposed strata near the Canadian Jamieson were originally more north-facing prior to transposition.

Facing directions in the KGC are based on modal gradations in layering, with "normal" mafic to felsic gradation in cumulus minerals pointing up stratigraphy (Plate 1a). This is generally the case in most intrusions that exhibit cumulate layering, although post-cumulus processes may create the opposite effect, and layering in alkaline intrusions commonly exhibits "reverse" gradations (Irvine 1982; Parsons 1979). Facing directions are also based on cross-bedding features. In western Whitesides Township, Lower Zone cumulates have facing directions to the east, with layering overturned and dipping steeply to the west (Plate 1b). Facing directions are to the north in northeast Carscallen and northwest Massey townships. Upper Zone cumulates show a consistent facing direction to the northeast in southern Robb Township.

In a broad sense, the stratigraphy (Lower volcanic suite, metasedimentary rocks, KGC, KfV) forms a regional monocline that faces to the north and northeast. Unfortunately thick glacial deposits cover a substantial portion of the MZ, which prevents a more detailed structural analysis in this area. Granitoids B, C, and D represent antiformal structures, and in general facing directions are away from their margins. It is possible that a synform, with a north-trending axis is present in the southwestern KGC. This would help explain the

apparent anomalous thickness of the KGC in the southern part. The apparent anomalous thickness of the northern KGC in southern Robb Township may be due to a higher proportion of felsic intrusive material than is mapped; bedrock in this area is also under glacial cover.

Linear features in the Kamiskotia area that are apparent on the Timmins area shadow-enhanced total magnetic field map (Barlow 1988) are presented on the accompanying 1:50,000 scale map. The shadow enhancement is designed to filter out the north-northwest-trending Matachewan dikes in the area. Several linear features on the shadow-enhanced magnetic map are interpreted to support or extrapolate significant geological features where there is poor bedrock exposure. Firstly, one north-northwest-trending linear feature is aligned with an offset of 2 km in a left-lateral sense, at the KGC - Lower volcanic suite contact in Carscallen Township. This linear feature is interpreted to represent a continuation of the fault that extends through western Turnbull Township into northwestern Robb Township, with 1 to 2 km of sinistral displacement on the sub-horizontal plane. Secondly, an oval, concentric magnetic pattern is apparent around granitoid B in Cote Township, consistent with granitoid B representing the core of a dome-like or antiformal structure. Thirdly, magnetic units in the Middle Zone of the KGC may form a steeply-plunging synform that opens to the west-southwest. This feature is present immediately south of granitoid B, in Massey Township, where there is a thick cover of glacial alluvium. (These features are included on the 1:50,000 map.) And fourthly, a prominent, northwest-trending linear feature extends from southeast Bristol Township to northwest Cote Township. This is interpreted to represent a post-Kenoran mafic dike that is not present at the surface in the Kamiskotia area.

STRUCTURAL FABRIC ANALYSIS

To determine the structural history of the Kamiskotia area, penetrative and non-penetrative foliations, lineations and L-S fabrics (Flinn 1965) were recorded, in the field in addition to the bedding attitudes described above. The foliation fabrics (Figures 6, 7) are primarily schistosity; however, gneissosity is present locally between the margins of granitoids B and C, within amphibolite-grade Lower volcanic suite rocks and leucocratic tonalite lenses. Lineations and L-S fabrics were recorded for outcrops with penetrative deformation wherever possible (Figures 8, 9). The L-S fabric scheme qualitatively estimates the prolateness factor of the strain ellipsoid by comparing the relative strength of lineation (L) with that of foliation (S - schistosity) for minerals or mineral aggregates, assuming that their initial state was quasi-spherical. For prolate strain ellipsoids, $L > S$ with the maximum extension direction parallel to the lineation measured in the field. When combined with regional lithologic, foliation and lineation patterns, the L-S fabric scheme has been used with success to discriminate between structural hypotheses in Archean terranes, particularly within and adjacent to granitoid intrusions (e.g., Schwerdtner et al. 1983; Stott 1985).

Results

Two prominent features appear in planar fabric maps (Figures 6, 7): zones of foliation that parallel the margins of granitoids B, C, and D; and east-trending foliations, particularly in the KVC. The zones of foliation parallel to the margins of granitoids B, C, and D extend up to 2 km to either side of the granitoid margins, and locally up to 5 km into the country rocks. The intensity of penetrative deformation (Plates 6a,b) diminishes with distance from the contacts. From the available outcrops, it appears that granitoid B is completely rimmed

by penetratively-deformed rocks. Fragments of gabbro and mafic metavolcanic rock are found within the foliated margins of granitoids B and C.

The second prominent feature noted in the planar fabric maps is the pervasive E-W foliation fabric across the Kamiskotia area, particularly in the metavolcanic rocks to the east (Figures 6, 7). This fabric is generally non-penetrative and cuts directly across the strike of the metavolcanic rocks. Furthermore it continues relatively undisturbed through granitoid A, in contrast to foliations that parallel the contacts of the other felsic intrusions. (Penetrative fabrics are confined to local, east-northeast- or east-southeast-trending high strain zones that contain quartz $\pm$ pyrite veins in the region of granitoid A). The east-west foliation fabric is most pervasive in the vicinity of base metal mineralization, where metavolcanic rocks that have undergone hydrothermal alteration (sericite and chlorite alteration). In metavolcanic rocks to the north, the fabric has a west-northwest trend; to the south, the foliations have a west-southwest trend.

A consistent, shallow west-plunging stretching fabric is pervasive in the Lower volcanic suite north of granitoid D, and at the C-D contact (Figures 8; 9). This fabric extends to the east and merges with highly strained rocks within the westernmost extent of the DPFZ. Accompanying this fabric is a reasonably consistent dextral sense of displacement (5 observations of dextral sense to 1 observation of sinistral sense) noted on the sub-horizontal plane in asymmetric, transposed fold hinges of siliceous material within strained Lower volcanic suite rocks, and within shear zones in granitoid D (Plate 6d). Between granitoids B and C, there are two prominent lineation trends, some plunging steeply to the north-northeast along the northwest contact of granitoid C, and the other with a shallow northwest-plunging orientation. Highly prolate ($L \gg S$) fabrics are concentrated between granitoids B and C, and along the northern contact of granitoid D, extending toward a part of the westernmost DPFZ (Figure 11). Highly strained rocks also occur in ductile shear zones that strike west-northwest with a dextral sense of displacement in KVC rocks near the Canadian Jamieson Mine (Plate 6c).

Highly prolate fabrics are present on the Holmer property in southwest Bristol Township. The Bristol Township lamprophyre suite on the adjacent Croxall property occurs within deformed and carbonate-altered basalts in the DPFZ. The margins of the suite are highly deformed, with a well-developed, sub-vertical flattening fabric that exhibits a component of elongation locally. The central portion of the suite is nearly undeformed, but is cut by sub-vertical shear zones up to 50 cm thick, and by brittle fractures with minor displacements of up to 20 cm.

Interpretation

The planar foliation fabrics that parallel the margins of granitoids B and C define contact strain zones; for granitoid D, where the fabric apparently encircles the intrusion, they define a contact strain aureole. These features may be the result of granitoid emplacement by diapirism and ballooning mechanisms. Schwerdtner et al. (1978) outlined three principal features that characterize diapiric structures. Firstly, an inverse density stratification is necessary for initial diapiric movement between a hot magma body and surrounding rocks. This is a common feature for granitic basement rocks, although densities may change after diapirism due to cooling of magma or an increase in metamorphism in the mantling material. This would pertain to the Kamiskotia area, where denser gabbroic rock ($\rho = 2.85 - 3.12$ g/cc) is cut by felsic intrusive rocks ($\rho = 2.52-2.96$ g/cc for granite to quartz diorite; $2.33 - 2.57$ g/cc for rhyolite to andesite liquids: Daly et al. 1966) at the base of the KGC. Secondly, the plan of a diapiric structure should have a continuous, smooth oval outline. This would pertain to granitoid B, and may also be the case for granitoids C and D, considering their regional aeromagnetic signatures and topographic expressions. (Unfortunately poor outcrop to the west and south of the mapped area limits the interpretation of these criteria for parts of granitoids C and D.) Thirdly, subhorizontal stretching fabrics should be present across the crest line of the diapiric structure. However

possible crest lines with a subhorizontal stretching fabric are not preserved or exposed for these intrusions. Granitoids B, C, and D meet two of the three criteria for diapirism, and they are interpreted to have been emplaced by diapiric processes.

As the flattening fabric in the margins of granitoids B, C, and D and in mantling country rock is well-developed, ballooning may have occurred in addition to diapirism. After partial crystallization at their margins, these intrusions may have expanded and ballooned in-situ as more magma entered the chamber, producing a radial compressive force that would create the observed tangential flattening fabric (e.g., Bateman 1985).

The fragments of gabbro and mafic metavolcanic rock within the foliated margins of granitoids B and C suggests that stoping of adjacent country rock may have been an operative process, in addition to diapirism and ballooning. Stopping is believed to be a common intrusive mechanism during the final stages of granitoid emplacement at high crustal levels (Bateman 1985), particularly by magmas with lower water contents (Marsh 1982).

The E-W foliation fabric may be a product of a late north-south compressional event that effected much of the southern Superior Province (Stott et al. 1987). The deflection of foliations away from an east-west trend in the metavolcanic rocks may be the result of a competency contrast between locally altered metavolcanic rocks and the more competent, relatively unaltered intrusions to the west. Within the contact strain zone of granitoid C, this fabric is manifested as a weak, non-penetrative foliation or parting cleavage which locally cuts the foliation produced by the emplacement of granitoid C. This provides evidence that at least some north-south compression post-dated the emplacement of granitoid C. Furthermore, the east-west fabric cuts across the strike of the subvertical KVC, indicating that north-south compression postdated a passive, bulk crustal rotation of the stratigraphy, that did not produce any rock fabric.

That granitoid A is cut by the east-west foliation fabric indicates that it was emplaced in a manner entirely different from that of the other felsic intrusions. Granitoid A is characterized by a variety of mixed magma textures with the basal part of the Middle Zone

of the KGC, indicating that it was emplaced during the solidification of the KGC and is therefore contemporaneous with it. Possible flexures on the western side of granitoid A may be related to strain from neighboring granitoid intrusions.

The two lineation trends between Granitoids B and C may be related to granitoid C emplacement. The steep north-northeast-plunging lineations are consistent with an upward diapiric movement of granitoid C with respect to the surrounding rocks. In laboratory experiments of diapiric structures using Plasticine, Dixon (1975) noted that sub-vertical prolate strain ellipsoids will develop along the flanks of diapiric structures of circular or oval plan. Similar features have been documented at the margins of crescentic granitoid plutons in northwestern Ontario (Schwerdtner et al. 1983). The shallow northwest-plunging lineations are found within granitoid C, within its intrusive breccia zone and into the KGC near granitoid B, overprinting any previous fabric formed from granitoid B emplacement. This fabric may be a product of ballooning by granitoid C against a competent wall of granitoid B, creating a subhorizontal stress regime where the less competent mantling material was attenuated parallel to the C axis of the strain ellipsoid. Alternatively, it may reflect strain created by flow of magma/crystal mush within granitoid C after the formation of a gneissic carapace and wall. Similar models have been proposed to explain subhorizontal stretching fabrics in crescentic granitoid plutons (Schwerdtner et al. 1983).

The shallow, west-plunging stretching (elongation) fabric within the Lower volcanic suite and in the margins of granitoid D may have been caused by several processes: 1) emplacement of granitoid D, 2) bulk north-south shortening, or 3) ductile shear along a splay of the DPFZ, possibly related to late north-south bulk shortening or transpression. Although this west-plunging fabric is parallel to the margin of granitoid D, it extends eastward for 10 km past the nose of granitoid D and on into the DPFZ. As the contacts of granitoid D are near vertical, it is unlikely that its contact strain effects would extend 10 km and beyond in one direction. Bulk north-south shortening could produce this fabric between two competent masses (granitoid D and the KGC) if, during deformation, the Lower volcanic suite was

confined by competent material above and below. However, the continuation of this strain fabric in metavolcanic rocks to the east away from any competent intrusive masses argues against pure north-south compression. The third process considered, related to ductile shear along the DPFZ, is supported by the following structural and geological evidence. In addition to merging with the structural fabric of the DPFZ, this prolate strain regime predominates over any flattening fabric that could be attributed to granitoid D emplacement or pure north-south compression. Furthermore, this deformation zone contains numerous occurrences of lode gold mineralization, with syn- to posttectonic, quartz - tourmaline + sulphide $\pm$ carbonate $\pm$ Au veins and associated carbonate alteration, similar to occurrences found in the western DPFZ.

Summary

Based on geology and structural fabric analysis, the sequence of major magmatic and tectonic events in the Kamiskotia area are interpreted as follows: 1) formation of the Lower volcanic suite; 2) formation of the KGC-KVC, accompanied by coeval intrusion of granitoid A into the KGC during solidification; 3) regional bulk rotation of the stratigraphy about a sub-horizontal axis, to form a northeast-facing monocline; possibly accompanied by 4) diapiric intrusion of granitoids B, C and D (timing with respect to each other ambiguous), imparting their respective contact strain aureoles or zones on mantling material; 5) regional north-south compression, probably in part synchronous with, but locally postdated by 6) ductile shear along the western DPFZ, and a related splay through the Lower volcanic suite, that exhibits a consistent sense of dextral displacement.

GEOCHRONOLOGY

Primary igneous ages from U-Pb geochronology, and primary and secondary ages from Sm-Nd and Rb-Sr isochron systematics have been determined for Kamiskotia area rocks in studies by Barrie (1990), Barrie and Davis (1990), and Barrie and Shirey (in prep.). In this section, the results from these studies are reviewed, and then the chronologic constraints on the magmatic and structural history of the area are discussed. Sample locations are marked in Figure 10. Universal Transverse Mercator grid coordinates for the sample locations with brief petrologic descriptions are provided in Appendix I.

Results of U-Pb Geochronology

A pegmatitic quartz gabbro sample taken from the Middle Zone of the KGC give an age of 2707 ± 2 Ma. A quartz- and feldspar-phyric flow-banded rhyolite sample of the Kamiskotia rhyolite has an age of 2705 ± 2 Ma. This age has an overlapping error with the age for the KGC.

Samples of well-foliated tonalite were taken from the contact strain aureole of Granitoid B (Cote Township tonalite) and the contact strain zone of Granitoid C (Groundhog River tonalite). Zircons from granitoid C yield an age of 2696 ± 1.5 Ma. One titanite fraction, from the same sample is nearly concordant, with a $^{207}\text{Pb}/^{206}\text{Pb}$ age of 2692 ± 5 Ma. One zircon fraction from granitoid B has a $^{207}\text{Pb}/^{206}\text{Pb}$ age of 2926 Ma and is 4.6% discordant. This most likely represents an inherited component. Four more analyses of abraded, small zircon fractions are all slightly discordant, with $^{207}\text{Pb}/^{206}\text{Pb}$ ages ranging from 2704 Ma to 2693 Ma. One grain with a $^{207}\text{Pb}/^{206}\text{Pb}$ age of 2704 Ma may be inherited from a slightly older component, and the preferred crystallization age is an average of the three lower $^{207}\text{Pb}/^{206}\text{Pb}$ ages at 2694 ± 4 Ma.

In an attempt to provide for time-stratigraphic correlation with other the felsic volcanic rocks in region, U-Pb ages are reported for three felsic metavolcanic rocks in the Kidd Creek area, including the addition of one abraded, concordant fraction to previous analyses of the Kidd Creek rhyolite by Nunes and Pyke (1981), and two new ages. For the Kidd Creek rhyolite, the new age is more precise than the age previously reported, at 2717 ± 2 Ma. Two other rhyolites, termed the Prosser (Township) rhyolite and the Reid (Township) rhyolite, are quartz- and feldspar-phyric tuffaceous horizons intercalated with metasedimentary rocks, located to the east and west of the Kidd Creek mine, respectively (Figure 10). The Prosser rhyolite is 2716 ± 4 Ma, and the Reid rhyolite is $2705 \pm 5/-3$ Ma. The Prosser rhyolite is coeval within error with the Kidd Creek rhyolite, whereas the Reid rhyolite is coeval with the KGC and Kamiskotia rhyolite 40 km to the south-southwest.

U-Pb isotopic data for two garnet fractions and one titanite fraction from a garnetite dike in the Bristol Township lamprophyre suite yield an age of 2687 ± 3 Ma. The garnetite dike is undeformed; however, the lamprophyre suite is deformed along its margins, with the fabric parallel to the fabric in the DPFZ. The lamprophyre suite is interpreted to have utilized the DPFZ as a conduit, thus the age represents a minimum age for the early formation of the DPFZ in this area. The U-Pb system in garnet is apparently more resilient than other isotopic systems (Rb-Sr, Ar-Ar, Sm-Nd) for whole rock suites and minerals associated with lode Au mineralization within the DPFZ, which show evidence for resetting or disturbance (Kerrick et al. 1987; Bell et al. 1989; Wong et al. 1989; Barrie 1990).

Sm-Nd and Rb-Sr Isochron and Regression Ages

Mineral separates for four whole rock samples were analyzed for Sm-Nd and Rb-Sr isotopes to produce mineral - whole rock isochrons, which are summarized in Table 3. These isotope systems are set at the time of crystallization, but may be reset during subsequent metamorphic events. The McIntyre 1 regression model (McIntyre et al. 1966) has

been used to calculate regression lines for these data. If the regression has an MSWD (mean square of the weighted deviates) of less than 3, the scatter in the data can be accommodated by analytical uncertainty, and the regression line is termed an isochron.

A seven point Sm-Nd isochron for rocks and mineral separates from the KGC and KVC, including plagioclase and clinopyroxene analyses, has an age of 2710 ± 30 Ma (2- σ), in complete agreement with the KGC and KVC U-Pb ages. An isochron from Rb-Sr analyses of a whole-rock KGC ferroan gabbro and clinopyroxene and zircon fractions derived from the ferroan gabbro sample has an age of 2450 ± 35 Ma. This age is identical within error to three U-Pb discordia ages from regressions of baddeleyite and zircon fractions, all at $2452 \pm 3/-2$ Ma, from the Hearst - Matachewan mafic dike swarm (Heaman 1989); and one baddeleyite fraction (2460 Ma: L. Heaman, personal communication, 1989) from a Matachewan dike 12 km from this location.

Regression of isotope data for mineral separates from samples of granitoids B and C produce ages younger than their zircon and titanite U-Pb ages in the Sm-Nd and Rb-Sr systems. For the granitoid B, a whole rock - plagioclase - apatite₁ - apatite₂ (two separate apatite reactions) Sm-Nd isochron has an age of 2615 ± 15 Ma, significantly younger than the 2696 ± 4 Ma U-Pb age from the same rock. This age represents essentially a two point isochron, controlled by the apatite fractions with their similar isotopic ratios, and the precision of the age error should be considered with some caution. The Rb-Sr system does not yield meaningful results, indicating that Rb or Sr or both were open to remobilization after crystallization in this sample.

For granitoid C, regressions of the Sm-Nd and Rb-Sr data give ages that are identical at 2530 ± 35 Ma, younger than the U-Pb zircon age of 2696 ± 2 Ma for the same sample, and a U-Pb titanite $^{207}\text{Pb}/^{206}\text{Pb}$ minimum age of 2692 ± 5 Ma. The agreement between the Sm-Nd and Rb-Sr isotope systems is strong evidence that this age has geological significance. Similar ages of 2530 Ma, 2508 Ma, and 2504 Ma have been reported for individual,

concordant metamorphic titanite grains in granulite and upper amphibolite grade rocks of the nearby Kapuskasing Structural Zone 40 km to the west (Krogh et al. 1988).

Magmatic and structural history

A chronologic summary of the magmatism and deformation in the Kamiskotia - Kidd Creek area based on the U-Pb data is presented in Figure 11. Three periods of magmatic activity occurred at approximately 10 Ma intervals: felsic volcanism in the Kidd Creek area from 2717 ± 2 to 2716 ± 4 Ma, volcanism and hypabyssal mafic and felsic intrusions at 2707 ± 2 to 2705 ± 2 Ma, and voluminous granitoid emplacement at 2696 ± 1.5 to 2694 ± 4 Ma. The timing of deformation events, constrained by geologic relationships along with U-Pb ages, is in part coincident with and related to the magmatism, with regional crustal warping between 2705 ± 2 to 2694 ± 4 Ma, and with contact strain aureole/zone development. N-S compression postdated contact strain development for granitoid C. A minimum age for the early formation of the DPFZ may be provided by a U-Pb age for the garnetite dike from the Bristol Township lamprophyre suite, at 2687 ± 3 Ma.

GEOCHEMISTRY

Major, trace and rare earth element analyses for KGC, KVC, Lower volcanic suite rocks, granitoid rocks, and the Bristol Township lamprophyre suite are presented in this section. Particular emphasis is given to the characteristics and petrogenesis of the KGC, and its relationship to KVC basalts.

Alteration

Major and incompatible trace elements for the rock suites presented in this section plot as coherent trends on inter-element variation diagrams, and are believed to represent the primary compositions, despite post-solidus alteration and metamorphism. This is in accord with studies on the mobility of major and trace elements in greenstone belt rocks that have been subjected to greenschist facies metamorphism and normal surficial weathering processes (Menzies et al. 1979; Ludden et al. 1982; Beswick 1982). Variation diagrams for KGC cumulates and KVC basalts, K_2O , Rb and Ba do not correlate well with elements of similar geochemical affinities: therefore, little emphasis is placed on these elements in the mafic rocks.

KGC Cumulates

Whole rock major and trace element compositions for 33 medium to coarse-grained cumulus samples are given in Table 4 and located in Figure 12. They include four olivine-bearing cumulates of the LZ: two peridotites (samples 1 and 2) and two troctolites (samples 3 and 4), other LZ cumulates (samples 15) and MZ cumulates (samples 16-23) which constitute the Western Traverse, and UZ cumulates (samples 24-33) which constitute the Northeastern Traverse (Figure 12). The samples are assigned a stratigraphic height by projection along strike to the traverse lines. These heights have errors of approximately ± 5 m to 50 m, with larger errors for the UZ samples located far from the traverse lines. Sample locations are given by Universal Transverse Mercator grid coordinates in Appendix II.

Samples 8, 9 and 10 were taken 3 to 5 m apart from the same outcrop of moderately altered gabbro-norite in the LZ as a test for trace element mobility. Their primary modes of

plagioclase, clinopyroxene and orthopyroxene respectively are estimated as: sample 8: 45-40-20, sample 9: 50-25-25, and sample 10: 45-25-30. Their similar trace element geochemistry suggests that the REE, the high field strength elements, V, Cr and Sr have not been disturbed by post-solidus alteration or metamorphism. Variation in Sc content is proportional to modal clinopyroxene content, as Sc partitions favorably into clinopyroxene (Irving 1978). Ni is higher in sample 9; this may be due to trace amounts of sulphide not detected in hand specimen or thin section.

The modes for the traverse samples are presented in Figure 13. These samples are meso- and adcumulates, with anhydrous silicates comprising the bulk of the cumulus and post-cumulus phases. One exception is sample 13 where hornblende comprises 11% of the mode as a primary, post-cumulus phase. Magnetite, Ti-magnetite and ilmenite are postcumulus phases in the Western Traverse, and they are both cumulus and post-cumulus phases in the Northeastern Traverse. Biotite and apatite are present as post-cumulus phases in sample 24; apatite may be a cumulus phase as well. Sulphides (pyrite, pyrrhotite, chalcopyrite) occur as trace phases interstitial to the silicates commonly in the olivine cumulates in the Western Traverse. They also occur sporadically in the Northeastern Traverse.

Whole rock Mg numbers and normative An, TiO_2 and P_2O_5 contents versus stratigraphic height are presented in Figure 14. Broadly, Mg numbers and normative An contents decrease from 82-86 and 91-96, respectively in the olivine-bearing cumulates to 37-53 and 60-76, respectively for the upper four UZ cumulates. This is consistent with normal fractionation and Fe-enrichment trends observed in many large layered intrusions such as the Bushveld and Skaergaard complexes (Wager and Brown 1968). TiO_2 and P_2O_5 are low in LZ cumulates and increase sporadically in the MZ and UZ cumulates, reflecting the presence of oxide phases and apatite, respectively.

Nickel contents are highest in LZ cumulates, consistent with the presence of olivine and sulphide which strongly favor Ni (Figure 15). Scandium abundances are low in the

olivine-bearing cumulates and the majority of LZ rocks (Figure 15). Scandium partitions favorably into clinopyroxene and oxide phases but not into olivine, orthopyroxene or plagioclase under reasonable magmatic temperatures and oxygen fugacities (Lindstrom 1976).

In Figure 16, incompatible elements and incompatible element ratios are plotted versus stratigraphic height. The abundances of the REE and high field strength elements in cumulus rocks are in a general sense proportional to the amount of intercumulus liquid trapped during post-cumulus crystallization. It is apparent from Figure 16 that the majority of LZ cumulates have very low La, Yb, Zr and Y contents and thus contain low amounts of intercumulus material: this is consistent with their adcumulus and mesocumulus textures. In comparison, the MZ and UZ cumulates are relatively enriched in these elements. In the MZ, more closely spaced samples from 2000 m to 3000 m show a consistent trend toward lower concentrations upsection. Three samples at the base of the UZ decrease in incompatible element abundances in a generally similar manner. This may reflect a decrease in the mode of intercumulus phases upsection which is not detectable petrographically. In both cases, Zr/Y follows the decrease in incompatible element contents; this may be due to Y^{3+} behaving more compatibly in clinopyroxene than Zr^{4+} . Lanthanum in the MZ samples remains relatively constant from 2000 to 3000 m, contrary to the behavior of the other trace elements. One explanation for the constant La content would be that cumulus pyroxenes are present in greater abundance where there is more trapped liquid. In the UZ, Yb, Zr and Y increase from 1000 m to 2500 m, accompanied by a decrease in Mg number from 64 to 37-52 (Figure 16), and corresponding decreasing Mg numbers in clinopyroxene and An contents of plagioclase (samples G-I, Figure 5). These trace element trends in the UZ are consistent with normal fractionation accompanied by assimilation of trace element-enriched siliceous material (discussed in detail below).

The cumulates of all three zones have flat REE patterns, with $La_N/Yb_N = 0.4 - 2.6$, and increasing total REE abundances upsection (Figure 17a, b). There is a strong negative correlation between La_N/Yb_N and Sc and clinopyroxene content, consistent with lower K_D s

for the light REE between clinopyroxene and basaltic liquids (Irving 1978). Europium anomalies are positive in the LZ with Eu/Eu^* of 1.2 to 5, and show a progressive depletion upsection to Eu/Eu^* of 0.8 to 0.85 for the upper four UZ cumulates. In general the REE are consistent with enrichment due to fractionation of plagioclase and mafic phases from a tholeiitic parent, accompanied by the addition of REE-enriched material (discussed below). Except for the upper four UZ samples, the KGC cumulates have REE contents similar to anorthositic gabbronorites from the Shawmere anorthosite of the Kapuskasing Structural Zone (Simmons et al. 1980) and massive gabbros from the Bad Vermilion complex, Northwestern Ontario (Ashwal et al. 1985). Their REE contents are distinct from ophiolitic gabbro cumulates which are characterized by LREE-depletion (e.g., Pallister and Knight 1981 and references therein).

KGC Chilled Rocks

Geochemistry for KGC chilled samples and average compositions of KVC basalts and evolved basalts are given in Table 5, and the sample locations are given in Appendix II.

There are several locations where chilled KGC gabbroic rocks are exposed. Samples 34-37 are from the mixed magma outcrops; sample 38 is from a contact with felsic metavolcanic rocks that interdigitate with the KGC margin, where the felsic rocks show no textural evidence of melting due to the juxtaposition of the hot gabbroic liquid (Figure 4). Care was taken to ensure that no felsic material was included during sample collection. These samples are composed of aphanitic basalt 1 to 2 cm in from the margin, to medium-grained gabbro, 50 cm in from the contact. They have similar geochemical characteristics, with Mg numbers from 54 to 58, TiO_2 from 0.88 to 1.01 wt.%, and generally similar REE and trace element compositions (Table 5).

Sample 34 has 8 to 12 times higher K_2O , Ba and Rb contents, but less than half the Sr content of an average of the other four samples. This may be attributed to preferential

diffusion of the alkalis and Ba from the adjacent felsic liquid into sample 34 during (and after?) solidification. Experiments in basaltic liquids at 1200°C to 1400°C indicate that initially, potassium diffuses much more rapidly than other melt species until the basalt has contents of around 1.5 wt.% K₂O, after which the basalt is buffered from further potassium diffusion (Watson 1982). With their similar cation charge to radius ratios, Rb and Ba would behave in a similar manner. In granitic melts, the rates of diffusion of the alkalis are known to be two or more orders of magnitude greater than for more highly charged cations (e.g., Ce, Eu) under a wide range of temperatures (Jambon 1982).

Watson's experiments (1976) on element partitioning between coexisting mafic and felsic melts indicate that the REE and HFS elements should partition into the less polymerized mafic melt by factors of 3 to 13. Wholesale REE and HFS enrichment does not seem likely for the KGC mixed magma chilled samples for two reasons. Firstly, whereas their major element compositions are similar to N-type MORB, they have slightly lower abundances of the REE and HFS elements in comparison to N-type MORB. Secondly, their concentrations are nearly identical to sample 38, taken from outcrops away from the mixed magma outcrops. It seems probable that mixed magma chill samples quenched rapidly enough to prevent the diffusion of the REE and HFS elements from the felsic liquid under liquid-liquid equilibrium conditions.

The REE for the KGC chill samples are plotted along with the range of the REE for the LZ in Figure 17a. They have a restricted range, with flat REE profiles ($\text{La}_N/\text{Sm}_N = 0.9 - 1.3$; $\text{La}_N/\text{Yb}_N = 0.8 - 1.2$) at 15-20x chondritic values, and consistent negative Eu anomalies. In comparison to southern Abitibi Subprovince basalts, these REE abundances are most similar to parts of the lower Skead Group south of Kirkland Lake, Ontario (Figure 1i in Capdevila et al. 1982). Many of the Skead Group basalts are interpreted as the products of partial melting of a slightly depleted mantle where the degree of melting is sufficiently high to leave only olivine and orthopyroxene in the residue, followed by minor fractional crystallization of olivine and plagioclase (Capdevila et al. 1982). The KGC chill

samples are distinguished from LREE-depleted, N-type MORB (e.g., Schilling 1976; Frey et al. 1974) and many ophiolite-related basalts (e.g., Pallister and Knight 1981 and references therein) by their flat REE patterns and consistent Eu anomalies. They have flatter REE patterns than the LREE-enriched parental ultramafic or basaltic liquids of the Bushveld Complex (average $Ce_N/Yb_N = 12$ and 3 , respectively: Sharpe and Hulbert 1985) and the majority of gabbro-norite sills of the Stillwater Complex (average $La_N/Yb_N = 3.5$: samples 3-10 in Helz 1985).

Granophyric Rocks

A brief characterization of the KGC felsic granophyric rocks is given here, based on fourteen samples from Godfrey Township collected and analyzed by Hart (1984). Felsic granophyres are characterized by high silica contents, (64.5 to 77.4 wt.% SiO_2), low magnesium contents (0.3 to 1.5 wt.% MgO), and relatively low alkali and alkaline earth contents (2.5 to 5.6 wt.% Na_2O , 0.3 to 2.6 wt.% K_2O , 4 to 39 ppm Rb, 19 to 84 ppm Sr). They have high REE contents (33 to 74 ppm La, $n = 13$) and flat REE patterns ($La_N/Yb_N = 1.6$ to 2.6) with negative Eu anomalies, and very high zirconium contents (525 to 825 ppm, $n = 13$).

The majority of granophyric rocks of intermediate composition are interpreted as altered felsic granophyres. Ten samples from Godfrey Township analyzed by Hart (1984) have incompatible element concentrations very similar to the felsic granophyric rocks, and differ from them only in their lower silica contents (60 to 68 wt.% SiO_2) and higher magnesium contents (1.2 to 4.5 wt.%, $n = 9$). This may be attributed chlorite alteration, probably related to sub-sea floor hydrothermal alteration (see Economic Geology section).

Kamiskotia Volcanic Complex

Basalts, and Basaltic Andesites and Andesites

Kamiskotia basalts are predominantly massive or pillowed flows that contain less than 15% plagioclase and pyroxene phenocrysts. They have been divided into "primitive" and "evolved" basalts on the basis of geochemistry by Hart (1984). The averages listed in Table 5 are from samples taken away from chlorite alteration zones related to Cu-Zn mineralization (locations given in Hart 1984; and Barrie 1990). Kamiskotia primitive basalts are mildly enriched in TiO_2 , P_2O_5 , REE and HFS elements in comparison to the KGC chill samples, and in comparison to N-MORB. "Evolved" basalts are comprised of basalts, basaltic andesites and andesites, that are significantly enriched in TiO_2 , P_2O_5 , REE and HFS elements; they are transitional with highly enriched andesites similar in composition to basaltic andesites and icelandites of eastern Iceland (e.g. Wood 1978). The "evolved" basalts have flat REE patterns (average $\text{La}_\text{N}/\text{Yb}_\text{N} = 1.47$ and 1.60 , respectively) with moderately negative to no Eu anomalies (Figure 17c).

One of the andesitic fragments with apparent quenched margins from within KVC welded rhyolite tuffs (Plate 5b) has been analyzed for major elements by XRF (same sample as plotted in Figure 12 of Comba et al. 1986; P. Binney, Falconbridge Limited, personal communication, 1988). The oxides for this sample are, in wt. %: $\text{SiO}_2 = 54.5$, $\text{TiO}_2 = 2.15$, $\text{Al}_2\text{O}_3 = 12.8$, $\text{Fe}_2\text{O}_3 = 11.0$, $\text{MgO} = 1.9$, $\text{MnO} = 0.34$, $\text{CaO} = 5.19$, $\text{Na}_2\text{O} = 2.25$, $\text{K}_2\text{O} = 2.9$, $\text{P}_2\text{O}_5 = 0.36$, $\text{LOI} = 6.23$, and $\text{Total} = 99.7$. The Mg number for this sample is 34, and $\text{Zr} = 290$ ppm, determined by a semi-quantitative, fast-count XRF method. This sample has similar TiO_2 and P_2O_5 contents, but higher potassium and LOI contents than the incompatible element-enriched andesites described above.

Rhyolites

KVC rhyolites occur as quartz and feldspar-phyric pyroclastic flows, tuffs and agglomerates, with phenocryst content up to 15%. Hart (1984) analyzed 15 rhyolite samples taken away from hydrothermal alteration associated with the Cu-Zn deposits, in south Jamieson and north Godfrey townships (also reported in Leshner et al. 1986). These samples are characterized by high silica contents, with $\text{SiO}_2 = 71.9$ to 78.6 wt.% (average of 74.8 wt.% SiO_2 , $n = 14$), and very high immobile, incompatible element contents, with $\text{La} = 39$ to 67 ppm, $\text{Yb} = 7$ to 18 ppm, $\text{Zr} = 215$ to 410 ppm, and $\text{Th} = 6$ to 11 ppm ($n=15$). The rhyolites exhibit a range in K_2O and Rb contents, with $\text{K}_2\text{O} = 2$ to 9 wt.% and $\text{Rb} = 30$ to 130 ppm ($n=15$). Rhyolite samples with higher K_2O and Rb contents are located from 1 to 2 km upsection to the northeast from the Canadian Jamieson and Genex Mine properties, and are also within one km of the main Kamiskotia Highway fault. Rhyolite flows and welded tuffs in central and southern Godfrey Township have similar chemical characteristics with those to the north. The rhyolites in north Carscallen and southeast Turnbull townships do not have as high abundances of these elements. However, they have identical incompatible element ratios as those in Godfrey and Jamieson townships and are included as part of the KVC (unpublished data).

Hart (1984) and Leshner et al. (1986) have noted that the incompatible element-enriched chemistry of the KVC rhyolites is similar to the chemistry of Kidd Creek agglomeratic rhyolites from the east outcrops of the Kidd creek mine site. One of the notable characteristics for both Kamiskotia and Kidd Creek rhyolites is their relatively flat, chondrite-normalized REE patterns with large negative Eu anomalies (Figure 18a). These REE patterns are distinctive from many other felsic metavolcanic sequences in the southern Superior Province, which generally have steeper chondrite-normalized REE patterns with lower heavy REE contents (e.g., "Types FI" and "FII": Figure 18b). Kamiskotia and Kidd

Creek rhyolites are also distinguished by their high Y contents (Figure 18c,d). In these respects they are similar to rhyolites in Tertiary and Recent bimodal volcanic fields, such as in the Basin and Range Province, U.S.A. (e.g., Coso volcanic field, California: Bacon et al. 1981; Kaiser Spring volcanic field, Arizona: Moyer and Esperanca: 1989); and in eastern Iceland (e.g., eastern Iceland rift zone: Wood 1978; 1875 eruption of Askja volcano: Macdonald et al. 1987).

Lower Volcanic Suite

Twelve samples from the Lower volcanic suite range in composition from basalts to andesites; no felsic metavolcanic rocks were sampled (Table 6). The samples have from 43.8 to 59.6% SiO₂, and Mg numbers from 30 to 64. They have relatively low REE contents, with La from 1.9 to 10.5 ppm, Yb from 0.6 to 3.6 ppm; and variable REE patterns, with La_N/Yb_N from 0.6 to 4.3. The variable REE patterns along with a range in incompatible element ratios is interpreted to reflect heterogeneous sources for the suite. It is possible that several basalt samples which have chemical characteristics similar to KVC basalts or KGC chilled samples, are part of the KVC (e.g., samples 84-72, 84-265, 86-12).

Granitoid Rocks

Major element analyses are reported for eleven granitoid samples taken from granitoids A, B, C and D, and trace element analyses are given for six of these (Table 7). On normative Q-A-P and Or-Ab-An diagrams, these rocks plot in the diorite, monzodiorite, tonalite and granodiorite fields. Chondrite-normalized REE profiles for two samples from granitoids A, B and C show distinguishable patterns (Figure 19). Samples of granitoid A, from the mixed magma outcrops, have higher REE abundances and flat REE patterns, with La_N/Yb_N = 2.9 and 6.3, and negative Eu anomalies. These REE patterns are similar in

character to KVC rhyolites, and KGC granophyre rocks, although they have lower concentrations. These samples may represent KGC felsic liquids from that mixed with KGC mafic liquids in the MZ, although it is noted that several other tonalitic samples of granitoid A have much steeper REE patterns that cannot be related to the KGC-KVC suite (Barrie 1990). Samples of granitoid B have steeper patterns $\text{La}_N/\text{Yb}_N = 6.3$ and 7.9 , with no Eu anomalies. The major and trace element chemistry of the granitoid B samples is typical of Archean tonalites that interpreted to be derived from partial melting of tholeiitic material with hornblende and garnet in the residuum (e.g., Arth and Hanson 1975; Martin 1987).

Samples of granitoid C, taken from rocks in the contact strain zone, have relatively low incompatible element contents; and steep REE patterns, with $\text{La}_N/\text{Yb}_N = 14.9$ and 15.9 , and slight positive Eu anomalies (Figure 19). These samples have a high modal content of plagioclase and alkali feldspar, reflected in their high Na_2O , K_2O , Sr and Ba and Eu contents (Table 6). The high alkali and alkaline earth but low incompatible element contents can be explained either by two processes. Firstly, if the feldspars represent meso- to adcumulus phases, then cumulus and/or post-cumulus processes, normally considered for mafic cumulates, may have been active in granitoid C and led to the removal of incompatible any element-enriched intercumulus liquid. A second process to explain the low incompatible element contents is in-situ deformation. If these rocks were deformed in-situ prior to total consolidation during diapiric emplacement, then a filter-pressing effect may have squeezed out any incompatible element-rich liquid.

Bristol Township Lamprophyre Suite

Geochemical analyses for fourteen samples (and one duplicate) of the Bristol Township lamprophyre suite, and three samples of adjacent, altered basalts are given in Table 8. Sample locations are given in Figures 32 and 34, and in Appendix II. Two biotite-rich lamprophyres, three garnetite samples and one altered garnetite sample have been

analyzed by ICP-MS for the REE, after total dissolution was achieved with a lithium metaborate flux.

The lamprophyre suite is characterized by very low silica contents (averaging 38.4 wt.% SiO₂, n = 14); high titanium and phosphorous contents (1.9 wt.% TiO₂ and 2.4 wt.% P₂O₅, n = 14) and very high, high field strength (HFS) and light REE contents (e.g., 970 ppm Zr n = 13; 157 ppm Y, n = 14; 40 ppm Th, n = 13; 680 ppm Ce, n = 6; 660 ppm Nd, n = 6). The biotite-rich lamprophyres have up to 6 wt.% P₂O₅, and up to 4.5 wt.% K₂O, reflecting their high apatite and biotite contents. The garnetite samples are highly enriched in TiO₂, Zr and the light REE. The altered garnetite samples are very highly enriched in these elements, and also contain appreciable S and Cu (Table 8).

Selected major and trace elements are plotted versus zirconium in Figure 20. With its high charge (+4) and small ionic radius (0.79 by 10⁻¹⁰ m), zirconium is a highly incompatible element in the crystal structures of the common rock-forming silicates. Because of this, zirconium is an excellent indicator of silicate mineral fractionation in most magmatic systems. Additionally, it can be used to track HFS and light REE enrichment in certain volatile-rich systems. Volatile activity is believed to have contributed to the HFS and light REE enrichment in this lamprophyre suite, considering the high carbonate (and halogen: unpublished data) contents in many of these rocks.

The MgO, Ni, K₂O and P₂O₅ contents all decrease with increasing Zr contents in the lamprophyre suite (Figure 20a). Enrichment in Zr is accompanied by increasing TiO₂, Y, Th and S contents (Figure 20b). This is interpreted to reflect the fractionation of a magnesium-rich silicate, possibly diopside, in addition to biotite and apatite fractionation. Diopside has a mineral/liquid partition coefficient of 1 to 3, much lower than that of olivine at 10 to 20 or sulphide (200 or greater) in basaltic magmas (Irving 1978). Olivine or sulphide fractionation would cause the nickel contents to drop precipitously. Titanium and yttrium are present along with zirconium in the melanite garnets, and they form linear trends with zirconium due to the addition of garnet in this suite. It is interesting to note that sulphur

and copper increase with zirconium, which suggests that the sulphide phases became saturated and precipitated along with garnet, during the late stages in this system.

Chondrite-normalized REE profiles for six samples from the lamprophyre suite are steep, with no Eu anomalies (Figure 21). They have $La_N = 400$ to 1500 , $Nd_N = 350$ to 2900 , and $Yb_N = 5.5$ to 19 . The flattest pattern is also the most enriched sample, an altered garnetite, with $La_N/Yb_N = 8$; and the steepest pattern is for a biotite lamprophyre with $La_N/Yb_N = 90$ (Figure 21). The patterns are elevated in comparison to tholeiitic and calc-alkalic rocks, and are higher than most alkalic suites, including lamprophyres (Rock 1987). They are similar to the patterns found in some fluorapatite-rich rocks found along the DFPZ in Taylor Township (King and Kerrich 1987); and to some of the less enriched carbonatites in the Oka Complex, Quebec (Eby 1975). They are similar to the patterns found in some fluorapatite-rich rocks found along the DFPZ in Taylor Township (King and Kerrich 1987); and to some of the less- enriched carbonatites in the Oka Complex, Quebec (Eby 1975). They are also similar to estimates of average kimberlite, lamproite and metasomatic fluids believed to be in equilibrium with richteriteic amphiboles from veined mantle nodules (e.g., Figure 4b in Menzies et al. 1987).

Geochemical Modeling of KGC Supracrustal Magmatic Processes

In this section, the relationships between the LZ, the upper UZ and their parental liquid compositions, represented by the KGC chill rocks, and the Kamiskotia evolved basalts, respectively are established. This is followed by assimilation-fractional crystallization (AFC) modeling of the parental liquids using their trace element contents.

Chilled Compositions and Evolved Basalts as Parental Liquids: CMAS-Type Phase Diagrams

The phase relations and major element compositions of the KGC chill samples and the evolved basalts suggest that they are parental to LZ and upper UZ cumulates, respectively. The chill samples and evolved basalts are similar to abyssal basalts in their major element geochemistry, and as such can be represented in the CMAS-type diopside + olivine + plagioclase + silica tetrahedron of Walker et al. (1979). The chill samples plot in the olivine- (samples 34-37) or orthopyroxene- (sample 38) saturated fields at one bar (and fO_2 at QFM buffer) when projected from plagioclase, and in the plagioclase-saturated field when projected from silica (Figure 22). At atmospheric pressure, the average chill composition would crystallize plagioclase at An_{81-78} , orthopyroxene at En_{83-79} and olivine at For_{82-80} , at ratios of approximately 6:3:1 (McBirney 1985, after Nathan and Van Kirk 1978). These values and ratios compare favorably to LZ mineral compositions (Figure 5) and the average LZ mode (60% plagioclase, 12% orthopyroxene, 13% clinopyroxene, 16% olivine). The average Kamiskotia primitive and evolved basalts plot in the olivine, and the plagioclase and olivine fields, respectively (Figure 22). The primitive basalt average would crystallize olivine at For_{81} . The evolved basalt average would crystallize orthopyroxene at En_{69-65} and plagioclase at An_{59} at a ratio of 4:1 at one atmosphere (McBirney 1985, after Nathan and Van Kirk 1978), which closely approximates mineral compositions in the UZ (Figure 22).

Mass Balance Calculations Using the REE

If the major element geochemistry of the mixed magma and contact chill samples are candidates for a LZ parental liquid, then mass balance calculations using trace elements should provide supporting evidence for this relationship. One approach is to use mass

balance equations to estimate the REE content of the LZ cumulates using the average chill composition, where:

$$\text{Chill}_{\text{REE}} \times (\text{Bulk } K_D \times M_{\text{cum}} + M_{\text{TL}}) = \text{Cum}_{\text{REE}},$$

with $\text{Chill}_{\text{REE}}$ as the REE content in the average chill composition, Bulk K_D as the bulk distribution coefficient for the average cumulus mineralogy of the LZ, M_{cum} and M_{TL} as the estimated average mass of the cumulate phases and trapped liquid in the LZ cumulates, respectively and Cum_{REE} as the average REE content of the LZ cumulates. The trapped liquid composition is assumed to be the same as the chill composition, and density differences between liquids and cumulates are considered insignificant. Models that use mass balance calculations are highly dependent on the K_D s chosen. For these calculations, the values have been chosen conservatively, and are near the lowest of the range for basalt/mineral K_D s in the literature. Using trapped liquid contents of 0 (C_1) and 5 (C_2)%, the model LZ cumulates bracket the average LZ cumulate composition for the REE, including the positive Eu anomaly (Figure 23). They closely approximate the overall slope of the REE pattern, with $\text{La}_N/\text{Yb}_N = 1.01$ and 1.05 respectively, in comparison to the average LZ cumulate with $\text{La}_N/\text{Yb}_N = 0.99$.

A similar set of models have been calculated for the upper four UZ cumulates, using the average evolved basalt as a model parental liquid (Figure 23). For these models, trace amounts of cumulus apatite and Ti-magnetite are included in the average cumulus mineralogy, the K_D s for orthopyroxene and plagioclase are slightly higher (see Appendix III), and trapped liquid contents of 10 (C_1) and 25 (C_2)% are used (higher amounts of trapped liquid which may be applicable to UZ orthocumulates would require lower REE contents in the liquid: discussed below). As in the example above, the cumulate REE pattern (except for Nd) is bracketed by the model cumulate compositions. For C_1 and C_2 , $\text{La}_N/\text{Yb}_N = 1.15$ and 1.38 respectively in comparison to the average upper UZ cumulates with La_N/Yb_N

= 1.21. These mass balance calculations indicate that LZ cumulates and upper UZ cumulates may be derived from the KGC chill average and Kamiskotia evolved basalts respectively, consistent with the phase relationships using the major elements described above.

Assimilation - Fractional Crystallization Modeling

Variation diagrams are used to determine the nature of the supracrustal magma chamber processes, with trace elements, and TiO_2 and P_2O_5 contents plotted versus Zr on the abscissa (Figure 24a - f). Zirconium is used to monitor trace element enrichment or depletion as it behaves incompatibly in this system, forming coherent enrichment trends with other incompatible elements (Th, Hf, Y, Yb). TiO_2 , P_2O_5 , Th, La and Eu contents all increase systematically with increasing Zr.

Nickel (also MgO and Cr, not shown) is highest for the KGC chill rocks and decreases systematically through the evolved basalts, with increasing Zr (Figure 24a). However, the Ni contents of the primitive basalts are low. The low Ni contents may be explained by the fractionation of olivine or sulphide which lowers the Ni content of the liquid prior to extrusion of the primitive basalts. This would have to be followed by recharge of a primitive liquid during fractionation before the extrusion of evolved basalts, in order to explain the Ni and Zr contents in the evolved basalts, (e.g., sample 88-16 with 100 ppm Ni and 180 ppm Zr: Figure 24a: Barrie 1990). An alternative explanation, that the evolved basalts are derived from a separate source with different trace element characteristics, seems unlikely for several reasons. Firstly, the Kamiskotia primitive and evolved basalts are intimately interbedded on a scale of tens of metres at different stratigraphic levels, in several locations. Chemically-similar hypabyssal sills are found in the footwall of the evolved basalts on the Canadian Jamieson property. Secondly, there is clear evidence for the coexistence of magmas of intermediate (more fractionated equivalents of the

enriched basalts) and felsic composition in outcrop: trace element-enriched, quenched andesitic globules are found within ignimbritic rhyolites in southern Godfrey Township (Comba et al. 1986). One fragment of the enriched andesitic material has 54.5% SiO₂, 2.15% TiO₂, 0.36% P₂O₅, 290 ppm Zr and 120 ppm Y (unpublished data). And third, the mass balance modeling between the evolved basalts and upper zone gabbros described above infers a cogenetic relationship.

The trajectories marked on Figure 24 are for bulk assimilation-fractional crystallization (AFC) models. An average of roof rock granophyre samples from Hart (1984) is used for the material assimilated. This composition is used for two reasons. Firstly, there is field evidence for partial assimilation of large granophyric blocks in and adjacent to the UZ cumulates. Secondly, the granophyres are highly enriched in incompatible elements, and have relatively flat REE patterns with negative Eu anomalies. Other material considered to be assimilated are the mixed magma outcrop tonalites and granodiorites, and average Archean continental crust compositions (Taylor and McLennan 1985; Shaw et al. 1986). However, all of these have low incompatible element contents in comparison to the granophyre that would require greater amounts of assimilation (increasing silica and K₂O contents). Furthermore, their P₂O₅ contents are too low and their REE patterns are too steep which is not consistent with the high P₂O₅ contents and flat REE patterns of the Kamiskotia basaltic andesites. Ratios of assimilation/fractionation used in the models range from 0 to 0.5. Generally, assimilation/fractional crystallization rates for mafic-ultramafic systems range from 0.01 for parts of the Kiglapait intrusion, which is hosted in refractory gneisses and anorthosites (DePaolo 1981a) to 0.5 for komatiites in supracrustal environments (Sparks 1986). Rates of 0.1 to 0.3 are considered normal for supracrustal tholeiitic intrusions (DePaolo 1985; Sharpe 1986; Sparks 1986).

The AFC models with ratios of 0.1 to 0.3 work well for some of the elements, but they do not approximate the enrichment for P₂O₅. Assimilation of the granophyric rocks with low P₂O₅ contents depresses the enrichment due to fractionation. Furthermore even

small amounts of granophyre assimilation will cause significant increases in silica content which are not observed.

Bulk fractionation of 70 to 80% provides a reasonable approximation for these trends, particularly Ni, P_2O_5 and Th (Figure 24a, c, d). Deviations from the data may be due to variation in the bulk K_D s during fractionation. For TiO_2 , an increase in the bulk K_D from 0.2 to 0.4 after 40% fractionation, provides a close approximation of the data (Figure 24b). This may reflect the onset of Fe-Ti oxide fractionation. Higher bulk K_D s provide a better fit for La and Eu also (Figure 24e). Higher K_D s for the La and Eu could reflect minor apatite fractionation and/or greater partitioning into plagioclase with decreasing An content (Schnetzler and Philpotts 1970). If higher bulk K_D s for the REE are necessary, then the average evolved basalt composition may be too enriched to be parental to the upper UZ cumulates (considering the mass balance calculations described above), and the upper UZ cumulates may have been derived from a composition intermediate between the primitive and evolved basalts.

Fractionation also explains decreasing Mg numbers and the relatively constant silica contents. Bulk fractionation of 70 to 80% from the KGC average chill composition causes Mg numbers to decrease from 56 to 29 to 30 (Nathan and Van Kirk 1978), within the range for the evolved basalts (Hart 1984; Barrie 1990). Silica contents are enriched only slightly during fractionation, in contrast to the five-fold enrichment in Zr. In general terms, if a bulk K_D for silica is estimated from $SiO_2(LZ)$ divided by $SiO_2(chill)$ at slightly less than unity, then only slight silica enrichment would result, even at high levels of fractionation. This is a well-known characteristic of tholeiitic fractionation trends, where Fe enrichment is accompanied by essentially no silica enrichment in the liquid prior to significant Fe-Ti oxide fractionation (e.g., Skaergaard Intrusion: Wager 1960; Hunter and Sparks 1987).

Other geochemical models for trace elements that may pertain to the Kamiskotia magma system, but not considered in detail here, are fractionation-recharge and - discharge models. In fractionation-recharge models, incompatible element enrichment and compatible

element depletion are enhanced in comparison to FC models due to higher degrees of cumulates removed for a given volume of liquid in the system. Thus, their trajectories would follow the fractionation-only trajectories in Figure 24 but extend to greater degrees of enrichment for a given volume of residual liquid in the system (O'Hara 1977). If discharge from the system is incorporated into the model, then enrichment or depletion may eventually reach a steady state, as originally envisioned for mid-ocean ridge systems by O'Hara (1977). Such a model could account for greater variation in incompatible element ratios (e.g., Zr/Hf), if the amount of magma discharged is small in comparison to the amount fractionated (O'Hara 1977).

Trace element enrichment by two-liquid chemical diffusion and thermal (Soret) diffusion should also be considered at Kamiskotia, as there is geologic and textural evidence for the coexistence of mafic and felsic liquids in both intrusive and metavolcanic rocks. Basalt flows and rhyolite tuffs interdigitate on a scale of metres in several locations, and quenched andesitic enclaves are found within rhyolite crystal lithic tuffs at one location (described in Geology section).

Two-liquid chemical diffusion models predict that the LILE prefer more polymerized acidic liquids, and just the opposite for the HFS elements (Watson 1976; Ryerson and Hess 1978). However, the entire Kamiskotia suite (excepting one extremely fractionated rhyodacite) shows normal enrichment trends with increasing SiO₂ content, which would not be the case if chemical diffusion were a dominant process. Thermal diffusion models predict that the LILE will generally partition into the hotter silicate liquid whereas most HFS elements will partition in the opposite direction (Leshner 1986). For MORB compositions, the REE are enriched by a factor of 2.5 under temperature gradients of 50-80°C (at 1 GPa, 1380°C to 1535°C: Leshner 1986). This suggests that in addition to fractionation, thermal diffusion may have been an operative process in the formation of the Kamiskotia evolved basalts.

Nd Isotope Signatures

In a related study of Kamiskotia area rocks, Barrie and Shirey (in prep.) investigated the geochemistry and Nd-Sr isotope systematics of several rock suites. They used geochemistry and Nd isotopic compositions to characterize mantle and crustal sources, and to provide constraints on petrogenetic models for tholeiitic, calc-alkalic and lamprophyric suites in the Kamiskotia area. One of the purposes of the study was to identify and characterize potentially distinct mantle and crustal sources for the supracrustal rocks using whole rock geochemistry and Nd isotopic compositions. Previous trace element and radiogenic isotope studies for Abitibi mantle-derived rocks have indicated that the mantle was variably depleted in Sm/Nd, Rb/Sr, and U/Pb with respect to a reservoir with chondritic ratios for these elements. When considered on a subprovince scale, mantle-derived Abitibi tholeiitic and komatiitic intrusions appear uniformly depleted in their Nd and Sr isotopic compositions (e.g., Machado et al. 1986). However, other investigations have detected different isotopic signatures (Basu et al. 1984), even within 'cogenetic' suites (Cattell et al. 1984), and mantle heterogeneities have been found in Archean rocks elsewhere in the southern Superior Province (Shirey and Hanson 1986). Crustally-derived Abitibi rocks show primitive and slightly enriched Pb isotopic signatures (see Gariepy and Allegre 1985; for review), and isotopic heterogeneities have been found in crustally-derived rocks in northwestern Ontario (e.g., Hanson et al. 1971; Morrison et al. 1985; Shirey and Hanson 1986; Smith et al. 1987).

The KGC and KVC are represented by two ferroan gabbros, a dacitic granophyre, a rhyolite and a basalt. The suite is characterized by relatively flat REE patterns and a restricted range of $\epsilon_{Nd}(t)$ values (defined as the deviation in parts per 10,000 from a chondritic uniform reservoir of the same age, with ages from U-Pb data) from +2.2 to +2.6. This is essentially identical to the best estimates of the MORB-like, isotopically depleted Abitibi mantle at 2.7 Ga (Dupre et al. 1984; Machado et al. 1986).

Three basalts of the Lower volcanic suite have flat to slightly light REE-enriched REE patterns, with Ce/Yb ranging from 2.3 to 10.6, and $\epsilon_{Nd}(t)$ values ranging from +1.7 to +3.0.

Lamprophyric rocks in the Kamiskotia are represented by dikes in the Montcalm gabbroic complex, located 25 km west of the KGC; and the Bristol Township lamprophyre suite (see Figure 4 for location). The lamprophyre samples are characterized by high Th, light REE and P_2O_5 values, and steep REE patterns. Both Montcalm lamprophyre dikes have nearly identical $\epsilon_{Nd}(t)$ values of +2.8 and +2.5. A garnetite sample from the Bristol Township lamprophyre suite has a whole rock $\epsilon_{Nd}(t)$ value of +1.0, slightly isotopically enriched in comparison to the Abitibi mantle at 2.7 Ga.

Petrogenesis of the KGC Gabbros and KVC Basalts

The KGC average chill composition is compared to N-type MORB and Alexo komatiites (located 140 km east of the Kamiskotia area, within the Abitibi Subprovince) on a primitive mantle-normalized diagram (Figure 25). The elements are ordered according to their compatibility with primitive mantle peridotite, with the most compatible elements to the right. Generally, a melt formed from single stage partial melting would curve downward toward the compatible elements to the right and would be flat toward the incompatible elements at the left. A two stage model is required for the abundances of N-MORB, which has a concave-down pattern. The first stage has approximately 1.5% melting that depletes the mantle in the most incompatible elements, and the second stage has 8 to 10% partial melting of the first stage residue (Hofmann 1988). For Alexo, Barnes (1984) and Brugmann (1985) have noted that the elemental abundances (except K_2O) also correspond to a two stage model, with a higher degree of melting during the second stage (up to 40%).

The KGC chill average has compatible element concentrations nearly identical to N-MORB; however the incompatible elements exhibit a flat pattern (except for K_2O , Rb and Ba: see below). The most straightforward explanation is a derivation by single-stage partial

melting of a chemically primitive mantle. A first order approximation for the minimum degree of single stage partial melting can be determined using a simplified equation for bulk partial melting: $F = C_0/C_l$, where F is the fraction of melt, C_0 is concentration in the primitive mantle before melting and C_l is the concentration of the element in the melt. Using the 13 elements in Figure 26 from La to Lu, an average value for the KGC chill average of 15.9% (1 sigma = 2.6%) partial melting is required; excluding P_2O_5 , TiO_2 and Eu, 14.7% ($n=10$, 1 sigma = 1.2%) partial melting is required.

This approximation does not account for fractionation that must have occurred, after mantle melting and prior to emplacement. Negative Eu anomalies and liquidus phase relationships (Nathan and Van Kirk 1978) for the KGC chill samples imply a previous history of plagioclase fractionation. Furthermore, primitive magmas must be in equilibrium with mantle olivine compositions of Fo₈₆₋₉₀ (Green 1970). Given the olivine - liquid $K_D^{Fe^{2+}/Mg}$ of 0.30 - 0.33 (Roeder and Emslie 1970; Roeder 1974; Longhi et al. 1978), the minimum Mg number for primitive magmas are 68-75. Using an FeO/FeO_{total} of 0.9 which is appropriate for mantle-derived magmas, the KGC chill average would have an Mg number of 51. This is significantly lower than the minimum required for a primitive magma, implying that olivine and possibly pyroxene fractionation occurred after mantle extraction in addition to plagioclase. A reasonable estimate for the amount of fractionation can be made by combining the average olivine-bearing adcumulate composition (samples 1-4: average Mg number = 81, using $FeO/FeO_{total}=0.9$) with the KGC chill average to obtain a liquid in equilibrium with mantle olivine. A 1:1 mix of these components produces a composition with an Mg number of 70. This liquid would have 27.9 wt.% normative olivine (24.7 mol.%) and a flat REE pattern at 9-10x chondrites. The degree of partial melting of primitive mantle required to produce this composition would be 34% (using the partial melting equation and the REE without Tb). The normative olivine content of this liquid and the degree of partial melting can be used to estimate the temperature and pressure of anhydrous melting from a

spinel lherzolite primitive mantle composition. This corresponds to melting at 17 kbars pressure and 1550°C (Figure 9: Jacques and Green 1980).

An alternative explanation is that like the Alexo komatiites and N-MORB, the KGC chill was derived from a mantle source that previously had been depleted in the most incompatible elements but has subsequently undergone enrichment to form, perhaps fortuitously, the flat pattern on the primitive mantle-normalized diagram (Figure 25). Selective enrichment would be due to either diffusion from the adjacent felsic liquids in the mixed magma and contact chill outcrops (Watson 1976), which as previously discussed seems unlikely, or by selective contamination en route through the lower crust as suggested by Watson (1982). Selective contamination processes cannot be ruled out, although at present there is no supporting experimental evidence for selective contamination involving HFS elements in mafic liquids.

Petrogenesis of KVC rhyolites

Considering the exposures of coeval mafic and felsic, intrusive and extrusive rocks, the Kamiskotia area is ideal for investigating the petrogenesis of high silica rhyolites in bimodal (basalt-andesite and rhyolite) volcanic fields. For this discussion, high silica rhyolites are defined as a suite of rhyolites having greater than 73 wt.% SiO₂, high HFS and REE contents, flat REE patterns with negative Eu anomalies, and Rb/Sr ratios greater than unity.

The petrogenesis of high silica rhyolites in bimodal volcanic fields like at Kamiskotia is a contentious issue. There are two main hypotheses. The first hypothesis, proposed particularly for high silica rhyolites of the Basin and Range Province, U.S.A., is that high silica rhyolites represent partial melts of a felsic granulite source in the lower or middle crust, and have undergone little or no modification by supracrustal magmatic processes (e.g., Doe et al. 1982; Christiansen et al. 1986). The second hypothesis is that high silica rhyolites are the product of high degrees of fractional crystallization from a MORB-like, mantle partial

melt, possibly accompanied by significant assimilation of roof rock partial melts, and by chemical and/or thermal diffusion-related enrichment processes within supracrustal magma chambers. Variations of this hypothesis have been proposed by numerous authors for the bimodal volcanic fields of eastern Iceland (e.g., Wood 1978; Macdonald et al. 1987; and by Hart 1984) for the Kamiskotia rhyolites. A third hypothesis, involving the separation of an immiscible, siliceous liquid from tholeiitic parent, has been proposed for rhyolites similar to high silica rhyolites in the Uchi-Confederation Lake area of the Superior Province (Thurston and Fryer 1983). Experimental studies (e.g., Watson 1976) on coexisting basaltic and silicic liquids indicate that incompatible elements will partition strongly into the mafic liquid, a phenomena that is not observed at Uchi - Confederation Lakes in particular, or in bimodal volcanic suites generally. Thus liquid immiscibility is not considered in this discussion.

The first hypothesis is proposed most elegantly for a variety high silica rhyolites, the topaz rhyolites of the Basin and Range Province (e.g. Christiansen et al. 1986). Topaz rhyolites are characterized by high silica contents, high F, Rb, U and Th contents, low Sr, Ba and Eu contents, and flat REE patterns with strong negative Eu anomalies. Christiansen et al. (1986) proposed that siliceous magmas with high F contents are the result of partial melting of a metamorphosed felsic protolith, due to the passage of hot mafic magmas through the lower or middle crust, with the decomposition of small amounts of F-rich biotite to give high F contents. The siliceous melts then migrated into supracrustal magma chambers, and extruded along with the contemporaneous mafic magmas. A similar hypothesis has been proposed for a suite of Archean high silica rhyolites, those that overlie the Bad Vermilion Anorthosite Complex in the Rainy River area of the southern Superior Province. Shirey (1984) postulated that partial melting of a tonalitic source with a relatively flat and elevated REE pattern, at either shallow or deep levels, as the most likely mechanism to generate these rhyolites.

The second hypothesis is a combination of several shallow level magma chamber processes: fractionation, assimilation of roof rock partial melts, and thermal/chemical

diffusion. Fractionation of olivine, two pyroxenes, plagioclase and an Fe-Ti oxide phase from a Kilauea basalt (Mg number = 53), under hydrous conditions at 1 kbar, has been shown to generate rhyolites nearly identical in major element geochemistry to the high silica rhyolites of eastern Iceland (Spulber and Rutherford 1983). These experiments required 90% fractionation of these phases in cotectic proportions. A similar, high degree of fractionation would account for incompatible element enrichment also. For example, using a reasonable bulk partition for Ce of 0.2 over the range of magma compositions, a tenfold enrichment from 12 ppm Ce (typical MORB) to 120 ppm Ce (average KGC rhyolites from Hart 1984) requires greater than 90% equilibrium crystallization (using the equation: $C_l/C_o = 1/K_D (1-F)$, with C_l the concentration in liquid remaining, C_o the concentration in the liquid initially, K_D the bulk distribution coefficient, and F the fraction of system that is liquid).

If fractionation is the primary mechanism to produce the high silica rhyolites, then a spectrum of volcanic rock compositions would be expected. This is clearly not the case for most of the high silica rhyolites which are found in bimodal volcanic fields, including at Kamiskotia. One explanation for the compositional gap found in the bimodal fields may be provided in a petrologic and theoretical study of the Pleistocene to Recent Medicine Lake Volcano, California (Groves and Donnelly-Nolan 1986). At Medicine Lake, a compositional gap between andesites and rhyolites is explained by a three stage fractionation process from a basaltic precursor. The first phases to fractionate were olivine, plagioclase and augite, and then olivine was replaced by orthopyroxene and amphibole. This led to an andesitic liquid, enriched in iron, titanium and phosphorous (and incompatible elements). Fractionation continued with apatite and magnetite crystallization joining the plagioclase, augite and amphibole, producing a liquid of rhyolite composition. Groves and Donnelly-Nolan (1986) noted that during magnetite and apatite crystallization, fractionation proceeds over a limited temperature interval (i.e., the liquidus surfaces have shallow slopes). Considering the rapid change in liquid composition over such a restricted temperature interval, the possibility of preserving an andesitic to rhyolitic liquid is diminished.

Fractionation from andesitic to rhyolitic liquids over a restricted temperature interval should be accompanied by a rapid increase in volatile contents. High volatile contents can aid chemical/thermal diffusion processes, which can lead to incompatible element enrichment in siliceous systems, provided that the magma chamber is compositionally stratified and convecting (e.g., Hildreth 1981).

Hart (1984) favored partial melting over fractionation as the principal mechanism to produce the KVC rhyolites and the KGC granophyres. Using a fractional melting model, and partition coefficients for andesitic liquids, Hart found that the REE concentrations in model rhyolites were generally similar to a ten percent partial melt of a KVC primitive basalt composition, or a thirty percent melting of a KVC evolved basalt. In both cases, however, the resulting model rhyolite had significantly higher Ce/Yb ratios than the average KVC rhyolites, and these models may not adequately represent the Kamiskotia system.

Radiogenic isotope studies can provide a way to distinguish between the mantle or crustal origins for high silica rhyolites. The ideal locations for differentiating between mantle and crustal sources are where old sialic basement is present, such as across much of the Basin and Range Province. In a study of the Yellowstone bimodal volcanic field, Doe et al. (1982) found that the initial Pb and Sr isotope ratios of basalts were significantly less radiogenic than the coeval rhyolites. They interpreted the basalts as mantle-derived, and the high silica rhyolites as partial melts of a siliceous lower crust, based principally on the Sr isotope ratios. Halliday et al. (1989) obtained nearly the opposite results for the young (1.2 to 0.8 Ma) basalts and high silica rhyolites of the Long Valley Caldera, California. From a detailed Nd-Sr-Pb isotopic study, they found that these rhyolites are relatively unradiogenic, and identical to coeval, mantle-derived basalts. Other studies have found a spectrum of isotopic ratios for high silica rhyolites, particularly in the Rb-Sr system (e.g., Bacon et al. 1984; Christiansen et al. 1986). (It is noted that a significant problem with the high Rb/Sr ratios that characterize high silica rhyolites, and topaz rhyolites in particular, is that it is very

difficult to obtain accurate initial ratios, due to long, age-dependent extrapolations on isochron diagrams.)

There is some evidence of an older sialic basement in the Kamiskotia area, from an inherited zircon fraction in granitoid B, and from a regional Sm-Nd isotope study in the Kamiskotia - Montcalm area (Barrie 1990; Barrie and Davis 1990). However, geochemical and radiogenic isotope studies have shown that there is no evidence for assimilation or contamination of an older sialic basement for any part of the KGC or the KVC (Barrie 1990). This provides supporting evidence to the field and geochronological data that support a coeval, cogenetic relationship between mafic and felsic, intrusive and extrusive rocks in the Kamiskotia area.

To summarize, there are several lines of evidence to support a coeval and cogenetic relationship between the KVC high silica rhyolites and the KGC, with no involvement of an older crustal component: 1) coeval, high precision U-Pb ages for KGC and KVC rhyolite samples; 2) abundant field evidence for the coeval nature of the KVC rhyolites and basalts; 3) geochemical evidence for a cogenetic relationship between the KGC and KVC basalts; and 4) initial Nd isotope ratios for KGC and KVC rocks that are compatible with a derivation from a depleted mantle source, with no evidence for contamination from any enriched crustal component. Fractionation-dominated magma chamber processes at low pressures adequately explain the composition of the KVC rhyolites, although more rigorous modeling is necessary to test this hypothesis. Partial melting of KVC basalts, and chemical/thermal diffusion processes within the Kamiskotia magma chamber may have played minor roles in the generation of the KVC rhyolites.

Petrogenesis of Bristol Township Lamprophyre suite

Perhaps the closest analogs to the Bristol Township lamprophyre suite are ultramafic lamprophyres, and particularly those with alkalitic affinities. Ultramafic lamprophyres (UML)

are characterized by their low SiO₂, and high K, Sr, Ba, REE, Mg, Cr, Ni, and volatile contents; they are distinguished from other lamprophyre types by an unusual mineralogy, including olivine (Fo₉₂₋₇₈), phlogopite (rich in Ti, Fe³⁺, Ba or F) and sodic amphiboles as phenocrysts, and carbonates, feldspathoids, Ca-Fe-Ti-Zr garnets in groundmasses (Rock 1986; 1987). Allikites are a carbonate-rich variety of UMLs. Classic examples of allikites are found at Aillik Bay, Labrador (Malpas et al. 1986); and associated with carbonatites at Oka, Quebec (Treiman 1982) and Magnet Cove, Arkansas (e.g., Steele and Wagner 1979). Ultramafic lamprophyres are rare: the only other UML rocks documented in the Superior Province are the 1650 Ma McKellar Harbour dikes near Marathon, Ontario, (Platt et al. 1983), and biotite-olivine-carbonate dikes at the northern end of Lake Nipigon, believed to be circa 1500 Ma (Sutcliffe 1988).

Many UMLs are believed to represent primary, mantle-derived magmas, generated at depths of 100 to 150 km, probably derived from metasomatized, LREE-enriched mantle material (e.g., Malpas et al. 1986; Rock 1986). The primary melts may have been extensively modified by fractionation or interaction with alkali- and volatile-rich fluids (Rock 1986). It is noted that the Bristol Township lamprophyre suite differs from most UML in that it is not known to have Mg-rich olivine, high Ni and Cr contents, and it more enriched in the HFS and REE than most UML (Rock 1986). In these aspects, it is similar to zones of P- and F-rich metasomatism along the DPFZ, which have been compared with fenitization associated with volatile-rich, mantle-derived alkaline magmas (King and Kerrich 1987). Additionally, it is chemically and mineralogically similar to parts of the Oka carbonatite, Quebec, believed to represent CO₂-rich, upper mantle partial melts (Eby 1975).

A slightly enriched Nd isotopic signature of +1.0 (Barrie 1990) must be accounted for when considering the ultimate source of the Bristol Township lamprophyre suite. One possibility is that the suite was derived by magmatic or metasomatic processes, from a crustal material 200 Ma older than the majority of rocks in the southern Abitibi Subprovince. This would require a very efficient scavenging of REE from the crust into a melt/fluid in order to

account for the incompatible element enrichment of the lamprophyre suite. Two processes that may account for this are CO₂ fluid streaming through the lower crust during granulite facies metamorphism (Newton 1987); and fluxing by an H₂O-rich fluid derived from subducting oceanic slabs (Sorensen and Grossman 1989). It is noted that there are no known rock suites associated with subduction zones that are as enriched in the light REE as the Bristol Township lamprophyre suite. A second possibility is that the lamprophyre suite was derived from a part of the mantle that had undergone significant light REE enrichment, approximately 200 Ma prior to the separation and emplacement of the lamprophyre suite (Barrie 1990). At present, it is difficult to distinguish between isotopically enriched mantle, or lower crustal sources for the lamprophyre suite.

ECONOMIC GEOLOGY

Volcanogenic Cu-Zn±Au±Ag mines and deposits, mesothermal Au deposits, magmatic Ni-Cu occurrences and one REE occurrence are present in the Kamiskotia area (Figure 4). This section describes the most economically significant of these mineral concentrations. New assays and chemical data are provided for several Ni-Cu occurrences within the KGC, and a previously undocumented REE occurrence in the Bristol Township lamprophyre suite.

VOLCANOGENIC Cu-Zn DEPOSITS

Four massive sulphide deposits hosted in the KVC have been mined for their Cu, Zn, ±Ag ±Au contents: the Kam-Kotia mine, the Jameland mine, the Canadian Jamieson mine, and the Genex mine (Figure 4). In general terms, all four orebodies have the following characteristics: they are composed of several smaller lenses or masses of sulphide material; the lenses or masses are generally within a restricted stratigraphic interval (< 150 m) at each mine site; their host metavolcanic rocks are broadly correlatable from deposit to deposit;

their host rocks are predominantly mafic metavolcanic rocks, with subordinate felsic tuffs located on the mine properties; and they exhibit strong chlorite alteration in the stratigraphic footwall, and a more widespread sericite alteration pattern. The sulphide minerals for the deposits are pyrite, pyrrhotite, chalcopyrite, sphalerite, and minor magnetite; trace galena has been reported for the Kam-Kotia mine (Pyke and Middleton 1970). Interestingly, incompatible element-enriched, basaltic andesite and andesite flows and sills, generally uncommon in the KVC, are present at each mine property, in close proximity to the ore. These rocks apparently postdate ore formation, as they are generally less-altered than other KVC rocks, and they cut the other metavolcanic rocks and the orebodies locally. The geochemistry of the basaltic andesites and andesites are described above (see Geochemistry section).

Kam-Kotia and Jameland Mines

The Kam-Kotia mine is located in eastern Robb Township (Figure 26; #1 in Figure 4). Pyritized, rhyolitic breccias and tuffs are the immediate host rocks to the mine, and these are found within a sequence of predominantly mafic metavolcanic rocks which are north-facing, have a strike of 315° , and dip 75°N . Massive pyrite lenses up to 2 m in thickness are exposed at the southern wall of the open pit. They are found within foliated mafic metavolcanic rocks which have been subjected to strong chlorite and carbonate alteration. Spherulitic rhyolite with apparent flow banding occurs in a prominent outcrop within the southern half of the open pit. The following description of the orebody is paraphrased from Somerville (1967), Pyke and Middleton (1970) and Binney (Falconbridge Limited, personal communication, 1990):

The Kam-Kotia orebody consisted of seven lenses or irregular-shaped masses. The mineralized zone appeared to be steeply dipping. It consisted of six copper-rich lenses

and one zinc-rich lens that had a strike slightly more westerly than the host rocks, and a plunge at 30°NW. The lower lenses were predominantly massive sulphide, whereas the upper lenses contained more stringer zone material (quartz - sulphide stockwork, with 10 to 25% sulphide). In general, copper was concentrated in closer to the surface and zinc at depth. Chloritic and sericitic alteration halos are present and surrounded the ore lenses. Andesite dike and sill-like masses apparently cut both ore lenses and altered metavolcanic rocks (Figure 26). The andesites are fine-grained, and unmineralized, with well-defined chilled margins. Mineralization predated andesite dike-sill emplacement and may have influenced their emplacement.

The Jameland mine is in Jamieson Township, 1.2 km to the southeast and along strike with the Kam-Kotia mine (#2 in Figure 4). The following description of the orebody is taken from Pyke and Middleton (1970) and Middleton (1973c):

The deposit was situated within sheared, chloritized and brecciated basalts, andesites and rhyolite tuffs. The deposit consisted of ten lenses or irregular-shaped masses, and in its entirety plunged 30 to 35° to the southeast. At the west end, the orebody was a single lens with a horizontal width of 15 m, whereas 300 m to the east, near the center of the zone, numerous sulphide masses are distributed over a horizontal width of 100 m. As with the Kam-Kotia mine, stringer-type copper-rich ore was confined to the upper lenses and massive zinc-rich ore to the lower lenses. A zone of massive pyrite, without significant base metal content, occurs between the Kam-Kotia and Jameland mines.

Ore from the Kam-Kotia mine was recovered briefly during World War II (186,000 tons (169,000 tonnes) recovered). The mine produced again from 1961-1972. During the second period of mining, 6,436,000 tons (5,840,000 tonnes) were recovered (including minor

production from the Jameland deposit), with average grades of 1.1% Cu, 1.2% Zn, and 0.1 oz./ton Ag (totals of 143,351,665 lbs (6.5 by 10⁶ kg), 156,000,000 lbs, (7.1 by 10⁶ kg) and 663,136 oz. (20600 kg), respectively; also 5604 oz. (174 kg) Au: Canadian Mines Handbooks 1961-1974).

Canadian Jamieson Mine

The Canadian Jamieson mine is located in northwest Godfrey Township (Figure 27, and #3 in Figure 4). The orebody was situated within predominantly mafic metavolcanic rocks that face to the northeast, with lesser rhyolite tuffs and flows(?). Basaltic andesites are found 100 m south of the mill foundation (Figure 27). Shear zones up to 1 m wide and ductile deformation in the metavolcanic rocks are present on the northeast outcrops. Deformation is probably related to a major fault zone located 100 m to the northwest (at 130-140°, near-vertical: Figure 27), which postdated KVC formation and massive sulphide mineralization. A description of the minesite stratigraphy and of the orebody is given here, based on the descriptions of Comba et al. (1986), and P. Binney and B. Filbey (Falconbridge Limited, personal communication, 1985-1990):

From surface exposures, the base of the stratigraphic succession in Figure 27 consists of pillowed mafic flows that are intercalated with thin felsic ash beds, a mafic fragmental unit and several, thin massive mafic flows or sills. Felsic lapilli tuff overlies the mafic metavolcanic sequence, which is, in turn, overlain by a sulphidic, iron carbonatized, mafic fragmental unit. The base of the south ore zone appears to comprise a pyritic, cherty exhalite with stringer and colloform pyrite in a black, siliceous matrix. The south ore zone is overlain by bleached and altered pillowed mafic flows. These are in turn overlain by a spherulitic flow banded rhyolite and lapilli tuff or flow breccia unit. All felsic units are sodium-depleted, iron-carbonatized,

and strongly foliated. A sulphide clast-bearing, iron-carbonatized, mafic fragmental unit overlies the felsic metavolcanic rocks and forms an extension along strike to the north ore zone. These rocks are capped by massive mafic flows, which have relatively little alteration.

The orebody consisted of three strata-bound sulphide lenses; the south, central and north zones. From observations made during underground mining, hanging all rocks were usually barren, but footwall rocks were pyritic and locally contained sufficient chalcopyrite to be mined. Zoning was recognized within individual orebodies, which were copper-rich near their stratigraphic bases and more zinc-rich near their tops. Footwall rocks have been altered to sericite and chlorite.

The Canadian Jamieson mine operated from 1966 to 1971, producing 816,000 tons (740,000 tonnes) of ore averaging 2.4% Cu and 4.2% Zn (Northern Miner Handbooks, 1965-1974).

Genex Mine

The Genex mine is located in central Godfrey Township (Figure 28, and #3 in Figure 4). The mine is located near a contact between KVC felsic breccia and mafic units: massive and pillowed mafic metavolcanic rocks and mafic sills. The stratigraphy here is near-vertical and faces to the east. Legault (1985) studied the geology and alteration associated with the Genex deposit. He compiled the results of surface geological mapping with mine plans from Genex Mines Limited (Figures 29, 30, 31). The following description is based largely on the work of Middleton (1975) and Legault (1985):

Four synvolcanic felsic intrusions are present on the Genex property. They have a general east-trend, perpendicular to stratigraphy, and range in apparent

thickness from 40 m to 150 m. They may represent synvolcanic rhyolite flow domes. They taper to the east, and may be gradational with larger granophyric felsic intrusive masses to the west. The synvolcanic felsic intrusions are physically and texturally similar to nearby massive felsic metavolcanic rocks: both have spherulitic and flow-banded margins, quartz and feldspar phenocrysts, local brecciated areas with flow banded clasts, and xenoliths of mafic metavolcanic material. Three of the four synvolcanic masses are strongly altered and mineralized, and one was the host rock for some of the "H" stringer ore zone (described below). Large sill-like mafic bodies are found stratigraphically below and above the Genex deposit (Figure 29). These rocks are possible feeder dikes in the area. Some of these intrusions apparently truncate patterns of alteration and mineralization that cross stratigraphy, but the intrusions are not visibly altered or mineralized.

The deposit is divided into several zones, with the A, C, and H zones comprising the most economic parts of the deposit (Figures 29, 30, 31). The A and H zones represent copper-rich, quartz stockwork stringer mineralization. They have a strike that crosses stratigraphy, and are situated within in felsic metavolcanic breccia and massive to pillowed mafic metavolcanic rocks. The C zone is conformable, and is located at a contact between a mafic pillow breccia and a massive mafic intrusion.

Production at Genex was minor, with approximately 120,000 tons (109,000 tonnes) of ore at 2.2% Cu recovered (Middleton 1975). Diamond drilling and underground exploration indicated reserves of 133,000 tons (121,000 tonnes) with an average grade of 2.2% Cu, with resources at 385,000 tons (349,000 tonnes) of 1.7% Cu by one estimate (Middleton 1975).

MESOTHERMAL Au DEPOSITS

Holmer Property

The Holmer gold property is located in southwest Bristol Township (Figure 32, and #6 in Figure 4), within highly deformed rocks of the DFPZ, at a contact between mafic metavolcanic rocks of the Lower volcanic suite, and metasedimentary rocks. The Bristol Township lamprophyre suite occurs on the mine property, 50 southwest of the main shaft, away from the gold mineralization (see Geochemistry section for description, and sample 89-5 in Table 8 for geochemical analysis).

The rocks on the property are variably deformed. Primary volcanic and sedimentary textures are observed in less-deformed rocks in several locations. Hyaloclastite textures are preserved 25 m southwest of the main shaft. Graded layering, with coarse sand-sized to silt-sized grains are present in turbiditic metagreywacke, 60 m northeast of the main shaft. Facing directions are difficult to determine in the metagreywacke, due to chlorite alteration concentrated along fractures that parallel bedding. One bed, in which coarse to fine gradation is observed, strikes 105° , dips 80°S and faces to the south. Elsewhere, the rocks are moderately to highly deformed. The most intense deformation is located in the nose of a west-plunging fold hinge, manifested as rods or pencils ($L > S$) of carbonatized metasedimentary rock, 25 m northeast of the main shaft.

Mineralization is found in quartz + tourmaline + sulphide (pyrite, chalcopyrite, and minor arsenopyrite, galena and sphalerite) veins and the adjacent, tourmalinized, carbonatized and locally silicified metasedimentary and metavolcanic rocks.

The property has been investigated intermittently since 1911, with no known production. Recent estimates of probable reserves range from 720,000 tons (650,000 tonnes) at 0.124 oz. (3.85 g)/ton Au to 865,000 tons (785,000 tonnes) at 0.08 oz. (2.5 g)/ton Au

(Northern Miner Handbook, 1987-88; S. Fumerton, Chevron Resources Limited, personal communication, 1989).

Au-REE Occurrence: The Croxall Property

The Croxall Au-REE property (also known as the Rusk property) is located 1.5 km to the southwest of the Holmer property (Figure 33, and #9 in Figure 4). The main showing has excellent exposure of the Bristol Township lamprophyre suite along three trenches and adjacent outcrops. (See Geology and Geochemistry sections for detailed descriptions of the lamprophyre suite, and Table 8 for geochemical analyses). Lower volcanic suite metabasalts are present on the northwestern part of the main showing, and continue to the west for several kilometres. Excellent exposures of hyaloclastitic metabasalts are located 0.7 km to the west of the main showing. Metasedimentary rocks are present to the east, and a granite porphyry stock is located to the south-southeast (Ferguson 1957a).

Several of the Bristol Township lamprophyre suite samples are highly enriched in the REE and Zr (Table 8). Sample 88-5, an altered garnetite dike, has total REE contents of 0.43 wt.% RE₂O₃, and 3070 ppm Zr. Preliminary microprobe work by D. Wark (Rensselaer Polytechnical Institute, Troy, New York) indicates that the majority of the REE in garnetite rocks are located in fluorine-rich apatites and epidotes, which have up to 9 and 13 wt.% RE₂O₃, respectively. In general the melanite garnets have relatively low REE, with up to 0.15 wt.% Sm₂O₃ (only measurable REE), and they average approximately 3000 ppm Zr. These rocks are possibly the most REE-enriched rocks of Archean age in the entire Superior Province.

The property has been worked as a gold prospect intermittently since 1941, including trenching, numerous blast pits, and 39 diamond drill holes. Gold mineralization occurs in two areas. One area is a minor, stockwork quartz-sulphide vein network, within the north-northeast part of the granite porphyry (J.C. Pederson, Highwood Resources unpublished

report). The second gold occurrence is the main showing, where hand samples up to 0.18 oz. (5.5 g)/ton Au have been reported, from quartz-carbonate-sulphide veins (LeBaron 1985).

De Santis Property

The De Santis property is located in southwest Turnbull Township (Figure 34, and #10 in Figure 4). The property lies entirely within the granitoid A region, and both KGC gabbros and quartz-feldspar porphyry (granitoid A) are exposed. A north-trending Matachewan dike cuts the gabbro and quartz-feldspar porphyry. The dike is parallel and adjacent to the main quartz vein, which is up to 1.5 m in width at the surface, and also cuts the gabbro and quartz-feldspar porphyry. The vein is generally barren of sulphide, but contains up to 1.5% sulphides (pyrite, chalcopyrite) locally. The gabbro and quartz-feldspar are locally deformed, with a well-developed flattening fabric parallel to the margins of vein, near the shaft. Mineral lineations (alignment of actinolite-chlorite) on altered gabbro surfaces are nearly horizontal, plunging 10° to the north. It is noted that the vein attitude, along the trend of the adjacent Matachewan dike, is unusual: there are few quartz veins having this orientation and with gold mineralization in the entire Porcupine District. (The Lally Au property, located in east-central Turnbull Township, within the eastern margin of the granitoid A region, is parallel to a north-trending Matachewan dike also: Middleton 1975). Considering the structural history of the Kamiskotia area, the vein may occupy a dilational fracture that formed during a north-south compressional regime, the same stress regime that was responsible for the predominant east-west fabric through the KVC and the granitoid A region at approximately 2700 Ma to 2685 Ma (Figures 6, 7; see Structural geology section).

The property was discovered in 1920 by Mr. P. De Santis. In 1923-1924, De Santis recovered a total of 2.8 tonnes of hand-picked ore, with an average grade of 250 g/tonne

(Middleton 1975). The property has a shaft and several hundred metres of underground workings. There has been no production since 1924.

MAGMATIC Ni-Cu OCCURRENCES

Several Ni-Cu (and very low PGE) sulphide occurrences are located within the more magnesian cumulates of the KGC. The three most significant are termed the northwest Carscallen occurrences, Bean Lake- Pirsson Lake occurrences, West Whitesides Township occurrences, and correspond to numbers 10, 11, and 12, respectively in Figure 4. Samples from these localities have been analyzed for Ni, Cu, Co, and in some cases S, Pt, Pd, Au, and the remaining platinum group elements: Os, Ir, Ru, and Rh. These data are reported in Table 9.

West Whitesides Township Occurrences

In central west Whitesides Township, 500 m west of the Kamiskotia River, eight trenches and numerous drill holes have outlined sulphidic pods with low grade Cu-Ni mineralization. The mineral occurrences are generally aligned parallel to the cumulate banding in the host rocks, with trenches oriented east-west, parallel to a weakly developed, non-penetrative structural fabric. A standard magmatic mineral assemblage of pyrrhotite, chalcopyrite pentlandite and magnetite occur interstitial to silicate minerals in a net-texture, and also injected into the host rock along jointing or fault planes locally. Graham (1931) reported 0.6 wt.% Ni, in a pyrrhotite-rich grab sample from this area. Samples 86-321 to 86-329 (Table 9) are considered representative of the mineralization in the trenches at surface; they do not contain any appreciable base or precious metal values.

Bean Lake- Pirsson Lake Occurrences

Two low grade magmatic sulphide occurrences are present in central Whitesides Township: one 0.5 km west of Bean Lake, and another 0.5 km west of Pirsson Lake. The Bean Lake occurrence was visited for this study. Extensive stripping in 1955 exposed iron formation, in contact with the base of the KGC. Sulphide mineralization occurs in both the iron formation, and in the gabbro, 50 m to the north. Previous sampling of outcrop and drill core in this area determined very low Cu, Ni and Au abundances in net-textured and semi-massive sulphide (Wolfe 1970), although silver assays up to 3.3 oz. (100 g)/tonne have been reported (Leahy 1968). Low base and precious metal values are found in one surface grab sample in this study (e.g., sample 86-27: Table 9).

Northwest Carscallen Township Occurrences

Magmatic sulphide mineralization is found in northwest Carscallen Township and northeast Whitesides Township, located immediately south of the prominent (stands in high relief) peridotite outcrop in the area. Here net-textured chalcopyrite and pyrrhotite are present within altered olivine gabbro. Twelve diamond drill holes encountered disseminated sulphides with low Cu and Ni values (Wolfe 1970). Samples 84-14a and 84-175 are typical of the net-textured sulphide material; they have very low Ni, Cu and PGE contents (Table 9).

POTENTIAL FOR MINERALIZATION

In this section, the potential for the discovery of volcanogenic Cu-Zn deposits, mesothermal Au deposits, REE deposits and magmatic Ni-Cu- PGE deposits is discussed.

Additionally, lithogeochemical exploration strategies for each of these deposit types is outlined briefly.

Volcanogenic Cu-Zn Deposits

The potential for the discovery of volcanogenic Cu-Zn deposits in the Kamiskotia area is considered high. The four deposits of the Kamiskotia Cu-Zn camp are located within the KVC, within a stratigraphic sequence less than two km thick, of mixed mafic and felsic metavolcanic rocks. Incompatible element-enriched basalts, basaltic andesites and andesites are present in close proximity the massive sulphide lenses, forming footwall and hanging wall flows and sills; relatively fresh andesite dikes and sills cut the lenses at the Kam-Kotia mine (Pyke and Middleton 1970). These incompatible-element enriched mafic rocks have been shown to be the product of high degrees of fractional crystallization in a tholeiitic system, with the residual cumulus phases represented in the KGC to the south. High silica rhyolites in the KVC may be the product of more advanced fractionation. The latent heat of fractionation in high level magma chambers has been modeled quantitatively by Cathles (1983), who demonstrated that this mechanism is adequate to drive base metal-precipitating, hydrothermal convection through overlying, volcanic strata.

Exploration for volcanogenic base metal deposits should focus on the KVC, particularly where the stratigraphy is composed of mixed, coeval, incompatible element-enriched mafic and felsic metavolcanic rocks. These areas should be characterized by evidence for syn-volcanic faulting, and mineralogical (chlorite, sericite, pyrite addition) and chemical (e.g., proximal sodium depletion, magnesium and iron enrichment) typical of low- CO_2 (e.g., carbonate and chloritoid not significant in Kamiskotia camp), volcanogenic massive sulphide-precipitating hydrothermal systems.

Mesothermal Gold Deposits

The potential for mesothermal gold deposits is considered moderate. Efforts to locate mesothermal gold mineralization in the Kamiskotia area should focus on late-tectonic structures: along the DPFZ and south of granitoid D in Bristol and Denton townships; along the splay of the DPFZ through the Lower volcanic suite in southern Carscallen and Whitesides townships; or along the Kamiskotia highway fault, in north Godfrey and southern Jamieson townships. The vast majority of gold mineralization in the southern Abitibi subprovince is found along late-tectonic deformation zones (e.g., Colvine et al. 1988). North-trending vein systems, such as at the De Santis and Lally properties, are considered less prospective. They are interpreted to have developed during regional north-south compression, prior to the late tectonic events in the Kamiskotia area.

Gold mineralization on the Holmer and Croxall properties, within the DPFZ, is proximal to the incompatible element-enriched, Bristol Township lamprophyre suite. In this respect, the mineralization is similar to gold mineralization associated with light REE- and P_2O_5 -enriched rocks in Taylor Township, and elsewhere along the DPFZ and Kirkland - Larder fault zone to the south (King and Kerrich 1987). The gold-bearing quartz - tourmaline veins cut the lamprophyre suite on the Croxall property, and are believed to be genetically distinct from the lamprophyre suite. Both gold mineralization and light REE-enrichment are believed to be produced during similar, regional tectonic processes, such as late transpression and crustal underplating by mafic magmas, that affected the crust and upper mantle in this region (discussed further below).

The potential for economic REE concentrations in the Bristol Township lamprophyre suite is considered low to moderate, considering: 1) its high REE concentrations, of up to 0.43 wt.% RE_2O_3 ; 2) that the REE are principally in apatite, which is easily dissolved in HCl; 3) and the size of the lamprophyre suite, interpreted from ground and air magnetic

surveys to extend for several kilometres. However, it is difficult to envisage competing with current REE producers, such as deposits with bastnaesite ($\text{REE}(\text{CO}_3)\text{F}$) as the primary REE mineral. The carbonatite at Mountain Pass, California is an example of a primary bastnaesite producer. The Mountain Pass deposit has approximately 31 million tonnes of ore, and contains, on average, approximately 12% bastnaesite (containing 74 wt.% RE_2O_3), or an average grade of 8.9 wt.% RE_2O_3 (O'Driscoll 1988).

Magmatic Ni-Cu-PGE Deposits

The potential for magmatic Ni-Cu-PGE deposits in the KGC is considered low. All sulphide occurrences in the Lower Zone contain very low Cu, Ni and PGE contents. Primary modal layering, an indicator of magma chamber processes that are favorable to reef-type PGE mineralization (e.g., establishment of stable convection within the magma chamber), is only poorly developed in the KGC. There are no known, stratiform horizons of pegmatitic-textured gabbros in the KGC, such as those found in the Bushveld and Stillwater Complexes. A reconnaissance program (52 samples), designed to test the potential for PGE mineralization in pegmatitic gabbros of any type did not detect any significant enrichment in Pt, Pd or Au (Barrie, unpublished data).

The low Ni, Cu and PGE contents are interpreted to be the result of limited and passive contamination of KGC mafic liquids by footwall rocks. This is consistent with the fractionation-dominated, geochemical trends in the KGC - KVC system (see Geochemistry section). Assimilation of siliceous, iron-rich or sulphur-rich rocks by mafic - ultramafic magmas causes sulphide immiscibility, and Ni, Cu and the PGE partition strongly into the resulting immiscible sulphide liquid. Under turbulent flow conditions in the magma chamber, the sulphide liquid can scavenge greater quantities in the siderophile elements and become increasingly enriched in them (e.g., Naldrett 1981).

DISCUSSION

Gabbroic Complex - Granitoid Relationships in the Southern Superior Province

One of the notable features in the Kamiskotia area is the close temporal relationship between voluminous tholeiitic magmatism followed by calc-alkaline magmatism. This phenomenon has been documented by geological and/or geochronological studies across the southern Superior Province: at the Bird River Sill, Manitoba (Trueman and Bannatyne 1982), the Bad Vermilion Complex, Ontario (Ashwal et al. 1985), Mulcahy Lake and related gabbroic intrusions, Ontario (Morrison et al. 1985), the Sturgeon Lake gabbro, Ontario (Davis and Trowell 1982); Lac des Iles and related intrusions, Ontario (Sutcliffe et al. 1989), the Montcalm Gabbroic Complex, Ontario (Barrie and Naldrett 1989), the Bell River Complex, Quebec (Sharpe 1968), and the Doré Lake Anorthosite, Quebec (Dimroth et al. 1986). It would appear that these relationships are widespread, and the process of nearly coeval, voluminous mafic and felsic magma generation is integral to the development of granitoid-greenstone terranes in the Superior Province.

For the Kakagi Lake - Savant Lake volcanic belt (KSVB) and the Wabigoon diapiric axis in northwestern Ontario, Morrison et al. (1985) proposed that partial melting of a slightly depleted mantle produced tholeiitic magmas, parental to supracrustal gabbroic intrusions, and concurrent partial melting of overlying, young oceanic crust produced calc-alkaline magmas, parental to major granitoid complexes. Both melting events were in response to the same thermal event. They considered a migrating mantle plume as the most plausible heat source. In contrast, Davis et al. (1988) suggested that tholeiitic metavolcanic rocks of the lower KSVB evolved, for the most part, in an ensimatic, oceanic environment, in the absence of older sialic material. They postulated that late calc-alkaline magmatism is the product of an arc-like tectonic setting, related to partial melting above a southward-subducting oceanic slab.

In the Kamiskotia area, trace element studies indicate that KGC parental magmas are compatible with a 34% bulk partial melt of a primitive mantle, whereas granitoid rocks may be derived from partial melting of either a hydrated basaltic eclogite or reasonable lower crustal compositions (Barrie 1990). Older sialic crust was at least partly involved in the calc-alkaline magmatism that produced the regional granitoid complexes. For example, granitoid B has one zircon population that shows inheritance from a crustal component at least 2926 Ma old. Further evidence is provided by Sm-Nd isotopic systematics for Kamiskotia area rocks. Low, enriched $\epsilon_{Nd}(t)$ values of +1.0 to -0.4 have been determined for two <2700 Ma granitoids and a 2702 Ma ferroan gabbro sample in the region, implying the presence of an enriched, older crustal reservoir (Barrie 1990). Elsewhere in the southern Abitibi Subprovince, isotopic studies have provided indirect evidence for a sialic basement. For example, a model 3.0 Ga crustal reservoir has been proposed for late granitoids in the eastern Abitibi Subprovince, based on Pb-Pb isotope systematics (Gariépy and Allegre 1985).

Chemical Comparison of the KGC with Other Synvolcanic Intrusions of the Southern Superior Province

The KGC is one of several late Archean, large mafic intrusions that underlie massive sulphide-bearing, bimodal volcanic centres in the southern Superior Province, including the Bad Vermilion, Bell River and Doré Lake complexes. At the Bad Vermilion Complex, the majority of gabbros have flat REE patterns at 2 to 10x chondrites and positive Eu anomalies, nearly identical to the MZ and lower UZ cumulates (Ashwal et al. 1985). The complex intrudes into and is overlain by basalts similar to the Kamiskotia basalts that exhibit similar fractionation trends, and by felsic metavolcanic rocks similar to the distinctive, trace element-enriched Kamiskotia rhyolites (Hart 1984). The complex and related metavolcanic rocks have $\epsilon_{Nd}(t) = +2.0 \pm 1.4$ (Ashwal et al. 1985), similar to but slightly less depleted than the KGC Nd isotopic signature. Rocks of this region of the Superior Province are generally

slightly isotopically enriched with respect to the Abitibi-Wawa Subprovinces (Shirey and Carlson 1988; Barrie and Shirey 1989). Detailed trace element and isotope geochemistry is lacking for the Bell River and Doré Lake complexes. However by virtue of similar geologic settings and similar geochemistry of overlying bimodal metavolcanic rocks (MacGeehan and MacLean 1980; Ludden et al. 1984), it is probable that like the Bad Vermilion complex have petrogenetic histories similar to the KGC.

Chemical and Tectonic Comparison of the KGC with the Skaergaard Intrusion, and the KVC with Eastern Iceland Volcanic Rocks and Volcanic Rocks of the Galapagos Spreading Center

Perhaps the closest modern analogs for the KGC and related basalts are found in rift-related, tholeiitic magmatism represented by the Skaergaard Intrusion, East Greenland and the bimodal volcanic fields of Eastern Iceland. The petrology and geochemistry of the KGC's MZ and UZ overlaps with the Lower Zone and Middle Zone at Skaergaard, which is renowned for its classic tholeiitic Fe-enrichment trend (Wager and Brown 1968). Estimates of the liquid compositions based on mass balance calculations (Wager and Brown 1968; Hunter and Sparks 1987) and nearby dike compositions (Brooks and Nielson 1978) indicate that the Skaergaard magma probably remained below 50% SiO₂ until magnetite precipitation, at the top of the Lower Zone. Silica behaves in a similar manner in the Kamiskotia system. Hunter and Sparks (1987) believed that as the Skaergaard magma continued to fractionate, it produced Fe- (and Ti- and P₂O₅-) rich andesitic liquids, parental to the apatite-bearing Upper Zone, and finally rhyolitic liquids, parental to the Sandwich Zone horizon. The major element trends of their model liquids are similar to the Kamiskotia basalt-evolved basalt-andesite-rhyolite trends (Hart 1984; Barrie 1990).

Hunter and Sparks (1987) noted the similarities in major element chemistry between their Skaergaard model liquids and several of the bimodal volcanic fields of Eastern Iceland.

They proposed that Iceland volcanic suites were the products of similar, fractionation-dominated systems. The basalt-ferrobasalt-icelandite-rhyolite suite of the Askja 1875 eruption, believed to reflect a fractionation-dominated system (MacDonald et al. 1987), has major and trace element characteristics similar to the Kamiskotia volcanic suite (Hart 1984). Sigurdsson and Sparks (1981) and MacDonald et al. (1987) envisioned a dynamic, compositionally stratified magma chamber beneath the Askja volcanic field. They postulated that prior to the eruption, a ferrobasaltic liquid, which represents 60 to 70% fractionation of parental, magnesian basaltic liquid, was present at the floor of the chamber, progressively overlain by andesitic to rhyolitic liquids. If such a chamber ceased to erupt and crystallized in situ, this could represent the KGC's UZ and granophyric cap of intermediate to felsic composition.

The compositions of basalts, Fe-Ti basalts and rhyodacites at the 85°W propagating ridge of the Galapagos Spreading Center, are also similar to the Kamiskotia suite. The rocks at 85°W are too far from the Galapagos hot spot to reflect mantle plume characteristics, so their petrology and geochemistry are believed to reflect a variation of mid-ocean ridge magma chamber processes (Schilling et al. 1976; Verma et al. 1983). Mineral compositions closely resemble the compositions in the KGC, and glass compositions follow fractionation trends similar to the Kamiskotia basalts (Byerly 1980; Juster et al. 1989). These fractionation trends have been duplicated in experiments on Fe-Ti basalts at one-atmosphere, but under more oxidizing conditions (Ni-NiO buffer) than would be expected by a closed system, implying an interaction with oxidizing surroundings (Juster et al. 1989). At Kamiskotia, fractionation occurred at shallow levels, and may have been influenced by extensive, and potentially oxidizing hydrothermal fluids that formed the contemporaneous volcanogenic Cu-Zn deposits (Barrie and Davis 1990).

Cogenesis of the Kamiskotia Gabbroic Complex and Kamiskotia Volcanics, and Significance with Respect to the Formation of Volcanogenic Massive Sulphide Deposits

That the KGC and the Kamiskotia rhyolite are coeval within geochronological error strengthens geochemical arguments for their cogenetic relationship. The Kamiskotia felsic metavolcanic rocks and the underlying KGC granophyric cap have nearly identical rare earth element abundances, which implies that the rhyolites are eruptive equivalents of the granophyre (Campbell et al. 1981). Both have distinctive high, flat chondrite-normalized patterns with strong negative Eu anomalies, consistent with low pressure fractionation of mafic phases and plagioclase. The rhyolites have been termed 'tholeiitic' rhyolites by Campbell et al. (1982), or 'Type IIIb' rhyolites by Leshner et al. (1986). Kamiskotia basalts were successfully modeled as fractionation products of a primitive MORB-like basalt, with the residual phases represented in the KGC (Hart 1984). Sm-Nd isotope systematics are also compatible with the cogenesis of the KGC and KVC: a 7-point, KGC-KVC whole rock and mineral isochron has an age of 2730 ± 30 Ma, in agreement with the U-Pb age, and all samples have identical $\epsilon_{\text{Nd}}(t)$ values within error (Barrie 1990).

The presence of metavolcanic-hosted massive sulphide deposits within a 2 km thick stratigraphic level in the KVC is consistent with the coeval nature of the KGC and KVC. During crystallization and cooling, the KGC was capable of providing the heat necessary to drive hydrothermal convection cells that would leach metals from a volcanic pile and precipitate them on the paleo- sea floor, given geologically reasonable permeability and temperature constraints (Cathles et al. 1983). Hypabyssal gabbroic intrusions that underlie massive sulphide-bearing 'tholeiitic' rhyolites are present elsewhere in the southern Superior Province: the Bad Vermilion Complex overlain by the Gagne Lake deposits, Ontario (Poulsen 1984), the Bell River Complex and Matagami Lake district (MacGeehan and MacLean 1980), numerous gabbro sills in the Noranda district, and the Doré Lake anorthosite overlain by the

Patino-Lamoine mine and other deposits (Guha et al. 1988). It seems likely that in these areas, as in the Kamiskotia area, the gabbroic intrusions are synvolcanic, and they provided the heat necessary for massive sulphide-precipitating hydrothermal systems.

Kamiskotia - Kidd Creek Relationships

Pyke (1978, 1982) tentatively correlated the Kidd Creek rhyolite with Kamiskotia rhyolites. He noted their similar geochemistry, the presence of massive sulphide mineralization and their general stratigraphic position below the metasedimentary rocks of the Tisdale Group in the Timmins area, and assigned them to the Upper Deloro Group. In the OGS-MERQ (1983) lithostratigraphic map of the Abitibi Subprovince, both metavolcanic piles are placed at the top of the Cycle III Volcanics, one of four cyclic metavolcanic packages tentatively correlated across the Abitibi Subprovince.

The U-Pb ages reported in Barrie and Davis (1990) here indicate that modification is necessary for the correlations of metavolcanic rocks for the western Abitibi Subprovince. It is apparent that the Kidd Creek and Prosser rhyolites are at least 10 Ma older than the KGC, the Kamiskotia rhyolite, and the Reid rhyolite 11 km to the west-northwest of the Kidd Creek mine. The Kidd mega-agglomeritic flow-banded rhyolite sample and the Kamiskotia flow-banded rhyolite sample both represent proximal volcanic facies and are proximal to mine sites. In contrast, the Prosser and Reid samples are crystal-ash tuffs intercalated with graphitic argillites, typical of distal, subaqueous volcanic facies. They may represent crystal-ash deposits distal to the Kidd and Kamiskotia volcanic centers, respectively. Samples of the more distal deposits have lower zirconium contents (Appendix I) and yielded far fewer zircons. This may reflect crystal sorting during ash fall eruptions, which would be consistent with their distal characteristics, or it may indicate that these eruptions tapped relatively zircon-poor siliceous magmas.

Time - Stratigraphic Correlation of Volcanic Rocks Across the Southern Abitibi Subprovince

Corfu et al. (1989) reported an age for a proximal(?), banded rhyolite from the Stroughton-Roquemaure Group of 2714 ± 2 Ma, and a revised age for a dacitic unit in the Hunter Mine Group to the east of the Ontario - Quebec border of 2713 ± 2 Ma. These samples were taken generally along strike from the Kidd Creek mine, to the east-southeast 75 km and 110 km, respectively. Thus, there is a consistent decrease in ages over this 110 km trend, from 2717 ± 2 Ma at the minesite to 2713 ± 2 Ma. This may imply that volcanism migrated eastward over a 4 Ma period. Furthermore, the geochemistry of the tholeiitic mafic metavolcanic rocks is similar from Kidd Creek to the Hunter Mine Group area (Pyke 1982; Jensen and Pyke 1982), and the distinctive 'tholeiitic' rhyolite geochemistry of the Kidd Creek rhyolite is found farther to the east along strike, in southeast Harker Township (Goodwin, unpublished data). By comparison with the geology and geochemical modeling at Kamiskotia, this would imply that the Kidd Creek - Hunter Mine Group trend may be associated with hypabyssal mafic intrusions that haven't been noted in outcrop.

Four U-Pb ages for metavolcanic lithologies and one hypabyssal dunite sill to the east of the Kamiskotia area are coeval to slightly younger than the KGC and Kamiskotia rhyolite, from 2707 ± 1.5 Ma to 2698 ± 4 Ma (Corfu et al. 1989). These include rocks previously included in the Upper Deloro, Tisdale, Skead, Larder Lake and Blake River Groups. The upper Blake River Group metavolcanic rocks are similar to the KVC rocks in several respects: they were deposited during active tectonism, with synvolcanic faulting (Gibson et al. 1989); they are bimodal; the rhyolites have similar, flat REE profiles with negative Eu anomalies (Leshner et al. 1986); they are underlain by a sub-volcanic sill, the coeval (both at 2701 ± 2 Ma: Corfu et al. 1989), and cogenetic (Paradis et al. 1988) Flavrian pluton, that has magma mixing textures (Goldie 1978); and they are host to volcanogenic Cu-Zn deposits of the Noranda camp. Considering these similarities, the KVC and upper Blake River Group may have erupted in a similar tectonic settings.

Late Transpression Across the Southern Superior Province

Late north-south compression/transpression has been documented by structural and geochronologic studies for areas across the southern and central Superior Province, at major subprovince boundaries (Stott 1985; Corfu and Stott 1986; Stott et al. 1987; Davis et al. 1989) and within greenstone belts (Hubert et al. 1984; Dimroth et al. 1986; Hudleston et al. 1988). The majority of these studies have indicated that transcurrent motion on regional east-west or west-northwest - east-southeast shear zones is in a dextral sense, whereas for west-northwest - east-southeast shear zones, a sinistral motion is indicated. These observations are compatible with a sub-horizontal stress regime oriented in a west-northwest - east-southeast sense (Stott et al. 1987). Dextral transcurrent motions along the Quetico - Wawa and Quetico - Wabigoon Subprovince boundaries are tightly bracketed by U-Pb analyses of deformed and undeformed rocks at 2689 to 2684 Ma and 2696 to 2686 Ma, respectively (Corfu and Stott 1986; Davis et al. 1989). The timing and sense of displacement along these subprovince boundaries are virtually identical to that in the Kamiskotia area. The age of the Bristol Township lamprophyre suite is 2687 ± 3 Ma, and is interpreted to have been emplaced after the formation of the DPFZ, but was subsequently deformed within it. Late-stage, predominantly dextral transpression is recorded in the splay of the DPFZ, and interpreted for the Kamiskotia Highway fault in the Kamiskotia area. This implies a synchronous, crustal shortening event across 1200 km of the southern Superior Province at approximately 2690 to 2685 Ma.

Timing of Late Archean Magmatism in the Southern Superior Subprovince

Figure 35 presents a compilation of high precision U-Pb ages for "Kenoran-aged" magmatism in the southern Superior Province, designed to highlight the timing of different

rock types. In a sense, this compilation of U-Pb ages supersedes the compilation of Rb-Sr whole rock isochron ages and K-Ar mineral ages used in the original definition of the Kenoran Orogeny, which has a mean age of 2480 Ma (Stockwell 1970). Rb-Sr and K-Ar data most probably reflect cooling histories and/or later thermal events that have effected the southern Superior Province (e.g., Beakhouse et al. 1988; York and Hyodo 1988; Barrie 1990).

All of the divisions are straightforward, with the exception of the division between LILE-/LREE-enriched rocks and calc-alkalic intrusive rocks. The LILE-/LREE-enriched rocks are represented by alkalic metavolcanic rocks, shoshonitic ("calc-alkalic") lamprophyres, and intermediate to felsic intrusions that are known to have a component of LILE-/LREE-enriched and/or alkalic affinity. Several of the intermediate to felsic intrusions are predominantly silica-saturated and have $\text{Na}_2\text{O} + \text{K}_2\text{O}$ contents that do not clearly distinguish them as alkalic; however, higher LILE and LREE contents coupled with high Cr, Ni and MgO contents distinguish them from trondhjemite-tonalite or granodiorite-granite series intrusions (e.g., Ottertail Lake pluton, Quetico Subprovince: Shirey and Hanson 1986).

From this compilation, calc-alkalic volcanism appears to have ended abruptly at 2695 to 2700 Ma: the youngest calc-alkalic metavolcanic rocks are found in the southern Abitibi Subprovince (Krist Fragmental tuff: Corfu et al. 1989; upper Blake River Group: Mortensen 1987). From the few U-Pb ages available, tholeiitic and ultramafic magmatism also appear to end at this time. Barrie and Davis (1990) speculated that this may be due to the cessation of the subduction of an oceanic plate at 2695 to 2700 Ma.

The timing of calc-alkalic intrusive activity approximates a gaussian distribution with a mean ($n=58$) at 2700 Ma. Gneissic rocks represent amphibolite to granulite grade calc-alkalic intrusions, from the Kapuskasing Structural Zone and related rocks to the southwest. Generally, their ages are younger than 2700 Ma. Corfu (1987) recognized that there may be a progressive younging of the gneisses with depth, and discussed the possibility of magmatic underplating by mantle-derived material (e.g., Bohlen 1987).

Evidence for Late Thermal Events in the Kamiskotia Area

Thermal events after crystallization have affected rocks in the Kamiskotia area and reset both the Nd and Sr mineral systematics locally. As mentioned above, a Rb-Sr isochron for the KGC ferroan gabbro whole rock and mineral separates agrees with U-Pb ages for the Hearst-Matachewan dike swarm (Heaman 1989). This sample was taken 7 m away from a 6 m thick, vertical Matachewan dike in an area where the dikes are particularly densely spaced at 200 m intervals on average. The Rb-Sr isochron age is interpreted as the age of thermal/hydrothermal metamorphism due to the emplacement of the neighboring dike. Delaney (1987) has reviewed heat transfer processes by conduction adjacent to cooling vertical mafic dikes. For cases where the thermal properties of the dike and host rock are identical, such as for Kamiskotia gabbroic rock and diabase dikes, the ambient temperature at 1.2 dike widths would rise approximately 200°C. This would be in addition to the temperature due to the geothermal gradient, and to heating from other Matachewan dikes and sills in the vicinity. The presence of intergranular fluid would allow for slightly more rapid heat transfer in crystalline igneous rocks (Hyodo, in prep.) and would aid Rb and Sr diffusion. Resetting of the K-Ar system in hydrous phases up to 15 m from a 10 m thick dike has been clearly documented (Hanes et al. 1988) and modeled by convection of fluids adjacent to mafic dikes (York and Hyodo 1988, and Hyodo in progress). The Sm-Nd system, generally considered more resistant to resetting than the Rb-Sr and K-Ar systems has not been disturbed for this sample. An internal, whole rock - plagioclase - clinopyroxene Sm-Nd regression line for this sample has an age of 2700 ± 30 Ma (MSWD = 8.0), in agreement with the KGC-KVC Sm-Nd isochron, and the U-Pb ages for the KGC and KVC rhyolite.

The emplacement of the Hearst - Matachewan dike swarm represents a major magmatic event after the stabilization of the Superior Province. It extends over greater than 250,000 km² of the Superior Province, the second largest dike swarm of the Canadian Shield

and one of the largest on earth (Fahrig 1987). The studies of Phinney and Morrison (1988) provide evidence for extensive Matachewan-related tholeiitic magma chambers at upper to mid-crustal levels, which may have been capable of resetting the Rb-Sr system regionally. Their experimental petrology on tholeiitic matrices of megacrystic basalts similar to Matachewan dike matrix material indicated that Matachewan plagioclase compositions can be reproduced only at pressures corresponding to depths of about 10 to 12 km. The thermal effects of Hearst-Matachewan magmatism may have been responsible for partially or completely resetting Rb-Sr and K-Ar ages for Archean rocks in a cryptic fashion, with no detectable change in mineral assemblages. For example, Rb-Sr isochron ages for many granitoids across the southern Superior Province are 2400 Ma to 2460 Ma (e.g., Frith and Doig 1975; Birk and McNutt 1981; Clark and Cheung 1980); and Rb-Sr systematics of three alteration zones along the Destor Porcupine and Larder Lake fault zones give regression ages of 2390, 2410 and 2440 Ma (Kerrick et al. 1987).

Younger regression ages for two granitoid rocks in the Nd and Sr systems are older than the emplacement age for the Hearst dike swarm. Particularly noteworthy are the identical ages of 2530 Ma in the Nd and Sr systems for a sample from granitoid C, coeval with U-Pb titanite ages at the base of the Kapuskasing Structural Zone (Krogh et al. 1988), and more than 160 Ma younger than U-Pb zircon and titanite ages from the same sample.

The young Sm-Nd and Rb-Sr ages may reflect thermal events related to the late addition of mantle-derived magmas into the lower crust of the southern Superior Province, after most of the magmatic activity in the late Archean. In the southern Abitibi Subprovince, radiogenic isotopic studies have investigated mineralization and alteration associated with mesothermal Au systems (e.g., Wong et al. 1989; Bell et al. 1989) and volcanogenic sulphide deposits (e.g., Maas et al. 1986; Schandl and Davis 1989). These studies have documented posttectonic hydrothermal fluid migration related to late thermal events from 2650 to 2400 Ma (see Bell et al. 1989 for a recent review). Crustal underplating and the intrusion into lower and mid-crustal levels by mantle-derived magmas is one

mechanism that can adequately explain the late addition of heat to this region (Fyon et al. 1989; Barrie 1990). Crustal underplating has been called on to explain anti-clockwise P-T paths documented in numerous granulite terranes, and is believed to be characteristic of continental arc environments (Bohlen 1987). For thermal modeling of the nearby Kapuskasing Structural Zone uplift, it is necessary to include regional uplift and continued elevated heat input from the mantle between 2680 Ma and 2500 Ma (Percival et al. 1988). This is consistent with the addition of mantle-derived magmas into the lower crust at 2530 Ma. Therefore, a continental arc environment would account not only for the igneous characteristics of the Abitibi and Wawa Subprovinces, but the Kapuskasing Structural Zone metamorphic and cooling histories as well (Percival et al. 1988).

Comparison to Modern Tectonic Settings: Synthesis

Perhaps the closest Phanerozoic analog to the tectonic setting of the KGC, considering its close temporal relationship with regional calc-alkalic plutonism, is found in the continental arc setting of the central Peruvian Andes. The vast Coastal Batholith of Peru exhibits classic exposures of Mesozoic and Cenozoic intrusive rocks that have been exhumed by continued uplift and erosion. The Coastal Batholith intruded into a tectonically thinned, extensional marginal basin, (Atherton et al. 1983), underlain by a relatively dense basement, interpreted to represent mantle or lower crustal material (Couch et al. 1981). A series of large (up to 5 km by 40 km), elongate gabbro to diorite intrusions comprise approximately 16% of the Coastal Batholith (McCourt 1981, Regan 1985). Geologic relationships and geochronology have shown that the gabbros predate the voluminous calc-alkaline magmatism. In the Arequipa (southern) segment, the Early Gabbros, at 101-106 Ma, were followed by a series of granitic intrusions at 37-101 Ma (Moore and Agar 1985; Mukasa 1986). There is abundant textural evidence in the gabbros for syn-crystallization deformation, volatile activity, and net veining by granitic magmas to form hybrid rocks during consolidation (Regan 1985).

All of these textures are found within the KGC (Barrie 1990). Atherton et al. (1981) have proposed that the parent magmas for the gabbros were olivine tholeiites, a product of partial melting in a mantle wedge above a subducting slab, leaving behind a garnet-free residuum. The partial melts were modified by olivine and clinopyroxene fractionation prior to emplacement at mid- to upper crustal levels. Generally, calc-alkaline to mildly alkaline granitoid rocks like those of the coastal batholith, may have parental magmas that were derived by partial melting of amphibolitic or eclogitic, subducted oceanic crust, and hydrated peridotitic upper mantle, respectively (Arth and Hanson 1975; Shirey and Hanson 1986). Pb and Sr isotope studies indicate minimal interaction with older crustal material for Coastal Batholith rocks (Barreiro and Clark 1984; Pitcher et al. 1985).

Other extensional tectonic settings are suggested through comparisons of the geochemistry, physical volcanology and mineralization of the KVC. The geochemistry and distinctive mixed magma textures in felsic pyroclastic rocks compare favourably to Askja, Iceland volcanic rocks, representing a hot spot, mid-ocean rift setting; (Sigurdsson and Sparks 1981). The presence of Cu-Zn volcanogenic massive sulphide deposits at Kamiskotia is commonly regarded to indicate subaqueous rifted settings (Cathles et al. 1983). This analogy would apply for the older, Kidd Creek - Hunter Mine Group metavolcanic trend also. Kamiskotia and Kidd Creek rhyolites are chemically and texturally similar to a number of high silica rhyolite suites from bimodal fields in the extensional, back-arc setting the of the Basin and Range (e.g., Crecraft et al. 1981, Bacon et al. 1981).

The trace element and isotopic signatures of the KGC and KVC are consistent with geologic and U-Pb geochronological evidence for their formation in an continental arc-like regime. The bimodal KVC, with its massive sulphide mineralization, is typical of settings generally considered to represent a rifted back-arc or arc tectonic regime (e.g., Cathles et al. 1983). The presence of older crustal rocks nearby (described above) would suggest that rifting must have been within or proximal to an ensialic crust, possibly a continental margin. One notable feature about the geochemistry of the KGC chill margins and basalt is their

MORB-like trace element signatures. The trace element contents of KGC chill margin rocks are consistent with partial melting of a chemically primitive mantle peridotite at shallow depths, and show little or no contamination by crustal components. Similarly, the Nd isotopic signature for the entire Kamiskotia suite is consistent with a direct derivation from the Abitibi depleted (in an isotopic sense) mantle with little or no interaction from enriched crustal material. In modern continental arc settings such as in the central Peruvian Andes or the Peninsular Ranges Batholith of Baja California, there is isotopic evidence that mantle-derived tholeiitic magmas rose through the lithosphere with little or no interaction from crustal material. Large gabbroic intrusions which represent the earliest intrusive rocks of the Coastal Batholith were emplaced in a tectonically thinned, extensional marginal basin (Atherton et al. 1983). Pb and Sr isotope studies indicate minimal interaction with older crustal material for Coastal Batholith rocks (Barreiro and Clark 1984; Pitcher et al. 1985). Large gabbroic intrusions in the Peninsular Ranges are slightly older and more isotopically depleted than the majority of nearby granitoid plutons (Walawender 1976; Walawender and Smith 1980; DePaolo 1981b).

The granitoid rocks were emplaced during times of extensional tectonics associated with the KGC, and later during a more compressional regime. In this respect the granitoid suite is similar to the Cretaceous Peninsular Ranges batholith of Baja California, the product of arc magmatism formed during subduction of the Pacific plate under the North American plate. Gromet and Silver (1987) found that tonalites and low-K granodiorites exhibit a systematic increase in Ce/Yb from west to east, generally accompanied by an increase in I_{Sr} values. These trends cannot be explained by high level fractionation of major and trace silicate phases. They are consistent with partial melting of arc basalt under increasingly high pressures, leaving behind gabbroic or amphibolitic residue at low pressures to the west, and an eclogitic residue to the east (Gromet and Silver 1987). The Kamiskotia granitoids have Ce/Yb ratios most similar to the central and eastern granitoids of the Peninsular Ranges, consistent with their derivation from a basaltic source at high pressures. Enriched Nd

isotopic signatures and generally higher Ce/Yb ratios for the central granitoids, which represent a regional granitoid terrane that encompasses the greenstone belts of the southern Superior Province, are consistent with derivation from an older basaltic source at greater depths in this area. Further Nd isotopic studies of the regional granitoids are necessary to test whether there are aurally distinct domains of older, enriched crust in the vicinity of the southern Abitibi Subprovince and elsewhere in the southern Superior Province.

The late, predominantly transpressional regime represents a major change in tectonic style in the Kamiskotia area and across the southern Superior Province. Transpressional tectonics were at least in part responsible for the development of Superior Province-wide structural discontinuities, and were generally coincident with alkaline magmatism (Frarey and Krogh 1986; Corfu and Stott 1986; Corfu et al. (1989); Davis et al., (1989); Wyman and Kerrich 1988; Barrie and Davis 1990). The close spatial association between major fault structures and mantle-derived alkaline rocks in the southern Abitibi Subprovince implies that at least locally, these faults served as conduits for alkaline magmas that extended to mantle depths (Kerrich et al 1987; Wyman and Kerrich 1988).

Similar spatial and temporal relationships between mantle-derived alkalic rocks and incipient transpressive regimes occur in Phanerozoic and modern arc settings, associated with a transition between relative plate motions. A rapid transition from subduction and collision-related calc-alkaline magmatism to alkaline magmatism in a transpressive regime has been documented by a structural and geochemical study of dike swarms, accompanied by Rb-Sr geochronology, for a part of the Pan-African belt in Mali (Liegeois and Black 1987).

Quaternary alkalic volcanic centers in the Colima graben area, Mexico, represent the transition from compressional to transpressional tectonics at the southern end of the San Andreas fault system (Luhr and Carmichael 1981). Morrison (1980) determined that arc-related shoshonitic rocks are commonly associated with the termination of subduction or the transition between two subduction regimes of different orientation in island arc and continental arc settings. This infers that for the Kamiskotia area and the southern Superior

Province, late transpression may have been related to a sudden change in plate motions in an Andean-type continental arc setting (Fig 37).

REFERENCES

- Arth, J.G., and Hanson, G.N. 1975. Geochemistry and origin of the early Precambrian crust of northeastern Minnesota; *Geochimica et Cosmochimica Acta*, v. 39, p.325-362.
- Ashwal, L.D., Wooden, J.L., Phinney, W.C., and Morrison, D.A. 1985. Sm-Nd and Rb-Sr isotope systematics of an Archean anorthosite and related rocks from the Superior Province of the Canadian Shield; *Earth and Planetary Science Letters*, v. 74, p.338-346.
- Atherton, M.P., McCourt, W.J., Sanderson, L.M., and Taylor, W.P. 1981. The geochemical character of the segmented Peruvian Coastal Batholith and associated volcanics; *in* *Origin of Granite Batholiths, Geochemical Evidence*, Atherton, M.P., and Tarney, J., eds., Shiva Publ. Ltd., p.45-64.
- Atherton, M.P., Pitcher, W.S., and Warden, V. 1983. The Mesozoic marginal basin of central Peru; *Nature*, v. 305, p.303-306.
- Bacon, C.R., Macdonald, R., Smith, R.L., and Baedeker, P.A. 1981. Pleistocene high-silica rhyolites of the Coso Volcanic Field, Inyo County, California; *Journal of Geophysical Research*, v. 86, p.10223-10241.
- Bacon, C.R., Kurasawa, H., Delevaux, M., Kistler, R., and Doe, B. 1984. Lead and strontium isotopic evidence for crustal interaction and compositional zonation in the source regions of Pleistocene basaltic and rhyolitic magmas of the Coso volcanic field, California; *Contributions to Mineralogy and Petrology*, v. 85, p.366-375.
- Barlow, R.B. 1988. Calculated second vertical derivative colour image from digital archives of the Ontario Geological Survey, Timmins Area, Districts of Cochrane and Timiskaming; Ontario Geological Survey, Map 81139, Geophysical/Geochemical Series, scale 1:100 000.

- Barnes, S.-J. 1984. The origin of the fractionation of platinum group elements in Archean komatiites of the Abitibi greenstone belt, northern Ontario, Canada; unpublished Ph.D. thesis, University of Toronto, 230 p.
- Barreiro, B.A., and Clark, A.H. 1984. Lead isotopic evidence for evolutionary changes in magma - crust interaction, Central Andes, southern Peru; *Earth and Planetary Science Letters*, v. 69, p.30-42.
- Barrie, C.T. 1990. Petrogenesis and tectonic evolution of the Kamiskotia and Montcalm gabbroic complexes and adjacent granitoid greenstone belt terrane, western Abitibi Subprovince, Ontario, Canada; Unpublished Ph.D. thesis, University of Toronto, 317 p.
- Barrie, C.T. and Davis, D.W. 1990. Timing of magmatism and deformation in the Kamiskotia - Kidd Creek area, western Abitibi Subprovince, Ontario, Canada; *Precambrian Research*, v. 46, p.217-240.
- Barrie, C.T., and Naldrett, A.J. 1989. Geology and tectonic setting of the Montcal gabbroic complex and Ni-Cu deposit, Western Abitibi Subprovince, Ontario, Canada; *in* *Magmatic Sulphides: the Zimbabwe Volume*, Prendergast, M.D., and Jones, M.J., eds., Institution of Mining and Metallurgy, London, p.151-164.
- Barrie, C.T., and Shirey, S.B. 1989. Geochemistry and Nd-Sr isotope systematics of the Kamiskotia area, western Abitibi Subprovince, Canada: implications for mantle processes during formation of the southern Superior craton; *Workshop on the Archean Mantle*, Lunar and Planetary Institute, Houston, Texas, p.31-33.
- Barrie, C.T. and Shirey, S.B. In preparation. Nd and Sr isotope systematics for the Kamiskotia - Montcalm area: implications for the formation of Late Archean crust in the western Abitibi Subprovince, Canada.
- Basu, A.R., Goodwin, A.M. and Tatsumoto, M., 1984. Sm-Nd study of Archean alkalic rocks from the Superior Province of the Canadian Shield: *Earth and Planetary Science Letters*, v.70 p.40-46.

- Bateman, R. 1985. Aureole deformation by flattening around a diapir during in situ ballooning: the Cannibal Creek Granite; *Journal of Geology*, v. 93, p.293-310.
- Beakhouse, G.P., McNutt, R.H., and Krogh, T.E. 1988. Comparative Rb-Sr and U-Pb zircon geochronology of late- to post-tectonic plutons in the Winnipeg River belt, northwestern Ontario, Canada; *Chemical Geology*, v. 72, p.337-351.
- Bell, K., Anglin, C.D., and Franklin, J.M. 1989. Sm-Nd and Rb-Sr isotope systematics of scheelites: possible implications for the age and genesis of vein-hosted gold deposits; *Geology*, v. 17, p.500-504.
- Beswick, A. 1982. Some geochemical aspects of alteration, and genetic relations in komatiitic suites; *in* Komatiites, Arndt, N.T., and Nesbit, E.G., eds., Allen Unwin, London, p.283-308.
- Birk, D., and McNutt, R.H. 1981. Geochronology of Wabigoon belt granitoids, northwestern Ontario: Rb/Sr isochrons for seven late-tectonic plutons; *Canadian Journal of Earth Science*, v. 18, p.157-175.
- Bohlen, S.R. 1987. Pressure-temperature-time paths and a tectonic model for the evolution of granulites; *Journal of Geology*, v. 95, p.617-632.
- Bright, E.G., and Hunt, D.S. 1973. Cote Township, District of Cochrane; Ontario Department of Mines map P.840.
- Brooks, C.K., and Nielsen, T.F. 1978. Early stages in the differentiation of the Skaergaard magma as revealed by a closely related suite of dike rocks; *Lithos*, v. 11, p.1-14.
- Brugmann, G.E. 1985. Geochemistry of the noble metals and lithophile elements in komatiite flows from Alexo, Canada, and Gorgona, Colombia; unpublished Ph.D. thesis, Johannes Gutenberg-Universitat, Mainz, F.R.G., 261 p.
- Byerly, G. 1980. The nature of differentiation trends in some volcanic rocks from the Galapagos Spreading Center; *Journal of Geophysical Research*, v. 85, pg. 3797-3810.
- Campbell, I.H., Coad, P., Franklin, J.M., Gorton, M.P., Scott, S.D., Sowa, J., and Thurston, P.C. 1982. Rare earth elements in volcanic rocks associated with Cu-Zn massive

- sulfide mineralization: preliminary report; Canadian Journal of Earth Sciences, v. 19, p.619-623.
- Campbell, I.H., Franklin, J.M., Gorton, M.P., Hart, T.R., and Scott, S.D. 1981. The role of subvolcanic sills in the generation of massive sulfide deposits; Economic Geology, v. 76, p.2248-2253.
- Campbell, I.H., Leshar, C.M., Coad, P., Franklin, J.M., Gorton, M.P., and Thurston, P.C. 1984. Rare earth element mobility in alteration pipes below massive sulfide deposits; Chemical Geology, v. 45, p.181-202.
- Capdevila, R., Goodwin, A.M., Ujike, O., and Gorton, M.P. 1983. Trace-element geochemistry of Archean volcanic rocks and crustal growth in southwestern Abitibi belt, Canada; Geology, v. 10, p.418-422.
- Cathles, L.M., Guber, A.L., Lenagh, T.C. and Dudas, F.O. 1983. Kuroko-type massive sulphide deposits of Japan: Products of an aborted Island-arc rift; *in* Economic Geology Monograph 5, The Kuroko and related Volcanogenic Massive Sulphide Deposits; Economic Geology Publishing Company, p.96-114.
- Cattell, A., Krogh, T.E., and Arndt, N.T. 1984. Conflicting Sm-Nd whole rock and U-Pb zircon ages for Archean lavas from Newton Township, Abitibi belt, Ontario; Earth and Planetary Science Letters, v. 70, p.280-290.
- Christiansen, E.H., Sheridan, M.F., and Burt, D.M. 1986. The geology and geochemistry of Cenozoic topaz rhyolites from the western United States, Special Paper 25; Geological Society of America, 82 p.
- Clark, G.S., and Cheung, S.P. 1980. Rubidium - strontium ages from the Oxford Lake - Knee Lake greenstone belt, northern Manitoba; Canadian Journal of Earth Science, v. 17, p.560-568.
- Comba, C.D.A., Binney, W.P., Stewart, R.D., Cunnison, K.M., and Mullen, D.V. 1986. Timmins, Ontario: Exceptional exposures of Archean subaerial and shallow

subaqueous volcanic rocks, and associated ore deposits; GAC-MAC 1986 Field Trip Guidbook #5, 25 p.

- Corfu, F. 1987. Inverse age stratification in the Archean crust of the Superior Province: evidence for infra-and subcrustal accretion from high resolution U-Pb zircon and monazite ages; *Precambrian Research*, v. 36, p.259-275.
- Corfu, F. and Grunsky, E.C. 1987. Igneous and tectonic significance of the Batchewana greenstone belt, Superior Province: a U-Pb zircon and titanite study. *Jour. Geology*, p.87-106.
- Corfu, F., Jackson, S., and Sutcliffe, R., In preparation, U-Pb ages and tectonic significance of Late Archean alkalic magmatism and non-marine sedimentation, Timiskaming Group, Superior Province.
- Corfu, F., Krogh, T.E., Kwok, Y.Y., Marmont, S. and Jensen, L.S. 1989. U-Pb zircon geochronology in the southwestern Abitibi greenstone belt, Superior Province; *Canadian Journal of Earth Sciences*, v. 26, p.1747-1763.
- Corfu, F. and Stott, G.M. 1986. U-Pb ages for late magmatism and regional deformation in the Shebandewan Belt, Superior Province, Canada; *Canadian Journal of Earth Sciences*, v. 23, p.1075-1082.
- Couch, R.W., Whitsett, R.M., and Hachu, B., and Briceno-Guarupe, L. 1981. Structures of the continental margin of Peru and Chile; *in* Nazca Plate: crustal formation of Andean convergence, Kulm. L., ed., *Geological Society of America Memoir* 154, p.703-726.
- Crecraft, H.R., Nash, W.P., and Evans, S.H. 1981. Late Cenozoic volcanism at Twin Peaks, Utah: geology and petrology; *Journal of Geophysical Research*, v. 86, p.10303-10320.
- Daly, R.A., Manger, G.E., and Clark, S.P. 1966. Density of Rocks; Clark, S. P., ed., *in* Handbook of Physical Constants, *Geological Society of America Memoir* 97, p.19-26.

- Davis, D.W., and Edwards, G.R., 1982. zircon U-Pb ages from the Kakagi Lake area, Wabigoon Subprovince northwest Ontario, Canadian Journal of Earth Science, v.19, p.1235-1245.
- Davis, D.W., and Edwards, G.R., 1986. Crustal evolution of Archean rocks in the Kakagi Lake area, Wabigoon Subprovince, Ontario, as interpreted from high precision U-Pb geochronology. Canadian Journal of Earth Science, v.23, p.182-192
- Davis, D.W. Krogh, T.E., Hinzer, J. and Nakamura, E. 1985. Zircon dating of polycyclic volcanism at Sturgeon Lake and implications for base metal mineralization. Economic Geology, v.80, p.1942-1952.
- Davis, D.W., Poulsen, K.H. and Kamo, S.L. 1989. New insights into Archean crustal development from geochronology in the Rainy Lake area, Superior Province, Canada; Journal of Geology, v. 97, p.379-398.
- Davis, D.W., Sutcliffe, R.H., Trowell, N.F. 1988. Geochronological constraints on the tectonic setting of a late Archean greenstone belt; Precambrian Research, v. 39, p.171-191.
- Davis, D.W. and Trowell, N.F. 1982. U-Pb zircon ages from the eastern Savant Lake - Crow Lake metavolcanic-metasedimentary belt, northwest Ontario; Canadian Journal of Earth Sciences, v. 19, p.868-877.
- Delaney, P.T. 1987. Heat transfer during emplacement and cooling of mafic dikes; *in* Mafic dike swarms, Halls, H.C., and Fahrig, W.F., eds., Geological Association of Canada Special Paper 34, p.31-46.
- DePaolo, D.J. 1981a. Trace element and isotopic effects of combined wallrock assimilation and fractional crystallization; Earth and Planetary Science Letters, v. 53, p.189-202.
- 1981b. A neodymium and strontium isotopic study of the Mesozoic calc-alkaline granitic batholiths of the Sierra Nevada and Peninsular Ranges, California; Journal of Geophysical Research, v. 86, p.10470-10488.
- 1985. Isotopic studies of processes in mafic magma chambers: I. The Kiglapait Intrusion, Labrador; Journal of Petrology, v. 26, p.925-951.

- Dimroth, E., Imreh, L., Rocheleau, M., and Goulet, N. 1982. Evolution of the south-central part of the Archean Abitibi Belt, Quebec. Part I: Stratigraphy and paleogeographic model; *Canadian Journal of Earth Sciences*, v. 19, p.1729-1758.
- Dimroth, E., Imreh, L., Goulet, N. and Rocheleau, M. 1983a. Evolution of the south-central part of the Archean Abitibi Belt, Quebec. Part II: Tectonic evolution and geomechanical model; *Canadian Journal of Earth Sciences*, v. 20, p.1355-1373.
- 1983b. Evolution of the south-central part of the Archean Abitibi Belt, Quebec. Part III: Plutonic and metamorphic evolution and geotectonic model; *Canadian Journal of Earth Sciences*, v. 20, p.1374-1388.
- Dimroth, E., Mueller, W., Heidelberg, R., Brisson, H., Poitras, A. and Rocheleau, M. 1986. Diapirism during regional compression: the structural pattern in the Chibougamau region of the Archean Abitibi belt, Quebec; *Geologische Rundschau*, v. 75, p.715-736.
- Dixon, J.M. 1975. Finite strain and progressive deformation in models of diapiric structures; *Tectonophysics*, v. 28, p.89-124.
- Doe, B.R., Leeman, W.P., Christiansen, R.L., and Hedge, C.E. 1982. Lead and strontium isotopes and related trace elements and genetic tracers in the upper Cenozoic rhyolite - basalt association of the Yellowstone Volcanic Field; *Journal of Geophysical Research*, v. 4785-4806.
- Dupre, B., Chauvel, C., and Arndt, N.T. 1984. Pb and Nd isotopic study of two Archean komatiitic flows from Alexo, Ontario; *Geochimica et Cosmochimica Acta*, v. 48, p.1965-1972.
- Eby, G.N. 1975. Abundance and distribution of the rare-earth elements and yttrium in the rocks and minerals of the Oka carbonitite complex, Quebec; *Geochimica et Cosmochimica Acta*, v. 39, p.597-620.

- Fahrig, W.F., 1987. The tectonic settings of continental mafic dike swarms: *in* Mafic dike swarms, Hall, H.C. and Fahrig W.F., eds. Geological Association of Canada Special Paper 34 P.331-348.
- Ferguson, S.A. 1957a. Bristol Township, District of Cochrane; Ontario Department of Mines map 1957-7.
- 1957b. Carscallen Township, District of Cochrane; Ontario Department of Mines map P.23.
- Flinn, D. 1965. On the symmetry principle and the deformation ellipsoid; *Geological Magazine*, v. 102, p.36-45.
- Frarey, M.J., and Krogh, T.E. 1986. U-Pb zircon ages of late internal plutons of the Abitibi and eastern Wawa subprovince, Ontario and Quebec; *in* Current Research, Geol. Survey of Canada, Paper 86-1A, p.43-48.
- Frey, F.A., Bryan, W.B., and Thompson, G. 1974. Atlantic ocean floor: geochemistry and petrology of basalts from legs 2 and 3 of the Deep Sea Drilling Project; *Journal of Geophysical Research*, v. 79, p.5507-5527.
- Frith, R.A., and Doig, R. 1975. Pre-Kenoran tonalitic gneisses in the Grenville province; *Canadian Journal of Earth Science*, v. 12, p.844-849.
- Fyon, J.A., Troop, D.G., Marmont, S., and Macdonald, A.J. 1989. Introduction of gold into Archean crust, Superior Province, Ontario- Coupling between mantle-initiated magmatism and lower crustal thermal maturation; *in* Economic Geology Monograph 6, Keays, R.R., Ramsay, R.H., and Groves, D.I., eds., p.479-491.
- Gariépy, C., Allegre, C.J. 1985. The lead isotope geochemistry and geochronology of late-kinematic intrusives from the Abitibi greenstone belt, and the implications for late crustal evolution; *Geochimica et Cosmochimica Acta*, v. 49, p.2371-2383.
- Gariépy, C., Allegre, C.J., and LaJoie, J. 1984. U-Pb systematics in single zircons from the Pontiac sediments, Abitibi greenstone belt; *Canadian Journal of Earth Science*, v. 21, p.1296-1304.

- Gibson, H.L., Watkinson, D.H., and Comba, C.D.A. 1989. Subaqueous phreatomagmatic explosion breccias at Buttercup Hill, Noranda, Quebec; *Canadian Journal of Earth Sciences*, v. 26, p.1428-1439.
- Goldie, R. 1978. Magma mixing in the Flavrian pluton, Noranda area, Quebec; *Canadian Journal of Earth Sciences*, v. 15, p.132-144.
- Graham, A.R. 1931. Groundhog - Kamiskotia area, District of Cochrane; Ontario Department of Mines Geological report, v. 40, p.23-36. .
- Green, D.H. 1970. The origin of basaltic and nephelinitic magmas; *Transactions of the Leicester Literature and Philosophical Society*, v. 64, p.28-54.
- Gromet, L.P., and Silver, L.T. 1987. REE variations across the Peninsular Ranges batholith: implications for batholithic petrogenesis and crustal growth in magmatic arcs; *Journal of Petrology*, v. 28, p.75-125.
- Grove, T., and Donnelley-Nolan 1986. The evolution of young silicic lavas at Medicine Lake volcano, California: Implications for the origin of compositional gaps in calc-alkaline series lavas; *Contributions to Mineralogy and Petrology*, v. 92, p.281-302.
- Guha, J., Dube, B., Pilote, P., Chown, E., Archambault, G. and Bouchard, G. 1988. Gold mineralization patterns in relation to the lithologic and tectonic evolution of the Chibougamau mining district, Quebec, Canada; *Mineralium Deposita*, v. 23, p.293-299.
- Halliday, A.N., Mahood, G.A., Holden, P., Metz, J.M., Dempster, T.J., Davidson, J.P., 1989. Evidence for long residence times of rhyolitic magma in the Long Valley magmatic system: the isotopic record in precalders lavas of Glass Mountain, Earth and Planetary Science Letters, v.94, p.274-290.
- Hanes, J.A., Queen, M., and Archibald, D.A. 1988. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of dikes and their contact aureoles in the Kapuskasing uplift; *in* Project Lithoprobe Kapuskasing Structural Zone Workshop II, p.165-172.

- Hanson, G.N., Goldich, S.S., Arth, J.G., and Yardley, D.H. 1971. Age of the Early Precambrian rocks of the Saganaga Lake - Northern Light Lake area, Ontario - Minnesota; *Canadian Journal of Earth Science*, v. 8, p.1110-1124.
- Hart, T.R. 1984. The geochemistry and petrogenesis of a metavolcanic and intrusive sequence in the Kamiskotia area; Timmins, Ontario; Unpublished M.Sc. thesis, University of Toronto, 179 p.
- Heaman, L.M. 1989. U-Pb dating of mafic dike swarms: what are the options? (abstract); International Association of Volcanology and Chemistry of the earth's interior, Santa Fe, New Mexico.
- Hildreth, W. 1981. Gradients in silicic magma chambers: implications for lithospheric magmatism; *Journal of Geophysical Research*, v. 86, p.10153-10192.
- Helz, R.T., 1985. Compositions of fine-grained mafic rocks from sills and dikes associated with the Stillwater Complex; Montana Bureau of Mines and Geology Special Publication 92, p.97-118.
- Hoffman, E.L. Naldrett, A.J., Alcock, R.A. and Hancock, R.G.V., 1979. The Noble-Metal Content of Ore in the Levack West and Little Stobie Mines, Ontario. *Canadian Minerologist*, v.17, p.437-451.
- Hofmann, A.W. 1988. Chemical differentiation of the Earth: the relationship between mantle, continental crust and oceanic crust; *Earth and Planetary Science Letters*, v. 90, p.297-314.
- Hogg, N. 1954. Geology of Godfrey Township; Ontario Department of Mines, v. 63, 55 p.
- Hubert, C., Trudel, P. and Gelinas, L. 1984. Archean wrench fault tectonics and structural evolution of the Blake River Group, Abitibi Belt, Quebec; *Canadian Journal of Earth Sciences*, v. 21, p.1024-1032.
- Hudleston, P.J., Schultz-Ela, D. and Southwick, D.L. 1988. Transpression in an Archean greenstone belt, northern Minnesota; *Canadian Journal of Earth Sciences*, v. 25, p.1060-1068.

- Hunter, R.H., and Sparks, R.S.J. 1987. The differentiation of the Skaergaard Intrusion; *Contributions to Mineralogy and Petrology*, v. 95, p 451-461.
- Hyodo, H., in prep., Numerical modeling for Argon diffusion by conduction and convection in crystalline rocks: Ph.D. thesis, University of Toronto.
- Irvine, T.N. 1982. Terminology for layered intrusions; *Journal of Petrology*, v. 127-162.
- Jackson, S.L., and Sutcliffe, R.H., in press, Central Superior Province geology: evidence for allochthonous, ensimatic, southern Abitibi greenstone belt; *Canadian Journal of Earth Sciences*.
- Jacques, A.L., and Green, D.H. 1980. Anhydrous melting of peridotite at 0-15 Kb pressure and the genesis of tholeiitic basalts; *Contributions to Mineralogy and Petrology*, v. 73, p.287-310.
- Jambon, A. 1982. Tracer diffusion in granitic melts: Experimental results for Na, K, Rb, Cs, Ca, Sr, Ba, Ce, Eu to 1300°C and a model of calculation; *Journal of Geophysical Research*, v. 87, p.10797-10810.
- Jensen, L.S., and Pyke, D.R. 1982. Komatiites in the Ontario portion of the Abitibi Belt; *in* Komatiites, Arndt, N.T., and Nisbet, E.G., eds., George Allan and Unwin, London, p.147-157.
- Juster, T.C., Grove, T.L., and Perfit, M.R. 1989. Experimental constraints on the generation of FeTi basalts, andesites, and rhyodacites at the Galapagos Spreading Center, 85°W and 95°W; *Journal of Geophysical Research*, v. 94, p.9251-9274.
- Kerr, H.L. 1906. Exploration in the Mattagami Valley; *in* Report of the Bureau of Mines, Ontario, p.116-135.
- Kerrick, R., Fryer, B.J., King, R.W., Willmore, L.M., and van Hees, E. 1987. Crustal outgassing and LILE enrichment in major lithosphere structures, Archean Abitibi greenstone belt: evidence on the source reservoir from strontium and carbon isotope tracers; *Contributions to Mineralogy and Petrology*, v. 97, p.156-168.

- King, R.W., and Kerrich, R. 1987. Fluorapatite fenitization and gold enrichment in sheeted trondhjemites within the Destor-Porcupine fault Zone, Taylor Township, Ontario; Canadian Journal of Earth Sciences, v. 24, p.479-502.
- Krogh, T.E. and Turek, A., 1981. U-Pb zircon ages from the Gamitagama greenstone belt, southern Superior Province. Canadian Journal of Earth Science, v.19, p.859-867.
- Krogh, T.E., Heaman, L.M., and Machado, N. 1988. Detailed U-Pb chronology of successive stages of zircon growth at medium and deep levels using parts of single zircon and titanite grains; *in* Project Lithoprobe, Kapuskasing Structural Zone transect Workshop, p.243.
- Leahy, E.J. 1968. Whitesides Township, District of Cochrane; Ontario Department of Mines, Map p.488.
- LeBaron, P. 1985. Summary report on work performed on the Croxall option, Bristol and Thorneloe townships; Noranda Exploration Company Limited assessment file report 15 p.
- Lee, C.A., and Sharpe, M.R. 1979. Spheroidal pyroxenite aggregates in the Bushveld Complex: a special case of silicate liquid immiscibility; Earth and Planetary Science Letters, v. 44, p.295-310.
- Legault, M.H. 1985. The geology and alteration associated with the Genex volcanogenic massive sulphide deposit, Godfrey Township, Timmins, Ontario; unpubl. M.Sc. thesis, Carleton University, 222 p.
- Leshner, C.E. 1986. Effects of silicate liquid composition on mineral-liquid element partitioning from Soret diffusion studies; Journal of Geophysical Research, v. 91, p.6123-6141.
- Leshner, C.M., Goodwin, A.M., Campbell, I.H. and Gorton, M.P. 1986. Trace-element geochemistry of ore-associated and barren, felsic metavolcanic rocks in the Superior Province, Canada; Canadian Journal of Earth Sciences, v. 23, p.222-237.

- Liegeois, J.P., and Black, R. 1987. Alkaline magmatism subsequent to collision in the Pan - African belt of the Adrar des Iforas (Mali); *in* Fitton, J.G., and Upton, B., eds., Alkaline Igneous Rocks; Geological Society Special Publication #30, p.381-401.
- Lindstrom D.J. 1976. Experimental study of the partitioning of the transition metals between clinopyroxene and coexisting silicate liquids; unpublished Ph.D. thesis, University of Oregon, 188 p.
- Longhi, J., Walker, D., and Hays, J.F. 1978. The distribution of Fe and Mg between olivine and lunar basaltic liquids; *Geochimica et Cosmochimica Acta*, v. 42, p.1545-1558.
- Ludden, J.N., Francis, D., and Allard, G.O. 1984. The geochemistry of the volcanic rocks of the Chibougamau region of the Abitibi metavolcanic belt; *in* Chibougamau - Stratigraphy and Mineralization: Guha, J., and Chown, E., eds., Canadian Institute of Mining and Metallurgy Special Volume 34, p.20-34.
- Ludden, J., Gelinas, C., and Trudel, P. 1982. Archean metavolcanics from the Rouyn - Noranda district, Abitibi greenstone belt, Quebec. 2. Mobility of trace elements and petrogenetic constraints; *Canadian Journal of Earth Sciences*, v. 19, p.2276-2287.
- Luhr, J.F., and Carmichael, I.S.E. 1981. The Colima volcanic complex, Mexico: Part II: Late Quaternary cinder cones; *Contributions to Mineralogy and Petrology*, v. 76, p.127-147.
- Maas, R., McCulloch, M.T., Campbell, I.H., and Coad, P.R. 1986. Sm-Nd and Rb-Sr dating of an Archean massive sulfide deposit: Kidd Creek, Ontario; *Geology*, v. 14, p.585-588.
- Macdonald, R., Sparks, R.S.J., Sigurdsson, H., Matthey, D.P., McGarvie, D.W., and Smith, R.L. 1987. The 1875 eruption of Askja volcano, Iceland: combined fractional crystallization and selective contamination in the generation of rhyolitic magma; *Mineralogical Magazine*, v. 51, p.183-202.
- MacGeehan, P.J. and MacLean, W.H. 1980. Tholeiitic basalt - rhyolite magmatism and massive sulphide deposits at Matagami, Quebec; *Nature*, v. 283, p.153-157.

- Machado, N., Brooks, C., and Hart, S.R. 1986. Determination of initial $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ in primary minerals from mafic and ultramafic rocks: experimental procedure and implications for the isotopic characteristics of the Archean mantle under the Abitibi greenstone belt, Canada; *Geochimica et Cosmochimica Acta*, v. 50, p 2334-2348.
- Malpas, J., Foley, S.F., and King, A.F. 1986. Petrology of the alkaline-ultrapotassic lamprophyres of Aillik Bay, Labrador; *Canadian Journal of Earth Sciences*, v. 23, p.1902-1918.
- Marsh, B.D. 1982. On the mechanics of igneous diapirism, stoping and zone melting; *American Journal of Science*, v. 282, p.808-855.
- Martin, H. 1987. Petrogenesis of Archean trondhjemites, tonalites and granodiorites from Eastern Finland: Major and trace element geochemistry; *Journal of Petrology*, v. 28, p.921-953.
- McBirney, A.R. 1985. *Igneous Petrology*; Freeman and Cooper, 600 p.
- McCourt, W.J. 1981. The geochemistry and petrography of the Coastal Batholith of Peru, Lima segment; *Journal of the Geological Society of London*, v. 138, p.407-420.
- McIntyre, G.A., Brooks, A.C., Compston, W., and Turek, A., 1966. The statistical assessment of Rb-Sr isochrons; *Journal of Geophysical Research*, v.71, p.5459-5468.
- Menzies, M.A., Rogers, N., Tindle, A., and Hawkesworth, C.J. 1987. Metasomatic and enrichment processes in lithospheric peridotites, an effect of asthenosphere - lithosphere interaction; *in* *Mantle Metasomatism*, Menzies, M.A., and Hawkesworth, C.J., eds., Academic Press, London, p.313-364.
- Menzies, M., Seyfried, W., and Blanchard, D. 1979. Experimental evidence of rare earth element immobility in greenstones; *Nature*, v. 282, p.398-399.
- Middleton, R.S. 1969. Jamieson Township, District of Cochrane; Ontario Department of Mines, Map P.521.
- 1970. Robb Township, District of Cochrane; Ontario Department of Mines, Map P.598.

- 1971a. Turnbull Township, District of Cochrane; Ontario Department of Mines, Map P.638.
 - 1971b. Godfrey Township, District of Cochrane; Ontario Department of Mines, Map P.639.
 - 1973a. Robb and Jamieson townships, District of Cochrane; Ontario Department of Mines, Map 2255.
 - 1973b. Turnbull and Godfrey townships, District of Cochrane; Ontario Department of Mines, Map 2330.
 - 1973c. Magnetic survey of Robb and Jamieson townships, District of Cochrane; Ontario Department of Mines, Geophysical Report 1, 56 p.
 - 1975. Geology of Turnbull and Godfrey townships, District of Cochrane; Ontario Geological Survey, Open File Report 5518, 267 p.
- Moore, N.D., and Agar, R.A. 1985. Variations along a batholith: the Arequipa segment of the coastal Batholith of Peru; *in* Magmatism at a Plate Edge, the Peruvian Andes: John Wiley and Sons, N.Y., p.108-118.
- Morrison, D.A., Davis, D.W., Wooden, J.L., Bogard, D.D., Maczuga, D.E., Phinney, W.C. and Ashwal, L.D. 1985. Age of the Mulcahy Lake intrusion, northwest Ontario, and implications for the evolution of greenstone - granite terrains; *Earth and Planetary Science Letters*, v. 73, p.306-316.
- Morrison, G.W. 1980. Characteristics and tectonic setting of the shoshonitic rock association; *Lithos*, v. 13, p.97-108.
- Mortensen, J.K. 1987. Preliminary U-Pb zircon ages for volcanic and plutonic rocks of the Noranda - Lac Abitibi area, Abitibi Subprovince, Quebec; Geological Survey of Canada, Paper 87-1A, p.581-590.
- Moyer, T.C., and Esperanca, S. 1989. Geochemical and isotopic complexities in bimodal magma systems: the Kaiser Spring Volcanic Field, Arizona; *Journal of Geophysical Research*, v.94, p.7841-7859.

- Mukasa, S.B. 1986. Zircon U-Pb ages of super-units in the Coastal batholith, Peru: Implications for magmatic and tectonic processes; Geological Society of America Bulletin, v. 97, p.241-254.
- Naldrett, A.J. 1981. Nickel sulfide deposits: classification, composition and genesis; Economic Geology 75th Anniversary Volume, p.628-686.
- Nathan, H.D., and Van Kirk, K. 1978. A model for magmatic crystallization; Journal of Petrology, v. 19, p.66-94.
- North American Commission on Stratigraphic Nomenclature, 1983. North American Stratigraphic Code; American Association of Petroleum Geologists Bulletin, v. 67, p.841-875.
- Nunes, P.D. and Pyke, D.R. 1981. Time-stratigraphic correlation of the Kidd Creek orebody with volcanic rocks south of Timmins, Ontario, as inferred from zircon U-Pb ages; Economic Geology, v. 76, p.944-951.
- O'Driscoll, M. 1988. Rare earths, enter the dragon; Industrial Minerals, v. 254, p.21-55.
- O'Hara, M.J. 1977. Geochemical evolution during fractional crystallization of a periodically refilled magma chamber; Nature, v. 266, p.503-507.
- OGS-MERQ 1983. Lithostratigraphic map of the Abitibi Subprovince; Ontario Geological Survey/Ministre de l'Energie et des Ressources, Quebec, scale 1:500 000.
- Pallister, J.S., and Knight, R.J. 1981. Rare earth element geochemistry of the Samail ophiolite near Ibra, Oman; Journal of Geophysical Research, v. 86, p.2673-2697.
- Paradis, S., Ludden, J., and Gelinas, L. 1988. Evidence for contrasting compositional spectra in comagmatic intrusive and extrusive rocks of the late Archean Blake River Group, Abitibi, Quebec; Canadian Journal of Earth Sciences, v. 25, p.134-144.
- Parks, W.A. 1900. Nivin's base line, 1899; *in* Report of the Bureau of Mines, Ontario, p.125-142.
- Parsons, I. 1979. The Klokken gabbro-syenite complex, south Greenland: cryptic variation and origin of inversely graded layering; Journal of Petrology, v. 20, p.653-694.

- Percival, J.A., Krogh, T.E., 1983. U-Pb zircon geochronology of the Kapuskasing structural zone and vicinity in the Chapleau-Foley area, Ontario: Canadian Journal of Earth Science v.20 p.830-843.
- Percival, J.A., Krogh, T.E., and Peterman, Z.E. 1988. When did the Kapuskasing Zone come up?; *in* Project Lithoprobe Kapuskasing Structural Zone Transect Workshop, p.43-48.
- Phinney, W.C., and Morrison, D.A. 1988. Constraints on the environment of the Matachewan dike intrusion (abstract): International Geological correlation Program 257 Annual Meeting, Lund, Sweden, p.25.
- Pitcher, W.S., Atherton, M.P., Cobbing, E.J., and Beckinsale, R.D. 1985. A model for the Coastal Batholith; *in* Magmatism at a Plate Edge, the Peruvian Andes, Pitcher, W.S., Atherton, M.P., Cobbing, E.J., and Beckinsale, R.D., eds., John Wiley and Sons, New York, p.340-341.
- Platt, R.G., Mitchell, R.H., and Holm, P.M. 1983. Marathon dikes: Rb-Sr and K-Ar geochronology of ultrabasic lamprophyres from the vicinity of McKellar Harbour, northwestern Ontario, Canada; Canadian Journal of Earth Sciences, v. 20, p.961-967.
- Poulsen, K.H. 1984. The geological setting of mineralization in the Mine Centre-Fort Frances Area- District of Rainy River; Ontario Geological Survey Open File Report 5512, 126 p.
- Pyke, D.R. 1982. Geology of the Timmins area, District of Cochrane; Ontario Geological Survey, Report 219, 141 p.
- 1978. Regional geology of the Timmins - Matachewan area, Districts of Cochrane and Timiskaming; *in* Summary of Field Work, Ontario Geological Survey, Miscellaneous Paper 82, p.73-77.
- Pyke, D.R. and Middleton, R.S. 1970. Distribution and characteristics of the sulphide ores of the Timmins area; Ontario Department of Mines, Miscellaneous Paper 41, 24p.

- Regan, P.F. 1985. The early basic intrusions; *in* *Magmatism at a Plate Edge The Peruvian Andes*, Pitcher, W.S., and Atherton, M.P., eds., John Wiley and Sons, N.Y., p.72-89.
- Rock N. M.S. 1986. The nature and origin of ultramafic lamprophyres: alnoites and allied rocks; *Journal of Petrology*, v. 27, p.155-196.
- 1987. The nature and origin of lamprophyres: an overview; *in* *Alkaline Igneous Rocks*, Fitton, J.G., and Upton, B.G., eds., Geological Society of London Special Publication 30, p.191-226.
- Roeder, P.L. 1974. Activity of iron and olivine solubility in basaltic liquids; *Earth and Planetary Science Letters*, v. 23, p.397-410.
- Roeder, P.L., and Emslie, R.F. 1970. Olivine - liquid equilibrium; *Contributions to Mineralogy and Petrology*, v. 29, p.275-289.
- Ryerson, F.J., and Hess, P.C. 1978. Implications of liquid-liquid distribution coefficients to mineral-liquid partitioning; *Geochimica et Cosmochimica Acta*, v. 44, p.611-624.
- Schandl, E.S., and Davis, D.W., 1989. Are sericite/fuchsite and chlorite alterations related to massive sulphide mineralization at the Kidd volcanic centre? Evidence from Pb-Pb and U-Pb systematics of hydrothermal rutile: GAC-MAC Program with Abstracts v.14, p.A120.
- Schilling, J.G., Anderson, R.N., and Vogt, P. 1976. Rare earth, Fe and Ti variations along the Galapagos spreading centre, and their relationship to the Galapagos mantle plume; *Nature*, v. 261, p.108-112.
- Schnetzler, C.C., and Philpotts, J.A. 1970. Partition coefficients of rare earth elements between igneous matrix material and rock-forming mineral phenocrysts- II; *Geochimica et Cosmochimica Acta*, v. 34, p.331-340.
- Schwerdtner, W.M., Stott, G.M. and Sutcliffe, R.H. 1983. Strain patterns of crescentic granitoid plutons in the Archean greenstone terrain of Ontario; *Journal of Structural Geology*, v. 5, p.419-430.

- Schwerdtner, W.M., Sutcliffe, R.H., and Troeng, B. 1978. Patterns of total strain in the cretal region of immature diapirs: *Canadian Journal of Earth Sciences* v.15 p.1437-1447.
- Sharpe, J.I. 1968. Field relations of the Matagami area; Quebec Department of Natural Resources, Geological Report #137, 50 p.
- Sharpe, M.R. 1986. Bushveld Complex - Excursion Guidebook; Institute for Geological Research on the Bushveld Complex, Pretoria, 143 p.
- Sharpe, M.R. and Hulbert, L. 1985. Ultramafic sills beneath the eastern Bushveld Complex: mobilized suspensions of early lower zone comulates in a parental magma with boninitic affinities; *Economic Geology*, v. 80, p.849-871.
- Shaw, D.M., Cramer, J.J., Higgins, M.D., and Truscott, M.G. 1986. Composition of the Canadian Precambrian Shield and the Continental Crust of the Earth; *in* Dawson, J.B., Carswell, D.A., Hall, J., and Wedepohl, K.H., eds., 1986, *The Nature of the Lower Continental Crust: Geological Society Special Publication* 24, p.275-282.
- Shirey, S.B. 1984. The origin of Archean crust in the Rainy Lake area, Ontario; unpubl. Ph.D. thesis, State University of New York at Stonybrook, 393 p.
- Shirey, S.B., and Carlson, R.W. 1988. Pb and Nd isotopic constraints on crustal evolution in the southern Superior Province and inferences for a heterogeneous Archean mantle (abstract); *GSA Abstracts with Programs*, v. 20, p.A137.
- Shirey, S.B., and Hanson, G.N. 1986. Mantle heterogeneity and crustal recycling in Archean granite-greenstone belts: evidence from Nd isotopes and trace elements in the Rainy Lake area, Superior Province, Ontario, Canada; *Geochimica et Cosmochimica Acta*, v. 50, p.2631-2651.
- Sigurdsson, H., and Sparks, R.S.J. 1981. Petrology and rhyolitic and mixed-magma ejecta from the 1875 eruption of Askja, Iceland; *Journal of Petrology*, v. 22, p.41-84.

- Simmons, E.C., Hanson, G.N., and Lumbers, S.B. 1980. Geochemistry of the Shawmere Anorthosite Complex, Kapuskasing Structural Zone, Ontario; *Precambrian Research*, v. 11, p.43-71.
- Smith, P.E., Tatsumoto, M., and Farquhar, R.M. 1987. Zircon Lu-Hf systematics and the evolution of the Archean crust in the southern Superior Province, Canada; *Contributions to Mineralogy and Petrology*: v. 97, p.93-104.
- Somerville, R. 1967. Kam-Kotia mine; Canadian Inst. Min. Met. Guide Book for Centennial Field Excursions Northwestern Quebec and Northern Ontario, p.132-134.
- Sorensen, S.S., and Grossman, J.N. 1989. Enrichment of trace elements in garnet amphibolites from a paleo-subduction zone: Catalina Schist, southern California; *Geochimica et Cosmochimica Acta*, v. 53, p.3155-3177.
- Sparks, R.S.J. 1986. The role of crustal contamination in magma evolution through geological time; *Earth and Planetary Science Letters*, v. 78, p.211-223.
- Spulber, S.D., and Rutherford, M.J., 1982. The Origin of Rhyolite and Plagiogranite in Oceanic Crust: An Experimental Study. *Journal of Petrology*, v.24, pt.1, p.1-25
- Steele, K.F., and Wagner, G.H. 1979. Relationship of the Murfreesboro kimberlite and other igneous rocks of Arkansas, U.S.A.; *in Kimberlites, Diatremes, and Diamonds: Their Geology, Petrology and Geochemistry*: Boyd, F.R., and Meyer, H.A., eds., American Geophysical Union, Washington, p.71-91.
- Stott, G.M. 1985. A structural analysis of the central part of the Archean Shebandawan greenstone belt and a crescent-shaped granitoid pluton, northwestern Ontario; Unpublished Ph.D. thesis, University of Toronto, 284 p.
- Stott, G.M., Sanborn-Barrie, M. and Corfu, F. 1987. Major transpression events recorded across Archean Subprovince boundaries in northwestern Ontario (abstract); Yellowknife '87, Geological Association of Canada, Summer Field Meeting, Program with Abstracts, p.26.

- Streckeisen, A. 1976. To each plutonic rock its proper name, *Earth Science Review*, v.12 p.1-33.
- Sutcliffe, 1988. *Geology of the Fletcher Lake Area, District of Thunder Bay; Ontario Geological Survey Report 251*, 64 p.
- Sutcliffe, R.H., Sweeney, J.M., and Edgar, A.D. 1989. The Lac des Iles Complex, Ontario: petrology and platinum group element mineralization in an Archean mafic intrusion; *Canadian Journal of Earth Sciences*, v. 26, p.1408-1427.
- Taylor, S.R., and McLennan, S.M. 1985. *The Continental Crust: its Composition and Evolution*; Blackwell Scientific Publications, Oxford, 312 p.
- Thurston, P.C., and Fryer, B.J. 1983. The geochemistry of repetitive cyclical volcanism from basalt through rhyolite in the Uchi-Confederation greenstone belt, Canada; *Contributions to Mineralogy and Petrology*, v. 83, p.204-226.
- Treiman, A.H. 1982. The Oka Carbonitite Complex, Quebec: aspects of carbonitite petrogenesis; unpubl. Ph.D.thesis, University of Michigan, 168 p.
- Trueman, D.L., and Bannatyne, B.B., 1982. Chromite reserves and the geology of the Bird River Sill, Manitoba: Manitoba Minerals Resource Division Open File Report no.1 73 p.
- Turek, A., Smith, P.E., and Van Schmus, W.R., 1982. Rb-Sr and U-Pb ages of volcanism and granite emplacement in the Michipicoten belt - Wawa, Ontario: *Canadian Journal of Earth Science* v.19 p.1608-1626.
- Verma, S.P., Schilling, J.G., and Waggoner, D.G. 1983. Neodymium isotopic evidence for Galapagos hotspot-spreading centre system evolution; *Nature*, v. 306, p.654-657.
- Wager, L.R. 1960. The major element variation of the layered series of the Skaergaard intrusion and a re-estimation of the average composition of the hidden layered series and of the successive residual magmas; *Journal of Petrology*, v. 1, p.364-398.
- Wager, L.R. and Brown, G.M. 1968. *Layered Igneous Rocks*; Oliver and Boyd, Edinburgh, 588 p.

- Walawender, M.J. 1976. Petrology and emplacement of the Los Pinos pluton, southern California; *Canadian Journal of Earth Science*, v. 13, p.1288-1320.
- Walawender, M.J., and Smith, T.E. 1980. Geochemical and petrologic evolution of the basic plutons of the Peninsular Ranges batholith, southern California; *Journal of Geology*, v. 88, p.233-242.
- Walker, D., Shibata, T., and DeLong, S.E. 1979. Abyssal tholeiites from the Oceanographer Fracture Zone; *Contributions to Mineralogy and Petrology*, v. 70, p.111-125.
- Watson, E.B. 1976. Two liquid partition coefficients: experimental data and geochemical implications; *Contributions to Mineralogy and Petrology*, v. 56, p.119-134.
- 1982. Basalt contamination by continental crust: some experiments and models; *Contributions to Mineralogy and Petrology*, v. 80, p.73-87.
- Weibe, R.A. 1987. Evidence for stratification of basic, silicic and hybrid magmas in the Newark Island layered intrusion, Nain, Labrador; *Geology*, v. 15, p.349-352.
- Weibe, R.A., and Wild, T. 1983. Fractional crystallization and magma mixing in the Tigalak layered intrusion, the Nain anorthosite complex, Labrador; *Contributions to Mineralogy and Petrology*, v. 84, p.327-344.
- Windley, B.F. 1976. Archean anorthosites: a review with the Fiskensasset Complex, west Greenland as a model for interpretation; *Geological Society of South Africa Special Publication 3*, p.319-322.
- Wolfe, W.J. 1970. Distribution of copper, nickel, cobalt, and sulfur in mafic intrusive rocks of the Kamiskotia - Whitesides area, District of Cochrane; Ontario Department of Mines, Miscellaneous Report 44, 28 p.
- 1971. Kamiskotia - Whitesides area, District of Cochrane; Ontario Department of Mines map P.633.
- Wong, L., Davis, D.W., Hanes, J.A., and Robert, F. 1989. An integrated U-Pb and Ar-Ar geochronological study of the Archean Sigma gold deposit, Val D'Or, Quebec (abstract); *GAC-MAC Program with Abstracts*, v. 14, p.A45.

- Wood, D.A. 1978. Major and trace element variations in the Tertiary lavas of eastern Iceland and their significance with respect to the Iceland geochemical anomaly; *Journal of Petrology*, v. 19, p.393-436.
- Wyman, D.A., and Kerrich, R. 1988. Alkaline magmatism, major structures, and gold deposits: implications for greenstone belt gold metallogeny; *Economic Geology*, v. 83, p.454-461.
- York, D., and Hyodo, H. 1988. Argon kinetics at a dike contact: $^{40}\text{Ar}/^{39}\text{Ar}$ laser step-heating study (abstract); *EOS Transactions, American Geophysical Union*, v. 69, p.518.

INTRUSIVE CONTACT

Turnbull Township tonalite (2707 ~~Ma~~)

- 7 Undifferentiated
- 7a Quartz diorite, tonalite
- 7b Quartz monzodiorite, granodiorite
- 7c Quartz monzonite, granite
- 7d Aplite, pegmatite
- 7e Hybrid rocks, agmatite^f

KAMISKOTIA VOLCANIC COMPLEX^g (Units 5, 6)FELSIC VOLCANIC ROCKS (2705 $\pm$ 2 Ma^d)

- 6 Undifferentiated
- 6a Rhyolite quartz-feldspar crystal tuffs and agglomerates
- 6b Welded quartz-feldspar rhyolite and dacite flows and tuffs
- 6c Spherulitic rhyolite flows
- 6d Tuffaceous volcanoclastic rocks

MAFIC VOLCANIC ROCKS

- 5 Undifferentiated
- 5a Massive and pillowed basalt
- 5b Basalt tuff, breccia
- 5c Plagioclase-porphyrific basalt
- 5d Massive and pillowed basaltic andesite

INTRUSIVE CONTACT

KAMISKOTIA GABBROIC COMPLEX^g (Units 3, 4)MAFIC INTRUSIVE ROCKS (2707 $\pm$ 2 Ma)

- 4 Undifferentiated
- 4a Peridotite
- 4b Troctolite and olivine gabbro
- 4c Magnesian gabbro-norite and gabbro (locally anorthositic)
- 4d Ferroan gabbro-norite and gabbro (locally anorthositic)
- 4e Hornblende gabbro and hornblendite
- 4f Pegmatitic gabbro
- 4g Chilled or agmatitic gabbro and gabbro-norite^f
- 4h Gabbro and gabbroic anorthosite sills
- 4i Amphibolite, pyroxenite

FELSIC INTRUSIVE ROCKS

- 3 Undifferentiated
- 3a Granophyric diorite, quartz diorite, tonalite
- 3b Granophyric quartz monzodiorite, granodiorite, quartz monzonite, granite

TABLE 1. TABLE OF LITHOLOGIC UNITS IN THE KAMISKOTIA AREA.

LEGEND^{a, b}

PHANEROZOIC

CENOZOIC

QUATERNARY

RECENT

Stream, lake and organic bog deposits

PLEISTOCENE

Eskers, drumlinoids, glacial till

UNCONFORMITY

PRECAMBRIAN

PROTEROZOIC

MATACHEWAN MAFIC DIKES (circa 2450 Ma^c)

INTRUSIVE CONTACT

ARCHEAN

BRISTOL TOWNSHIP LAMPROPHYRE SUITE (2687 $\pm$ 3 Ma^d)

11 Undifferentiated

11a Clinopyroxene - biotite -carbonate lamprophyre

11b Garnetite

INTRUSIVE CONTACT

FELSIC INTRUSIVE ROCKS

10 Undifferentiated

10a Quartz diorite, tonalite

10b Quartz monzodiorite, granodiorite

10c Quartz monzonite, granite

10d Aplite, pegmatite

10e Intrusive breccia^e

INTRUSIVE CONTACT

Cote Township tonalite (2694 $\pm$ 4 Ma^d)

9 Undifferentiated

9a Quartz diorite, tonalite

9b Quartz monzodiorite, granodiorite

9c Quartz monzonite, granite

9d Aplite, pegmatite

9e Intrusive breccia^e

INTRUSIVE CONTACT

Groundhog River tonalite (2696 $\pm$ 2 Ma^d)

8 Undifferentiated

8a Quartz diorite, tonalite

8b Quartz monzodiorite, granodiorite

8c Quartz monzonite, granite

8d Aplite, pegmatite

8e Intrusive breccia^e

INTRUSIVE CONTACT

METASEDIMENTARY ROCKS

- 2 Undifferentiated
- 2a Oxide- and sulphide-bearing iron formation
- 2b Metagraywacke
- 2c Argillite
- 2d Metachert

LOWER VOLCANIC SUITE (>2707 Ma)

- 1 Undifferentiated
- 1a Pillowed and massive basalt
- 1b Massive and tuffaceous andesite
- 1c Tuffaceous dacite and rhyolite
- 1d Garnet amphibolite

NOTES

- a) Lithologic names are based on field and petrographic observations and geochemical classification following Streckeisen (1976).
- b) Coding of rock types (e.g., numbers 1-11) is in chronologic order based on U-Pb geochronology and field relationships.
- c) The age is from U-Pb zircon and baddeleyite data from a dike within the map area (L. Heaman, personal communication).
- d) The U-Pb ages are from Barrie (1990) and Barrie and Davis (1990).
- e) Intrusive breccia is composed of a matrix of felsic intrusive material (generally tonalitic or granodioritic) with angular to sub-rounded fragments of gabbro and basalt (and their metamorphosed equivalents) up to three meters in width.
- f) Units 7e and 4g have textures that indicate magma mixing with between Turnbull Township tonalite and Kamiskotia Gabbroic Complex (KGC) liquids prior to crystallization and consolidation.
- g) The Kamiskotia Volcanic Complex (KVC) and the KGC are named formally using the North American Stratigraphic Code (1983) in this report. The KVC is intruded and is underlain by the KGC. The Complexes are interpreted to be cogenetic i.e., the KGC represents a residual magma chamber from which the KVC was derived.

Table 2: SUMMARY OF U-Pb AGES, FROM BARRIE AND DAVIS, 1990

<u>SAMPLE</u>	<u>AGE¹</u>	<u>REMARKS</u>
KGC Kamiskotia Gabbroic Complex	2707 +/-2	
KR Kamiskotia Volcanic Complex Rhyolite	2705 +/-2	
GC Granitoid C	2696 +/-1.5 2692 +/-5	titanite 207Pb/206Pb minimum age.
GB Granitoid B	2694 +/-43 2698 +/-8 2640 +/-25	contains inheritance(?).
KCR Kidd Creek Rhyolite	2717 +/-2	one concordant analysis along with four from sample 77-22 of Nunes and Pyke (1980).
PR Prosser Rhyolite	2716 +/-4	
RR Reid Rhyolite	2705 +/-3	

Samples located in Figure 10, and by Universal Transverse Mercator Grid coordinates in Appendix I. Also given in Appendix I are petrologic descriptions and whole rock geochemistry for each sample. All ages are from zircon populations unless noted otherwise.

1. All ages for upper intercepts unless noted otherwise. 2. Analyses listed in Table 1. All analyses of zircon populations unless noted otherwise. 3. Preferred age.

TABLE 3: SUMMARY OF Sm-Nd AND Rb-Sr AGES, FROM BARRIE (1990)

Rock Unit	System	Age (Ma)	MSWD ²	Initial Ratio
Kamiskotia Gabbroic and Volcanic Complex Rocks and Mineral Separates from a Kamiskotia Ferroan Gabbro				
	Sm-Nd	2710 +/- 30	2.77	0.50925 +/- 4
	Rb-Sr	2450 +/- 35	0.42	0.70085 +/- 4
Granitoid B: Cote Township Tonalite Whole Rock and Mineral Separates				
	Sm-Nd	2615 +/- 15	2.24	0.50940 +/- 2
Granitoid C: Groundhog River Tonalite Whole Rock and Mineral Separates				
	Sm-Nd	2530 +/- 35	14.0	0.50941 +/- 4
	Rb-Sr	2530 +/- 35	8.7	0.70124 +/- 2
Garnetite Dike of Bristol Township Lamprophyre Suite Whole Rock and Mineral Separates				
	Sm-Nd	2500 +/- 160	0.37	0.50959 +/- 7

1. Ages and initial ratios with 2-sigma errors calculated using a McIntyre 1 model least squares cubic regression program (McIntyre et al. 1966), with 2-sigma errors for $^{143}\text{Nd}/^{144}\text{Nd}$ and $^{87}\text{Sr}/^{86}\text{Sr} = 0.000040$, $^{147}\text{Sm}/^{144}\text{Nd} = 0.3\%$, $^{87}\text{Rb}/^{86}\text{Sr} = 2.0\%$. 2. Ages with MSWD values <3 indicate geological scatter inside of analytical error and represent isochrons. See text for discussion.

TABLE 4. GEOCHEMISTRY¹ OF KAMISKOTIA GABBROIC COMPLEX CUMULATES FROM BARRIE *et al.*, SUBMITTED.

	Lower Zone-->				Western Traverse-->		
	----Olivine-bearing cumulates----						
	1	2	3	4	5	6	7
	86-260	84-12	84-18	84-19	86-309	86-310	86-311
SiO ₂	44.3	40.7	40.3	38.1	48.1	43.0	41.6
TiO ₂	0.07	0.05	0.07	0.04	0.14	0.13	0.06
Al ₂ O ₃	7.36	9.89	21.70	15.00	20.30	23.00	21.40
Fe ₂ O ₃	9.67	10.30	6.31	9.06	7.08	4.93	6.65
MnO	0.09	0.14	0.10	0.13	0.16	0.07	0.09
MgO	25.90	24.60	13.20	20.80	8.61	10.80	12.70
CaO	3.29	5.82	11.80	8.26	11.30	11.90	10.90
Na ₂ O	0.09	0.17	0.64	0.19	1.52	1.13	1.01
K ₂ O	0.03	0.01	0.06	0.05	0.33	0.20	0.21
P ₂ O ₅	0.02	0.02	0.02	0.02	0.02	0.02	0.01
LOI	9.39	8.00	5.85	8.39	2.70	5.31	5.00
Total	100.2	99.7	100.1	100.0	100.3	100.5	99.6
ppm							
La	0.37	0.18	0.42	0.29	0.65	0.33	0.31
Ce	...	0.4	0.6	0.4	2.2	0.9	0.8
Nd	0.4	0.4	0.5	...	...	...	...
Sm	0.15	0.11	0.15	0.09	0.33	0.22	0.13
Eu	...	0.04	0.30	0.10	0.23	0.18	0.22
Tb	...	...	...	...	...	...	...
Yb	0.10	0.10	0.16	0.12	0.36	0.20	0.08
Lu	0.02	0.02	0.03	0.02	0.05	0.02	0.01
Zr	2	2	5	...	5	7	2
Y	2	1	...	2	4	3	2
Hf	...	0.1	0.1	0.1	...	0.2	...
Rb	...	...	...	...	6	5	9
Sr	7	6	31	128	136	119	128
Ba	...	...	...	...	74	...	...
Sc	5.6	7.1	4.7	5.0	17.4	7.8	4.5
V	18	15	18	11.5	...	45	17
Cr	105	205	114	115	707	105	31
Ni	1290	2300	400	680	370	540	820
Co	114	97	59	92	40	43	73
Zn	62	40	30	75	68	33	40
Mg' ²	86	85	82	84	73	83	81
Height ³ (m)					300	340	400

¹ Major elements by XRF at X-Ray Assay Laboratories, Toronto; trace elements by XRF and INAA at the University of Toronto: see Barrie (1990) for accuracy and precision of analyses. ² Mg' = molecular percent MgO*100/ MgO+FeO, with FeO/FeO+Fe₂O₃ = 0.85. ³ Height -estimated stratigraphic height from base of Western or Northeastern Traverses (Figure 12); see text for details. ⁴ Samples with TH prefix from Hart (1984).

TABLE 4, CONTINUED.

	8	9	10	11	12	13	14
	84-271a	84-271b	84-271c	86-313	86-315	86-317	86-316
SiO ₂	49.9	49.7	49.8	43.6	42.9	41.8	43.0
TiO ₂	0.18	0.18	0.19	0.08	0.09	0.11	0.09
Al ₂ O ₃	14.90	15.80	14.20	23.60	20.80	17.00	20.30
Fe ₂ O ₃	6.39	6.05	6.62	5.46	6.06	8.24	6.42
MnO	0.12	0.11	0.13	0.08	0.09	0.11	0.09
MgO	10.70	10.10	11.30	9.25	12.90	17.20	14.00
CaO	14.30	14.10	13.70	12.20	11.10	9.08	10.60
Na ₂ O	1.19	1.48	1.15	1.18	1.27	0.87	1.27
K ₂ O	0.09	0.25	0.08	0.03	0.10	0.15	0.11
P ₂ O ₅	0.02	0.02	0.02	0.02	0.02	0.02	0.02
LOI	2.39	2.31	2.31	4.93	5.00	5.23	3.93
Total	100.2	100.1	99.5	100.4	100.3	99.8	99.8
ppm							
La	0.44	0.53	0.40	0.30	0.31	0.32	0.52
Ce	1.3	1.3	1.5	0.9	1.6	1.2	0.8
Nd	0.8	...	0.9	...	...	...	...
Sm	0.44	0.38	0.36	0.19	0.19	0.22	0.21
Eu	0.23	0.23	0.22	0.18	0.13	0.15	0.15
Tb	0.1	0.1	0.1	...	...	...	...
Yb	0.62	0.46	0.47	0.19	0.16	0.18	0.15
Lu	0.09	0.08	0.09	0.03	0.03	0.04	0.03
Zr	10	10	10	3	4	3	8
Y	6	5	6	2	2	4	2
Hf	0.3	0.2	...	...	...	...	0.2
Rb	...	...	...	...	...	3	5
Sr	94	98	86	125	114	55	104
Ba	...	...	...	30	...	37	39
Sc	46.0	40.3	39.6	6.3	8.1	8.3	5.2
V	172	151	174	41	36	40	26
Cr	383	353	381	71	108	125	64
Ni	110	310	100	430	610	380	530
Co	38	35	40	33	49	65	50
Zn	50	10	40	20	59	53	15
Mg'	79	78	79	79	83	83	83
Height (m)	400	401	402	540	660	680	710

TABLE 4, CONTINUED.

	<--Lower Zone		Middle Zone-->				
	15	16	17	18	19	20	21
	84-237	84-226	84-227	84-219	86-295	86-296	86-297
SiO ₂	41.0	48.0	47.8	48.8	49.5	48.9	49.8
TiO ₂	0.09	0.28	0.55	0.43	1.16	1.46	0.26
Al ₂ O ₃	21.00	16.80	15.20	15.25	15.60	16.90	15.40
Fe ₂ O ₃	5.41	8.16	10.90	9.97	11.10	9.21	9.94
MnO	0.08	0.15	0.19	0.18	0.16	0.12	0.18
MgO	12.10	10.30	8.74	8.96	7.37	6.04	8.49
CaO	10.70	8.44	10.10	11.80	8.90	11.50	11.20
Na ₂ O	0.86	3.09	2.44	1.54	2.29	2.51	1.64
K ₂ O	0.16	0.25	0.20	0.30	0.12	0.19	0.19
P ₂ O ₅	0.02	0.02	0.02	0.01	0.04	0.02	0.01
LOI	9.00	4.54	3.00	2.89	3.08	2.93	3.16
Total	100.4	100.0	99.1	100.1	99.3	99.8	100.3
ppm							
La	0.35	0.86	0.98	0.48	0.99	0.62	0.57
Ce	...	2.7	3.0	...	2.5	1.5	1.9
Nd	...	1.2	2.3	0.8	2.4	2.4	1.7
Sm	0.17	0.61	1.04	0.62	0.88	0.82	0.60
Eu	0.12	0.38	0.64	0.39	0.77	0.59	0.51
Tb	...	0.2	0.2	0.2	0.2	0.2	0.2
Yb	0.16	0.70	1.19	0.91	0.83	0.58	0.89
Lu	0.03	0.11	0.17	0.14	0.12	0.09	0.14
Zr	4	14	15	14	10	11	7
Y	2	5	9	8	8	6	8
Hf	0.1	...	0.5	0.2	0.6	0.3	0.3
Rb							
Sr	86	106	137	114	188	288	135
Ba	19	20	30	40	32	41	58
Sc	6.3	48.3	40.4	40.4	24.9	31.8	39.1
V	39	180	236	273	203	255	193
Cr	224	279	165	445	314	210	291
Ni	600	220	220	290	230	72	150
Co	47	60	60	44	35	32	43
Zn	23	50	80	10	29	...	68
Mg'	84	74	64	67	60	60	66
Height (m)	725	1250	2150	2325	2550	2700	2875

TABLE 4, CONTINUED.

	<--Middle Zone <--Western Traverse		Upper Zone--> Northeastern traverse-->				
	22	23	24	25	26	27	28
	86-298	86-300	84-402	84-406	TH ³ -150	TH-172	TH-174
SiO ₂	50.3	50.1	49.6	44.6	48.3	46.8	43.6
TiO ₂	0.21	0.14	1.11	2.29	0.76	0.38	0.53
Al ₂ O ₃	16.50	19.10	14.20	13.70	19.87	22.50	11.40
Fe ₂ O ₃	9.29	8.08	12.80	17.30	9.55	7.64	17.53
MnO	0.18	0.16	0.21	0.21	0.15	0.10	0.21
MgO	8.00	6.14	7.79	6.60	5.33	6.02	14.00
CaO	9.24	10.70	12.10	11.10	10.64	11.00	8.02
Na ₂ O	2.53	2.51	1.90	1.65	2.17	3.17	0.79
K ₂ O	0.41	0.12	0.10	0.11	0.18	0.55	0.05
P ₂ O ₅	0.02	0.02	0.13	0.02	0.06	0.05	0.05
LOI	3.62	3.31	0.39	0.77	3.20	2.87	4.63
	100.3	100.4	100.3	98.4	100.2	101.1	100.8
ppm							
La	0.55	0.63	2.40	0.76	1.30	1.80	1.30
Ce	1.4	1.2	8.1	2.6	4.1	4.6	...
Nd	1.1	...	4.0	2.6	2.4	2.7	2.3
Sm	0.47	...	1.82	1.00	0.74	1.00	0.77
Eu	0.46	0.37	0.78	0.57	0.59	0.79	0.39
Tb	...	0.1	0.4	0.2	0.2	0.3	0.2
Yb	0.71	0.42	1.98	1.25	0.71	0.77	0.90
Lu	0.10	0.06	0.26	0.21	0.15	0.14	0.16
Zr	5	2	53	23	16	17	22
Y	5	3	20	10	7	7	8
Hf	...	...	1.4	0.6	0.5	0.4	0.6
Rb	10	...	...	...	2	14	1
Sr	162	184	104	99	142	205	64
Ba	92	42	300	200	26	148	37
Sc	34.7	18.8	48.8	52.1	24.2	17.3	21.7
V	117	91	318	860	...	...	...
Cr	224	101	307	197	130	...	...
Ni	150	200	170	240	90	...	...
Co	37	37	45	61	46	40	167
Zn	26	65	5	30	...	...	...
Mg'	66	69	58	46	56	64	64
Height (m)	3000	3100	100	275	300	1075	1150

TABLE 4, CONTINUED.

	29 84-415	30 TH-85	31 TH-37	32 TH-133	33 TH-146
SiO ₂	41.9	47.8	50.1	40.7	49.2
TiO ₂	0.78	0.77	1.08	3.20	0.56
Al ₂ O ₃	9.37	15.30	14.60	11.57	17.47
Fe ₂ O ₃	19.50	10.53	12.56	24.49	11.15
MnO	0.28	0.09	0.17	0.22	0.12
MgO	15.20	8.33	6.32	6.25	5.29
CaO	7.84	8.46	10.20	7.58	11.26
Na ₂ O	0.12	2.47	1.78	0.81	2.44
K ₂ O	0.03	0.23	0.63	0.44	0.32
P ₂ O ₅	0.05	0.02	0.17	0.10	0.10
LOI	5.16	6.77	2.75	5.80	2.02
Total	100.2	100.8	100.4	101.2	99.9
ppm					
La	0.91	3.20	6.20	2.30	3.30
Ce	2.2	7.2	18.3	7.0	10.4
Nd	1.3	3.5	11.7	...	5.8
Sm	0.60	1.42	3.10	1.68	2.38
Eu	0.32	0.44	1.03	0.94	0.90
Tb	0.2	0.3	0.7	0.4	0.5
Yb	0.76	1.57	2.82	1.94	2.05
Lu	0.13	0.25	0.45	0.34	0.35
Zr	15	21	110	36	45
Y	6	12	26	18	19
Hf	0.3	0.4	2.2	1.0	1.1
Rb	...	9	14	22	5
Sr	34	155	110	75	116
Ba	...	76	189	127	82
Sc	23.7	28.5	39.2	44.7	35.3
V	219	...	...	...	...
Cr	221	...	...	...	...
Ni	630	105	70	120	55
Co	124	...	51	102	47
Zn	150	...	...	...	...
Mg'	64	64	53	37	52
Height (m)	1275	1975	2300	2375	2500

TABLE 5. GEOCHEMISTRY* OF CHILLED GABBROS AND KAMISKOTIA BASALTS

	34 87-14	35 87-15	36 87-18	36 84-186	37 86-285	Prim. Basalt Average	Evolved Basalt Average
wt. %							
SiO ₂	47.4	46.9	47.3	46.7	49.7	50.1	49.2
TiO ₂	0.92	0.90	0.93	0.88	1.01	1.36	2.47
Al ₂ O ₃	16.00	16.00	15.70	15.60	14.4	13.4	12.2
Fe ₂ O ₃	13.40	13.30	14.10	13.10	13.3	14.0	15.5
MnO	0.21	0.20	0.20	0.19	0.21	0.25	0.24
MgO	8.11	8.09	7.49	7.98	6.89	6.32	4.73
CaO	7.01	9.81	9.64	9.93	8.87	7.10	6.79
Na ₂ O	1.15	1.78	1.78	1.28	1.55	2.56	2.49
K ₂ O	2.36	0.34	0.18	0.33	0.19	0.31	0.74
P ₂ O ₅	0.09	0.09	0.09	0.09	0.09	0.14	0.54
LOI	3.23	2.54	2.70	3.16	3.77	3.96	5.10
	99.9	100.0	100.1	99.2	100.0	99.3	99.3
ppm							
La	5.02	4.83	3.76	3.80	5.51	7.30	14.5
Ce	12.4	12.8	10.6	10.0	11.7	18.4	37.2
Nd	6.6	7.7	6.9	7.4	10.3	11.2	25.1
Sm	2.51	2.63	2.51	2.47	3.19	3.67	7.13
Eu	0.83	0.75	0.82	0.74	0.83	1.09	2.30
Tb	0.6	0.5	0.5	0.5	0.8	0.8	1.5
Yb	2.99	3.08	3.10	3.03	3.61	3.35	5.87
Lu	0.48	0.49	0.47	0.46	0.49	0.54	0.90
Zr	61	62	61	55	69	102	302
Y	23	22	23	22	31	30	60
Hf	1.6	1.9	1.7	1.8	1.8	2.6	3.4
U	0.1	0.2	0.1	0.1	0.2	...	0.4
Th	0.5	0.1	0.1	0.4	0.4	0.6	1.6
Ta	...	0.2	0.2	0.2	0.3	0.2	1.1
Rb	88	11	11	10	...	6	18
Sr	65	136	133	112	188	70	130
Ba	770	110	39	70	34	121	285
Sc	42.5	44.9	44.4	44.0	46.4	40.0	38.0
V	236	237	243	239	344	...	315
Cr	194	200	229	301	230	...	139
Ni	170	170	140	170	80	42	49
Co	54	60	58	60	40	...	43
Zn	68	145	145	55	43	...	140
Mg'	57	58	56	58	54	49	42

* Kamiskotia primitive basalt average of samples TH-2, TH-8, TH-25, and TH-208 from Hart (1984); evolved basalt average of samples TH-225, TH-233, TH-244 from Hart (1984), and 88-17, 88-18, 88-19 from Barrie, (1990). Locations for gabbro chill samples in Appendix II; see Hart (1984) and Barrie (1990) for other sample locations.

TABLE 6: GEOCHEMISTRY OF LOWER VOLCANIC SUITE MAFIC AND INTERMEDIATE VOLCANIC ROCKS*, FROM BARRIE (1990)

	84-48.	84-72	84-86	84-136	84-217	84-265
SiO ₂	48.30	49.00	55.10	59.60	54.20	43.80
TiO ₂	1.16	0.81	0.93	0.56	0.65	0.70
Al ₂ O ₃	13.50	14.70	16.40	12.80	15.80	14.40
Fe ₂ O ₃	15.30	12.70	6.87	12.80	11.80	11.80
MnO	0.23	0.19	0.10	0.60	0.50	0.18
MgO	6.96	6.87	3.42	2.43	5.00	8.06
CaO	9.66	12.70	6.44	6.23	6.94	8.75
Na ₂ O	2.29	1.33	3.65	3.26	3.60	0.88
K ₂ O	1.17	0.32	1.22	0.54	0.26	0.61
P ₂ O ₅	0.08	0.06	0.16	0.09	0.12	0.05
LOI	1.39	1.16	5.35	0.39	1.31	10.70
Total	100.0	99.8	99.6	99.3	100.2	99.9
ppm						
La	2.49	2.48	8.83	8.98	4.88	1.90
Ce	6.1	6.4	18.5	21.2	7.0	6.0
Nd	5.90	4.60	10.80	9.20	4.30	3.60
Sm	2.48	1.97	3.00	2.18	1.40	1.45
Eu	0.69	0.75	0.72	0.72	0.66	0.62
Tb	0.4	0.4	0.4	0.4	0.2	0.3
Yb	2.69	2.20	1.75	1.42	1.13	1.40
Lu	0.41	0.32	0.27	0.22	0.19	0.23
Zr	65	46	111	92	23	40
Hf	1.3	1.4	2.0	2.2	0.4	1.4
Y	24	17	18	14	10	13
Nb	3	2	4	3	1	
Ta	0.2	0.1	0.3	0.4	0.1	0.1
Th	6.6	0.2	0.8	1.4	0.2	0.1
Sc	37.0	48.6	17.9	17.2	23.6	34.7
V		259	209	116	175	24
Sr	132	171	150	92	222	88
Ba	270	90	180	250	200	90
Rb		8	33	1	3	18
Cr	168	326	135	273	265	292
Ni	112	148	98	49	107	195
Cu	48	78	58	14	38	5
Co	40.2	53.8	16.4	16.2	19.4	51.5
Zn	40	110	20	70	32	130
Au (ppb)	6				3	5
Mg'	57	55	53	30	50	61

TABLE 6 CONTINUED.

	86-6	86-7	86-12	86-93	86-306	86-307
SiO ₂	48.50	49.60	50.40	46.60	52.40	53.40
TiO ₂	1.28	1.23	0.97	1.96	0.55	0.25
Al ₂ O ₃	13.30	14.50	13.60	12.50	15.00	15.80
Fe ₂ O ₃	16.10	16.00	14.00	20.10	10.20	9.26
MnO	0.26	0.25	0.20	0.28	0.19	0.15
MgO	5.57	5.14	7.13	5.63	6.48	7.44
CaO	8.78	8.18	9.96	10.30	8.60	9.73
Na ₂ O	3.07	3.44	1.65	1.15	3.54	2.36
K ₂ O	0.49	0.15	0.39	0.35	0.87	0.25
P ₂ O ₅	0.10	0.10	0.07	0.18	0.05	0.02
LOI	1.77	1.23	1.31	1.08	1.62	1.47
Total	99.2	99.8	99.7	100.1	99.5	100.1
ppm						
La	4.07	4.24	2.52	1.24	10.50	6.87
Ce	14.2	7.8	7.8	2.5	27.4	17.4
Nd	9.50	9.10	5.00	1.70	17.90	11.60
Sm	3.06	3.17	1.80	0.58	4.80	3.54
Eu	1.16	0.70	0.73	0.39	1.17	0.72
Tb	0.8	0.5	0.4	0.1	0.6	0.7
Yb	3.24	3.44	1.93	0.62	3.93	3.63
Lu	0.51	0.53	0.29	0.09	0.60	0.52
Zr	80	60	60	124	160	54
Hf	2.7	1.6	1.5	0.2	4.2	1.7
Y	28	19	19	41	36	32
Nb	4	2	2	6	11	3
Ta	0.3	0.2	0.1	0.0	0.7	0.1
Th	0.5	0.3	0.2		0.8	0.2
Sc	47.8	52.5	44.6	24.2	26.8	28.0
V					159	177
Sr	118	405	89	93	134	118
Ba	90	80	62		300	203
Rb	13	47	9		18	
Cr	105	69	148	686	120	143
Ni	80	37	220	170	137	162
Cu						
Co	54.0	36.8	46.4	42.7	38.3	39.1
Zn	150	35	60	110	125	
Au	4	6	8	6		4
Mg'	44	42	53	39	59	64

*See Appendix II for sample locations.

TABLE 7: GEOCHEMISTRY OF REGIONAL GRANOTOID ROCKS*, FROM BARRIE (1990)

	--Granitoid A--		--Granitoid B--		--Granitoid C--	
	84-31	86-267	86-101	84-246	84-43	86-220
wt. %						
SiO ₂	68.1	69.5	73.9	76.8	71.4	71.9
TiO ₂	0.36	0.36	0.35	0.32	0.17	0.15
Al ₂ O ₃	15.5	15.1	12.5	11.6	15.6	15.7
Fe ₂ O ₃	3.74	3.26	3.26	3.01	1.75	0.84
MnO	0.06	0.06	0.06	0.04	0.03	0.02
MgO	1.41	1.02	0.78	0.63	0.70	0.41
CaO	3.10	3.61	2.64	0.74	2.82	2.00
Na ₂ O	4.48	4.56	4.00	3.62	5.25	6.48
K ₂ O	1.95	1.59	1.24	0.82	1.75	1.91
P ₂ O ₅	0.10	0.08	0.07	0.06	0.06	0.06
LOI	1.62	1.08	1.00	1.54	0.39	0.54
Total	100.4	100.2	99.8	99.2	99.9	100.0
ppm						
La	24.80	8.97	13.70	11.50	5.88	3.31
Ce	49.0	19.1	32.7	25.3	10.2	11.8
Nd	18.4	8.4	10.3	12.2	3.6	4.6
Sm	4.49	2.40	2.05	2.05	0.75	1.14
Eu	1.08	0.55	0.72	0.66	0.28	0.43
Tb	1.2	0.8	0.3	0.3		0.1
Yb	2.65	2.09	1.17	1.23	0.25	0.15
Lu	0.46	0.31	0.25	0.19	0.05	0.05
Zr	154	125	183	157	98	91
Hf	5.0	3.0	5.0	3.6	1.9	2.9
Y	17	22	13	11	3	4
Nb	8	7	12	4		
Ta	1.4	0.9	0.8	0.7	0.3	0.2
U	1.1	0.6	0.6	0.3	1.1	0.5
Th	8.8	3.8	6.0	1.4	1.2	1.1
Sr	213	222	160	173	405	811
Ba	1090	365	540	260	620	640
Rb	87	55	56	39	47	40
Cr	22	6	15	70	25	
Ni	28			21		
Co	14	7	6	71	2	3
Zn	32	62	43	26	80	
Sc	9.0	6.2	6.3	5.1	1.7	1.4
V	46		30		9	

TABLE 7 CONTINUED.

	Granitoid B	-----Granitoid D-----		
	84-174	84-130	84-266	84-01
wt. %				
SiO ₂	75.3	66.8	66.1	67.4
TiO ₂	0.11	0.59	0.45	0.36
Al ₂ O ₃	13.5	14.6	15.7	16.1
Fe ₂ O ₃	1.45	4.55	3.76	3.22
MnO	0.03	0.07	0.06	0.04
MgO	0.27	1.45	1.70	1.84
CaO	1.28	3.08	4.00	1.97
Na ₂ O	4.52	4.51	4.86	5.78
K ₂ O	2.34	1.61	1.31	1.10
P ₂ O ₅	0.04	0.15	0.13	0.15
LOI	1.08	1.31	1.62	2.23
TOTAL	99.9	98.7	99.7	100.2

*Sample locations given in Appendix II.

TABLE 8. GEOCHEMISTRY OF BRISTOL TOWNSHIP LAMPROPHYRE SUITE AND NEARBY ROCKS.

	87-4	87-6	88-2	88-3	88-4	88-4D	88-5	88-6
wt. %								
SiO ₂	36.10	32.60	38.0	35.0	40.0	40.2	39.8	34.2
TiO ₂	2.27	1.35	1.17	2.38	2.45	2.57	4.85	1.25
Al ₂ O ₃	7.68	4.44	3.88	9.74	8.34	8.40	10.8	6.54
Fe ₂ O ₃	20.30	15.70	14.1	18.1	18.9	18.7	10.1	13.6
MgO	0.46	0.13	11.0	1.66	1.38	1.49	1.79	12.0
MnO	1.78	10.60	0.15	0.43	0.42	0.42	0.16	0.13
CaO	25.70	16.90	20.1	23.7	22.8	22.9	11.7	13.9
Na ₂ O	1.44	0.12	0.19	0.99	1.37	1.32	5.03	0.06
K ₂ O	1.16	2.98	0.64	1.02	1.66	1.70	0.35	4.52
P ₂ O ₅	2.32	2.30	5.81	0.29	0.24	0.26	1.90	2.02
S	0.59	0.04	0.04	0.61	0.42	0.43	0.79	0.07
LOI	1.6	2.3	4.8	3.6	0.7	0.7	7.4	9.7
CO ₂	5.6	4.07	4.73	2.91	0.33	0.32	8.02	8.99
TOTAL*	99.9	99.6	99.9	97.5	98.7	99.1	94.7	98.0
ppm								
La	234	133	-	218	165	155	377	99
Ce	772	332	-	809	664	617	1252	248
Pr	156	47.3	-	155	133	127	244	36
Nd	840	202	-	762	745	753	1272	156
Sm	188	35.8	-	128	187	189	302	29.3
Eu	46.4	9.13	-	27.5	42.9	43.4	64.1	6.8
Gd	110	24.8	-	54.4	107	103	190	18.7
Tb	8.9	2.2	-	4.8	9.6	10.0	19.5	1.7
Dy	38.2	8.6	-	15.5	41.1	40.2	96	7.2
Ho	5.07	1.08	-	2.12	5.8	5.6	16.3	0.9
Er	11	2.3	-	4.0	13	12.1	40	1.8
Tm	1.6	0.3	-	0.7	1.5	1.6	5.3	0.1
Yb	8.3	1.0	-	3.8	10.3	9.5	32	0.9
Lu	1.2	bd1	-	0.7	1.5	1.5	4.9	0.1
Zr	2100	67	-	1200	1586	1599	3070	138
Y	160	26	58 ¹	88	210	204	522	23
Hf	49	4	-	-	-	-	-	-
Nb	18	bd1	-	24	15	9	17	bd1
Ta	3	bd1	-	-	-	-	-	-
Th	44.5	11.2	-	41	35	38	72	12
U	9.5	0.7	-	-	-	-	-	-
Sr	998	1181	1293 ¹	3830	1320	1318	1249	1150
Ba	698	2200	-	-	-	-	-	-
Rb	30	93	-	28	30	27	6	125
Cs	2	13	-	-	-	-	-	-
Cr	bd1	32	-	-	-	-	-	-
Ni	11	77	9	-	-	-	-	-
Cu	190	9.5	11	-	-	-	-	-
Co	39	49	43	-	-	-	-	-
Zn	110	110	102	-	-	-	-	-
Sc	bd1	36	52	-	-	-	-	-
V	56	34	92	-	-	-	-	-
ppb								
Au	2	bd1	-	3	5	5	4	bd1
Pt	-	-	-	bd1	bd1	bd1	bd1	-
Pd	-	-	-	bd1	bd1	bd1	bd1	-

TABLE 8, continued.

	<--Lamprophyre Suite						
	89-5	89-8	89-9	89-10	89-13	89-14	89-15
wt. %							
SiO ₂	37.8	49.6	35.2	39.3	49.6	42.5	37.6
TiO ₂	1.25	0.84	2.2	1.48	0.99	0.82	4.38
Al ₂ O ₃	3.85	5.42	4.76	11.5	14.6	5.66	10.2
Fe ₂ O ₃	14.2	11.8	24.1	12	9.5	10.1	14.2
MgO	10.9	8.79	9.72	4.16	2.19	8.70	2.54
MnO	0.15	0.19	0.24	0.22	0.22	0.23	0.17
CaO	20.0	14.6	13.2	13.2	13.5	23.7	11.2
Na ₂ O	0.23	2.6	0.01	4.77	4.85	0.49	3.98
K ₂ O	0.65	0.20	3.16	2.70	0.86	0.34	0.42
P ₂ O ₅	1.90	2.13	1.07	0.78	1.11	6.06	1.8
S	0.04	0.32	0.66	0.18	0.11	0.05	2.34
LOI	4.8	3.7	6.3	9.4	1.2	1.4	5.4
CO ₂	4.74	3.96	6.04	7.91	0.89	0.86	6.85
TOTAL*	99.7	100.2	100.6	99.7	98.7	100.1	94.2
ppm							
Zr	224	261	254	635	340	218	3081
Y	74	61	45	129	129	79	569
Nb	bd1	bd1	bd1	16	25	bd1	9
Th	29	17	15	54	54	36	86
Sr	1425	566	935	1409	3050	2864	1328
Rb	11	bd1	107	82	20	7	6
Ni	9	49	88	31	23	27	12
Cu	14	91	237	34	51	12	12000
Co	44	42	68	24	25	38	88
Zn	102	98	187	89	78	136	174
Sc	55	22	33	12	9	40	64
V	94	147	333	247	208	175	328

TABLE 8, continued.

	----Altered Basalt----		
	89-7	89-18	89-19
wt. %			
SiO ₂	45.6	53.0	61.9
TiO ₂	1.18	0.73	0.53
Al ₂ O ₃	13.7	19.3	18.5
Fe ₂ O ₃	13.3	7.31	6.41
MgO	5.70	2.62	2.78
MnO	0.21	0.1	0.07
CaO	10.1	3.68	0.93
Na ₂ O	2.49	4.5	3.59
K ₂ O	0.17	0.20	0.78
P ₂ O ₅	0.15	0.14	0.12
S	0.12	0.14	0.07
LOI	7.2	5.2	2.5
CO ₂	5.6	4.07	1.83
TOTAL*	99.9	96.9	98.2
ppm			
Zr	104	140	227
Y	21	25	78
Nb	6	bd1	7
Th	bd1	bd1	30
Sr	358	356	1413
Rb	5	6	14
Ni	289	93	69
Cu	87	41	40
Co	54	23	18
Zn	94	69	74
Sc	25	19	16
V	220	135	130

*using LOI as total volatile content. Samples with low totals contain high concentrations of trace elements and/or halogens.

Geochemistry from OGS analytical laboratories except for samples 87-4 and 87-6 from X-Ray Assay Laboratories, Toronto. Major elements by XRF; REE by ICP; other trace elements by XRF and ICP except Pt, Pd and Au by AA. Sr¹ and Y¹ by ICP, other Sr and Y data from XRF. Zr data by XRF: samples 88-3, 88-4, 88-5, and 89-15 have significantly 15-30 percent higher Zr contents by ICP-MS. bdl- below detection limits. - - not analyzed.

Lamprophyre samples: 87-4: Carbonate-rich biotite lamprophyre; 88-2: Carbonate-apatite rock; 88-3: Garnetite; 88-4: Garnetite (88-4D = duplicate from separate powder); 88-5: Highly altered garnetite(?) with 3% sulfides; 88-6: Biotite lamprophyre; 89-5: Diopside-rich lamprophyre; 89-8: Diopside-rich lamprophyre; 89-9: Biotite- and carbonate-rich lamprophyre; 89-10: Syenite dike with biotite; 89-13: Plagioclase - garnet dike; 89-14: Diopside-rich lamprophyre; 89-15: Highly altered garnetite(?) with 5% sulfides; 89-17: Biotite-carbonate-apatite rock. Other rock types: 89-7: Epidotized basalt; 89-18a: Tourmalinized basalt; 89-19: Silicified and tourmalinized basalt.

Samples from Croxall property except samples 89-5, 89-7, 89-18 and 89-19 from Holmer property: see Figures 32 and 34 for locations.

TABLE 9: S, Ni, Cu, PGE Au and Re ABUNDANCES FOR KGC SULFIDE OCCURRENCES

	84-2a	84-2b	84-2c	84-2f	84-14a	84-63c	84-135	84-175	86-27	86-27a	86-27b	86-321	86-322	86-324	86-326	86-327	86-328	86-329
wt. %																		
S			12.9	2.04	14.1	1.53	22.0	1.32	17.1	15.1	3.52	2.87	6.56	8.03	1.75	7.10	3.69	
ppm																		
Ni			3500	3200	91	240	180	130	1300	1200	1000	910	2600	3700	850	3300	830	
Cu			2000	2000	2700	280	180	33	1200	580	6300	2200	20000	12000	910	7000	2500	
ppb																		
Os	<3	3	3	0.7	<3	<3	<3	<3										
Ir	0.3	2.6	3.6	<0.3	0.6	0.1	0.1	0.1										
Ru	5	<5	10	<5	<5	<5	<5	<5										
Rh	1	4	6	2	2	1	2	1										
Pt	25	45	15	14	76	<5	5	<5										
Pd	5	5	5	<5	55	<5	<5	<5										
Au	51	44	12	5														
Re	<5	<5	15	5	<5	<5	<5	<5										

Analyses by X-Ray Assay Laboratories, Toronto: S, Ni and Cu by XRF, PGE, Au and Re by neutron activation using NI-bead preconcentration: see Hoffmann et al. (1979).

84-2: Occurrence in NW Whitesides Township.

84-14: NW Carscallen Township occurrence (#10 in Figure 4).

84-63: Occurrence in NE Enid Township.

84-135: Occurrence in Western Carscallen Township, adjacent to partially assimilated iron formation.

84-175: Whitesides Bean Lake occurrence (#11 in Figure 4).

84-171a: NW Carscallen township occurrence (#10 in Figure 4).

86-27: Whitesides Bean Lake occurrence (#11 in Figure 4).

86-321 to 329: Western Whitesides Township occurrences (#12 in Figure 4).




Figure 1. Location of the Kamiskotia area in the Superior Province, with major subprovinces in the southern Superior Province labeled.




Figure 2. Distribution of outcrops (Denoted in black) and major esker ridges (denoted by v-arrays) in the Kamiskotia area.

Geology of Jamieson, Godfrey and northern Bristol townships (to east of irregular line) compiled from previously published maps. The townships are: Fo- Fortune; Co- Cote; R- Robb; J- Jamieson; E- Enid; M- Massey; T- Turnbull; G- Godfrey; Fr- Frey; W- Whitesides; Ca- Carscallen; B- Bristol.




Figure 3. Matachewan dikes in western Godfrey Township (after Middleton 1971b, 1973). Dike density typical for Kamiskotia area. North to top of figure.




Figure 4. Geology of the Kamiskotia area. Mines (none currently producing) and mineral occurrences labelled as follows: 1. Kam-Kotia Cu-Zn Mine; 2. Jameland Cu-Zn Mine; 3. Canadian Jamieson Mine; 4. Genex Mine; 5. Lally Gold Mine; 6. Holmer Gold Mine; 7. De Santis gold property; 8. Union Gold Mine; 9. Croxall REE and gold property; 10-12. Ni-Cu occurrences.




Figure 5. Microprobe wavelength dispersive analyses for selected samples. For ferromagnesian minerals, numbers are 100 x molecular $\text{MgO}/\text{MgO} + \text{FeO}$; for plagioclase, numbers are molecular percent anorthite content. Full chemical analyses, sample locations and description of analytical conditions given in Barrie (1990).



Figure 6. Foliation map. Outcrop exposure reflected by density of foliations. Bedrock in the region of granitoid B and the central KGC under glacial cover; bedrock in region of southern KVC covered by poorly drained area. All foliations are near-vertical ($90^\circ \pm 15^\circ$). No preferred dip directions are evident across the region.



Figure 7. Foliation trajectory map. Visual estimate of foliation trends, based on foliations in Figure 6. Light lines represent predominantly non-penetrative foliations due to north-south compression; bold lines represent penetrative planar fabrics due to granitoid emplacement.



Figure 8. Mineral lineation map, with high strain zones marked with hatchured pattern. Note that the western extent of the Destor Porcupine Fault Zone extends to the east and south of Granitoid D; a splay continues to the north within the Lower volcanic suite.




Figure 9. Map of apparent total strain fabric, based on field observations of mineral aggregates. Areas of high strain intensity shaded.




Figure 10. U-Pb sample locations, and general geology of the Kamiskotia - Kidd Creek area. KGC- Kamiskotia Gabbroic Complex; KR- Kamiskotia rhyolite of the Kamiskotia Volcanic Complex; GB- Granitoid B (Cote Township tonalite); GC- Granitoid C (Groundhog River tonalite); GAR- Garnetite of the Bristol Township lamprophyre suite; KCR- Kidd Creek Rhyolite; RR- Reid (Township) rhyolite; PR- Prosser (Township) rhyolite.

Age	Magmatic Events	Deformation Events
2926 (+?) Ma	OLDER SIALIC CRUST (Inherited component in Granitoid C)	
2717 +/- 2 Ma, 2716 +6/-3 Ma	KIDD CREEK RHYOLITE, PROSSER RHYOLITE	
>2707 Ma	LOWER MAFIC VOLCANICS, IRON FORMATION	
2707 +/- 2 Ma, 2705 +/- 2 Ma, 2706 +/- 2 Ma	KAMISKOTIA GABBROIC COMPLEX, KAMISKOTIA RHYOLITE, REID RHYOLITE, GRANITOID A	ONSET OF REGIONAL CRUSTAL FOLDING
2696 +/- 2 Ma	GRANITOID B	B CONTACT STRAIN DEVELOPMENT
2694 +/- 4 Ma	GRANITOID C (zircon)	C CONTACT STRAIN DEVELOPMENT
	GRANITOID D	D CONTACT STRAIN DEVELOPMENT
2692 +/- 5 Ma		(?) THERMAL METAMORPHISM OR PROTRACTED COOLING (titanite, Granitoid C)
		TRANSPRESSION: DEVELOPMENT OF SHEAR ZONES ALONG THE WESTERN DESTOR- PORCUPINE FAULT ZONE
2687 +/- 3 Ma	GARNETITE DIKE	

Figure 11. Summary of the U-Pb ages in the Kamiskotia - Kidd Creek area.




Figure 13. Petrographic traverse for cumulate rocks. ol -olivine; opx -orthopyroxene; cpx -clinopyroxene; plag -plagioclase; hb -hornblende; mt + sulf -magnetite and sulfide; oc - olivine-rich cumulates. Traverse locations shown in Figure 12.




Figure 14. KGC cumulates: whole rock magnesium number, normative anorthite composition, TiO₂ and P₂O₅ versus stratigraphic height.




Figure 15. Ni and Sc versus stratigraphic height in the KGC.




Figure 16. Incompatible trace elements and trace element ratios: La, Yb, La/Yb, Zr, Y, Zr/Y; versus stratigraphic height in the KGC.




Figure 17. Rare earth element profiles, normalized to CI chondrite values of Evensen et al. (1978). Gd values determined from interpolation between Sm and Tb.

a. Olivine-rich cumulates (samples 1-4): horizontal-ruled; LZ cumulates (samples 5-15): northeast-southwest diagonal-ruled; MZ cumulates (samples 16-23): vertical-ruled; chilled samples 33-38 -diagonal-ruled.

b. Lower UZ cumulates (samples 24-29): horizontal ruled; upper LZ cumulates (samples 30-34): vertical-ruled.

c. Range for Kamiskotia basalts: horizontal ruled; range for Kamiskotia evolved basalts: vertical-ruled.




Figure 18. REE and trace element plots for Kamiskotia and Kidd Creek rhyolites, and felsic metavolcanic rocks that are not known to be the host to Cu-Zn deposits (FI and FII) from the southern Superior Province. a. Rare earth element profiles for Kamiskotia and Kidd Creek; b. $(La/Yb)_N$ versus Yb_N ; c. Zr/Y versus Y ; and d. $Ti/100$ - $Zr/10$ - Y ternary. After Lesher et al. 1986.




Figure 19. Rare earth element profiles for selected granitoid rocks from the Kamiskotia area.





Figure 20. X-Y diagrams for Bristol Township lamprophyre suite. Biotite - carbonate - apatite lamprophyres- circles; garnetite samples- hexagons; altered garnetite samples- triangles. a. MgO, Ni, K₂O and P₂O₅ versus log Zr. b. TiO₂, Y, Th, and S versus log Zr.






Figure 21. REE profiles for Bristol Township lamprophyre suite.




Figure 22. CMAS-type tetrahedron projections for KGC chill samples and Kamiskotia basalt (BAS) and evolved basalt (E.BAS), after Walker et al. (1979). a. Projection from plagioclase; b. projection from silica.




Figure 23. Mass balance calculations using the REE. a. Model LZ cumulate compositions from average KGC chill average composition, using percent trapped liquid (C_1) and 5% trapped liquid (C_2). b. Model upper UZ cumulate compositions from average Kamiskotia evolved basalt composition, using 10% trapped liquid (C_1) and 25% trapped liquid (C_2).




Figure 24. X-Y plots for liquid compositions with AFC modeling curves. KGC chill samples: dots; Kamiskotia "primitive" basalts: squares; Kamiskotia evolved basalts: triangles. AFC curves using equations of DePaolo (1981) labeled by ratio of mass assimilated (MA) to mass fractionated (MF), with 0 representing fractionation only. Tick marks represent 10% fractionation increments; double-tick marks are for 80% fractionation. Assimilant used is average of 9 granites and tonalites from the KGC granophyre, with 6 ppm Ni, 0.29 wt.% TiO_2 , 0.04 wt.% P_2O_5 , 9.2 ppm Th, 58 ppm La, 2.9 ppm Eu and 643 ppm Zr (Hart, 1984). Bulk K_D s used: Ni: 2; TiO_2 : 0.2; P_2O_5 : 0.01; Th: 0.001; La: 0.1; Eu: 0.2; Zr: 0.01. For TiO_2 , La and Eu, secondary FC curves from 100 ppm Zr use bulk K_D s as labeled. Additional data from Hart (1984) and Appendix V. a. Ni versus Zr. b. TiO_2 versus Zr. c. P_2O_5 versus Zr. d. Th versus Zr. e. La versus Zr. f. Eu versus Zr.




Figure 25. Primitive mantle-normalized profiles for Kamiskotia chill average in comparison to Alexo, Ontario komatiites (data from Brugmann, 1985; Barnes, 1984; and Whitford and Arndt, 1978), and average N-MORB. KGC chill average for Rb, Ba and K₂O including sample 34 marked with open circles; without sample 34 marked in closed circles. Normalizing values and average N-MORB as listed in Hofmann (1988).





Figure 26. Geology of the number six mine level at Kam-Kotia Porcupine Mines Limited. From Pyke and Middleton (1971).

Figure 27. Geology of the Canadian Jamieson mine property, slightly modified after Comba et al. (1986).



F27





Figure 28. Geology of the Genex Mine area. Slightly modified after Legault (1985).



38m Level Plan

Figure 29. Geology of the Genex Mine, 38 m level. Slightly modified after Legault (1985).




Figure 30. Section of the H and A zones looking northeast. Slightly modified after Legault (1985).




Figure 31. Section of the C zone, view looking north. Slightly modified after Legault (1985).




Figure 32. Geology of the Holmer gold property main outcrops. Quartz veins marked in black.




Figure 33. Geology of the DeSantis gold property outcrops. Quartz vein marked in black.




Figure 35. Histogram of high precision, U-Pb ages for the southern Superior Province (excluding well-defined basement terrane >2780 Ma, and samples with complex U-Pb data that do not clearly define primary igneous ages). The majority of these ages have 2-sigma errors of ± 2 to 4 Ma. The $^{207}\text{Pb}/^{206}\text{Pb}$ ages for the detrital sedimentary grains generally have significantly larger errors: see references for details. Gneissic: predominantly calc-alkalic intrusive rocks at amphibolite or granulite facies in the Kapuskasing (or related) terrane; LILE-/LREE-enriched: intrusive and extrusive rocks, including one alkalic metavolcanic rock, an albitite dike, lamprophyres, and monzodiorite-syenite-granodiorite intrusions with alkalic affinity (e.g., Otto stock, Kirkland Lake area); Tholeiitic, Ultramafic: supracrustal tholeiitic intrusions and related metavolcanic rocks, and one ultramafic dike; Calc-alkalic, Intrusive: tonalite-trondjemite and diorite-granodiorite-granite intrusions, quartz-feldspar porphyries associated with gold mineralization, and sedimentary conglomerate clasts of these compositions; Calc-alkalic, Extrusive: rhyolites and dacites; Sedimentary Detrital:

$^{207}\text{Pb}/^{206}\text{Pb}$ ages for single zircon grains from metagraywackes and meta-arkoses. See text for discussion.

Data from: Barrie and Davis (1990); Barrie et al. (in prep.), Cattell et al. (1984); Corfu, (1987); Corfu and Grunsky, (1986); Corfu and Stott (1986); Corfu et al. (in press); Davis et al. (1985; 1989); Davis and Edwards (1982; 1986), Davis and Trowell (1982); Frarey and Krogh (1986); Garipey et al. (1984); Krogh and Turek (1981); Mortensen, (1987; and unpublished data); Nunes and Pyke (1981); this study; Turek et al. (1982); and Percival and Krogh (1983).





Figure 36. Tectonic model for the Kamiskotia area and parts of the southwestern Abitibi Subprovince. a) Generalized block diagram looking east from Kamiskotia area at 2707 Ma to 2698 Ma. Kamiskotia complex represented in continental marginal basin-like extensional setting. KGC- Kamiskotia gabbroic complex; KLV- Kamiskotia mafic metavolcanic rocks of the KVC; KGV- lower mafic metavolcanic rocks of the KVC; g- Kamiskotia granophyre; LMV- lower mafic metavolcanic rocks of the Lower volcanic suite; shaded- sedimentary rocks; patterned- mafic intrusions; cross pattern- early granitoid intrusions. Dashed line represents division between predominantly supracrustal rocks above and depleted mantle below. Hatchured areas represent areas of partial melting.



b) Same area, at 2690 Ma to 2670 Ma. Symbols same as in a; additionally: B- granitoid B; D- granitoid D, DPFZ- Destor Procupine Fault Zone; KLB- Kirkland - Larder Lake Break; OS- Otto stock (mildly alkalic); Timiskaming- sedimentary rocks and alkalic metavolcanic rocks of the Timiskaming Group; cross pattern; predominantly granitoid intrusions and gneisses; no pattern- predominantly metavolcanic rocks, unsubdivided. After voluminous granitoid emplacement, N-S bulk shortening and subduction, a dextral, transpressive tectonic regime predominates across the region, with development of late ductile shear fabrics along the major fault zones, and wrench-style folding in metavolcanic rocks. From Barrie and Davis (1990).




Plate 1. Textures in the KGC. a. Rhythmic modal layering in Lower Zone cumulates: pyroxene-rich gabbronorite grading up into more plagioclase-rich gabbronorite. Facing direction is up. b. Cross-bedded layering in Lower Zone gabbronorite cumulates. Facing direction is up. c. Hand sample of slightly foliated gabbroic anorthositic cumulate from the Lower Zone. d. Mafic enclaves in anorthositic gabbro, from the Middle Zone.





Plate 2. KGC contact zone textures. a. Thermally-metamorphosed pillowed mafic metavolcanic rocks, Lower Volcanic Suite; northern Whitesides Township. Chlorite alteration rims of pillows are now tremolite-actinolite, and pillow cores are hornfelsic with clinopyroxene porphyroblasts. b. KGC gabbro cumulate cut by mafic pegmatite dike with coxcomb texture; hornblende crystals terminate toward center of dike;

northern Carscallen Township. c. Hornblende-rich bud enclosed by epidote - clay, cutting KGC cumulates 50 m from basal contact, western Carscallen Township. d. "Tennis ball-like" aggregates of plagioclase and altered pyroxene within sub-pegmatitic gabbro, western Whitesides Township.




Plate 3. KGC mixed magma outcrop textures, western Turnbull Township. a. Medium-grained gabbro cut by felsic dikes along brittle fractures. b. Agmatitic textured fine-grained gabbro and felsic intrusion. c-f. Chilled gabbro liquid forming pillow-like structures or irregular-shaped masses due to quenching in contact with cooler felsic liquid.



Plate 4. Photomicrographs of KGC rocks. all photomicrographs 2.2 mm across long dimension. a. Cumulus olivine (center) and intercumulus plagioclase (bottom right) and sulphide (top left) in sulphide-bearing troctolite of the Lower Zone. b. Ferroan gabbro of the Upper Zone, with plagioclase, pigeonite and Ti-rich magnetite. c.

Granophyric quartz-feldspar intergrowth surrounding partly resorbed quartz in KGC granophyre. d. Quench-texture in chill margin of mixed magma outcrop chilled gabbro: radiating clinopyroxene and plagioclase crystals.




Plate 5. Textures in KVC rocks. a. Rhyolite agglomerate, with blocks of flow-banded rhyolite: eastern Godfrey Township. b. Rhyolite crystal tuff with quenched inclusions of andesitic material, providing evidence for coexisting felsic and intermediate liquids in magma chamber prior to eruption: southern Godfrey Township. c. Spherulitic rhyolite,

Kam-Kotia open pit: Robb Township. d. Pillowed mafic matavolcanic rocks, with slight tectonic flattening parallel to hammer handle: Canadian Jamieson mine property, northern Godfrey Township.




Plate 6. Deformation textures. a. Boudinaged blocks of Lower Zone gabbro-norite in migmatitic rocks in western part of contact strain zone of graitoid B, Enid Township. b. Chevron folds in mafic metavolcanic rocks? of the Lower volcanic suite, situated between granitoids B and C, northwest Massey Township. c. Ductile shear zone in pillowed mafic metavolcanic rocks on the northeastern part of the Canadian Jamieson

mine property. Note possible pillow rims near hammer, now highly attenuated. Deformation intensity increases to left. These rocks occur 4 meters away from those in photograph 5d. d. Small zone of ductile deformation in the eastern extent of granitoid D. Undeformed granodiorite is progressively deformed toward quartz-sulphide vein in center, with C fabric parallel to pens.




a.



c.



b.



d.

Plate 7. Croxall property ultramafic lamprophyre suite. a. Undeformed biotite-rich lamprophyre. Very coarse-grained, dark brown to black biotite in matrix of ferroan carbonate, chlorite, magnetite and apatite. b. Contact between biotite ultramafic lamprophyre and melanite-bearing syenite dike. Note alignment of biotite and

magnetite (pseudomorphous after biotite) perpendicular to contact. c. Melanite-bearing syenite dike (to left) and apophyses (to right), cutting carbonate-rich ultramafic lamprophyre. d. Thin melanite-bearing syenite dike cutting medium-grained diopside-feldspar-melanite-biotite rock. Note coxcomb-like growth of garnet toward dike center.


CONVERSION FACTORS FOR MEASUREMENTS IN ONTARIO GEOLOGICAL SURVEY PUBLICATIONS.

Conversion from SI to Imperial			Conversion from Imperial to SI		
SI Unit	Multiplied by	Gives	Imperial Unit	Multiplied by	Gives
LENGTH					
1 mm	0.039 37	inches	1 inch	25.4	mm
1 cm	0.393 70	inches	1 inch	2.54	cm
1 m	3.280 84	feet	1 foot	0.304 8	m
1 m	0.049 709 7	chains	1 chain	20.116 8	m
1 km	0.621 371	miles (statute)	1 mile (statute)	1.609 344	km
AREA					
1 cm ²	0.155 0	square inches	1 square inch	6.451 6	cm ²
1 m ²	10.763 9	square feet	1 square foot	0.092 903 04	m ²
1 km ²	0.386 10	square miles	1 square mile	2.589 988	km ²
1 ha	2.471 054	acres	1 acre	0.404 685 6	ha
VOLUME					
1 cm ³	0.061 02	cubic inches	1 cubic inch	16.387 064	cm ³
1 m ³	35.314 7	cubic feet	1 cubic foot	0.028 316 85	m ³
1 m ³	1.308 0	cubic yards	1 cubic yard	0.764 555	m ³
CAPACITY					
1 L	1.759 755	pints	1 pint	0.568 261	L
1 L	0.879 877	quarts	1 quart	1.136 522	L
1 L	0.219 969	gallons	1 gallon	4.546 090	L
MASS					
1 g	0.035 273 96	ounces (avdp)	1 ounce (avdp)	28.349 523	g
1 g	0.032 150 75	ounces (troy)	1 ounce (troy)	31.103 476 8	g
1 kg	2.204 62	pounds (avdp)	1 pound (avdp)	0.453 592 37	kg
1 kg	0.001 102 3	tons (short)	1 ton (short)	907.184 74	kg
1 t	1.102 311	tons (short)	1 ton (short)	0.907 184 74	t
1 kg	0.000 984 21	tons (long)	1 ton (long)	1016.046 908 8	kg
1 t	0.984 206 5	tons (long)	1 ton (long)	1.016 046 908 8	t
CONCENTRATION					
1 g/t	0.029 166 6	ounce (troy)/ ton (short)	1 ounce (troy)/ ton (short)	34.285 714 2	g/t
1 g/t	0.583 333 33	pennyweights/ ton (short)	1 pennyweight/ ton (short)	1.714 285 7	g/t

OTHER USEFUL CONVERSION FACTORS


	Multiplied by	
1 ounce (troy) per ton (short)	20.0	pennyweights per ton (short)
1 pennyweight per ton (short)	0.05	ounces (troy) per ton (short)

Note: Conversion factors which are in bold type are exact. The conversion factors have been taken from or have been derived from factors given in the Metric Practice Guide for the Canadian Mining and Metallurgical Industries, published by the Mining Association of Canada in co-operation with the Coal Association of Canada.



462000m.E. 81 30'

OFR 5829



Legend for Geology of Kamiskotia area
OFR 5829

LEGEND a,b

PHANEROZOIC
CENOZOIC
QUATERNARY
RECENT
Stream, lake and organic bog deposits
PLEISTOCENE
Bakers, drumlinoids, glacial till
UNCONFORMITY

PRECAMBRIAN
PROTEROZOIC
MATACHEWAN MAFIC DIKES (circa 2450 Ma^c: not shown on map; see figure 2)

INTRUSIVE CONTACT

ARCHAEO
BRISTOL TOWNSHIP LAMPROPHYRE SUITE (2687 ± 3 Ma^d)
11 Undifferentiated
11a Clinopyroxene - biotite lamprophyre
11b Garnetite

INTRUSIVE CONTACT

FELSIC INTRUSIVE ROCKS
10 Undifferentiated
10a Quartz diorite, tonalite
10b Quartz monodiorite, granodiorite
10c Quartz monzonite, granite
10d Aplite, pegmatite
10e Intrusive breccia^e

INTRUSIVE CONTACT
Cote Township tonalite (2694 ± 4 Ma^d)
9 Undifferentiated
9a Quartz diorite, tonalite
9b Quartz monodiorite, granodiorite
9c Quartz monzonite, granite
9d Aplite, pegmatite
9e Intrusive breccia^e

INTRUSIVE CONTACT
Groundhog River tonalite (2696 ± 2 Ma^d)
8 Undifferentiated
8a Quartz diorite, tonalite
8b Quartz monodiorite, granodiorite
8c Quartz monzonite, granite
8d Aplite, pegmatite
8e Intrusive breccia^e

INTRUSIVE CONTACT
Turnbull Township tonalite (2707 Ma)
7 Undifferentiated
7a Quartz diorite, tonalite
7b Quartz monodiorite, granodiorite
7c Quartz monzonite, granite
7d Aplite, pegmatite
7e Hybrid rocks, agmatites^f

KAMISKOTIA VOLCANIC COMPLEX^g (Units 5, 6)
FELSIC VOLCANIC ROCKS (2705 ± 2 Ma^d)
6 Undifferentiated
6a Rhyolite quartz-feldspar crystal tuffs and agglomerates
6b Welded quartz-feldspar rhyolite and dacite flows and tuffs
6c Spherulitic rhyolite flows
6d Tuffaceous volcanoclastic rocks

MAFIC VOLCANIC ROCKS
5 Undifferentiated
5a Massive and pillowed basalt
5b Basalt tuff, breccia
5c Plagioclase-porphritic basalt
5d Massive and pillowed basaltic andesite

INTRUSIVE CONTACT
KAMISKOTIA GABBROIC COMPLEX^g (Units 3, 4)
MAFIC INTRUSIVE ROCKS (2707 ± 2 Ma)
4 Undifferentiated
4a Peridotite
4b Troctolite and olivine gabbro
4c Magnesian gabbro and gabbro (locally anorthositic)
4d Ferroan gabbro and gabbro (locally anorthositic)
4e Hornblende gabbro and hornblende
4f Pegmatitic gabbro
4g Chilled or agmatitic gabbro and gabbroites^f
4h Gabbro and gabbroic anorthositic sills
4i Amphibolite, pyroxenite

FELSIC INTRUSIVE ROCKS
3 Undifferentiated
3a Granophyric diorite, quartz diorite, tonalite
3b Granophyric quartz monodiorite, granodiorite, quartz monzonite, granite

INTRUSIVE CONTACT
METASEDIMENTARY ROCKS
2 Undifferentiated
2a Oxide- and sulphide-bearing iron formation
2b Metagraywacke
2c Argillite
2d Metachert

LOWER VOLCANIC SUITE (>2707 Ma)
1 Undifferentiated
1a Pillowed and massive basalt
1b Massive and tuffaceous andesite
1c Tuffaceous dacite and rhyolite
1d Garnet amphibolite

NOTES
a) Lithologic names are based on field and petrographic observations and geochemical classification following Streckeisen (1976).
b) Coding of rock types (e.g., numbers 1-11) is in chronologic order based on U-Pb geochronology and field relationships. Rock types noted by lighter type are subordinate to those in bold type, and generally as breccia or agmatitic blocks within the other rock types.
c) The age is from U-Pb zircon and baddeleyite data from a dike within the map area (L. Heaman, personal communication).
d) The U-Pb ages are from Barrie (1990) and Barrie and Davis (1990).
e) Intrusive breccia is composed of a matrix of felsic intrusive material (generally tonalitic or granodioritic) with angular to sub-rounded fragments of gabbro and basalt (and their metamorphosed equivalents) up to three meters in width.
f) Units 7e and 4g have textures that indicate magma mixing with between Turnbull Township tonalite and Kamiskotia gabbro liquids prior to crystallization and consolidation.
g) The Kamiskotia Gabbroic Complex (KGC) and Kamiskotia Volcanic Complex (KVC) are named formally using the North American Stratigraphic Code in the accompanying geologic report. The KGC both intrudes and is overlain by the KVC. The complexes are interpreted to be genetically related, i.e., the KGC represents a residual magma chamber from which the KVC was derived.